



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 05:06 AM GMT

PDB ID : 2EVS  
Title : Crystal structure of human Glycolipid Transfer Protein complexed with n-hexyl-beta-D-glucoside  
Authors : Malinina, L.; Malakhova, M.L.; Kanack, A.T.; Abagyan, R.; Brown, R.E.; Patel, D.J.  
Deposited on : 2005-10-31  
Resolution : 2.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2938 (2.20-2.20)
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	209	
1	E	209	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	HEX	A	301	-	X
4	D10	A	302	-	X

## 2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 3496 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycolipid transfer protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	0	3	0
			1650	1074	270	297	9			
1	E	188	Total	C	N	O	S	0	3	0
			1525	995	245	275	10			

- Molecule 2 is n-hexyl-beta-D-glucoside (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	E	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is HEXANE (three-letter code: HEX) (formula:  $C_6H_{14}$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 6 6	0	0
3	A	1	Total C 6 6	0	0
3	E	1	Total C 6 6	0	0

- Molecule 4 is DECANE (three-letter code: D10) (formula: C<sub>10</sub>H<sub>22</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C 10 10	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	227	Total 227	O 227	0	0
5	E	42	Total 42	O 42	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.91Å 42.18Å 70.82Å 90.00° 104.57° 90.00°	Depositor
Resolution (Å)	15.00 – 2.20 17.14 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.3 (15.00-2.20) 98.2 (17.14-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.21Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.190 , 0.236 0.200 , 0.192	Depositor DCC
$R_{free}$ test set	1183 reflections (5.44%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.3	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 72.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22940 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3496	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D10, GLC, HEX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	3/1701 (0.2%)	0.79	1/2305 (0.0%)
1	E	0.51	0/1577	0.58	0/2135
All	All	0.68	3/3278 (0.1%)	0.70	1/4440 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	174	GLU	CG-CD	6.41	1.61	1.51
1	A	174	GLU	CB-CG	6.08	1.63	1.52
1	A	196	GLU	CG-CD	5.53	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	ARG	NE-CZ-NH2	-6.71	116.95	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1667	38	2
1	E	1525	0	1539	23	0
2	A	12	0	11	0	0
2	E	12	0	11	2	0
3	A	12	0	24	3	0
3	E	6	0	14	2	0
4	A	10	0	22	0	0
5	A	227	0	0	15	3
5	E	42	0	0	1	0
All	All	3496	0	3288	59	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (59) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:57:LYS:HE3	5:A:533:HOH:O	1.29	1.26
1:A:71:GLN:NE2	1:A:201[B]:MET:SD	2.53	0.80
1:A:209:VAL:HG21	3:A:301:HEX:H12	1.66	0.77
1:A:95:MET:SD	5:A:515:HOH:O	2.44	0.75
1:A:174:GLU:HB3	5:A:534:HOH:O	1.89	0.72
1:A:174:GLU:HG3	5:A:344:HOH:O	1.90	0.71
1:A:74:LEU:HB3	1:A:201[B]:MET:HE2	1.73	0.71
1:A:19:GLU:OE1	5:A:503:HOH:O	2.11	0.68
3:A:311:HEX:H12	1:E:209:VAL:HG21	1.76	0.67
1:A:74:LEU:HB2	1:A:201[B]:MET:HE1	1.79	0.64
1:A:71:GLN:NE2	1:A:201[A]:MET:SD	2.73	0.62
1:E:125:ARG:HH21	1:E:153:TYR:HB3	1.66	0.60
1:A:44:PRO:O	3:A:301:HEX:H62	2.04	0.57
1:A:66:LYS:NZ	5:A:400:HOH:O	2.38	0.56
1:A:206:ASN:HB2	5:A:518:HOH:O	2.05	0.55
1:A:124:ILE:HA	1:A:127:ASN:HD22	1.72	0.55
1:A:200:GLN:HG3	5:A:468:HOH:O	2.07	0.54
1:A:128:ALA:HB1	1:A:152:LEU:HD22	1.89	0.54
1:E:41:VAL:O	1:E:42:PHE:HB2	2.08	0.53
1:A:74:LEU:CB	1:A:201[B]:MET:CE	2.86	0.53
1:E:41:VAL:O	1:E:42:PHE:CB	2.55	0.53
1:A:200:GLN:HG2	5:A:509:HOH:O	2.08	0.52
1:A:74:LEU:HB3	1:A:201[B]:MET:CE	2.39	0.52
1:E:124:ILE:HD11	1:E:156:PRO:HD2	1.93	0.50
1:E:18:ILE:HD13	1:E:190:THR:HG23	1.93	0.50
1:A:40:PRO:CB	2:E:310:GLC:H1	2.42	0.50
1:A:43:THR:N	1:A:44:PRO:CD	2.75	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:41:VAL:HG12	1:E:144:VAL:HG22	1.93	0.49
1:E:40:PRO:O	1:E:43:THR:HG23	2.11	0.49
1:A:74:LEU:CB	1:A:201[B]:MET:HE2	2.41	0.49
1:E:196:GLU:O	1:E:200:GLN:HB2	2.13	0.49
1:A:137:LYS:NZ	5:A:389:HOH:O	2.38	0.49
1:A:148:PHE:O	1:A:152:LEU:HG	2.14	0.48
1:E:41:VAL:O	1:E:42:PHE:CD1	2.67	0.47
1:A:74:LEU:HB2	1:A:201[B]:MET:CE	2.45	0.46
1:A:38:GLY:HA3	5:A:525:HOH:O	2.14	0.46
1:E:107:PHE:CD2	3:E:312:HEX:H52	2.51	0.46
1:E:68:ARG:CZ	5:E:337:HOH:O	2.64	0.46
1:A:71:GLN:NE2	1:A:201[B]:MET:CE	2.80	0.45
1:A:41:VAL:HG12	1:E:144:VAL:CG2	2.46	0.45
1:E:85:TRP:CG	1:E:86:PRO:HA	2.52	0.45
1:E:43:THR:N	1:E:44:PRO:CD	2.81	0.44
1:A:57:LYS:NZ	5:A:448:HOH:O	2.49	0.44
1:E:77:GLU:HA	1:E:80:MET:HE2	1.99	0.44
1:E:39:SER:O	1:E:41:VAL:O	2.36	0.43
1:A:145:GLN:O	1:A:149:GLN:CG	2.67	0.43
1:A:196:GLU:HG3	5:A:462:HOH:O	2.18	0.42
1:A:138:LYS:HD3	5:A:438:HOH:O	2.18	0.42
1:A:19:GLU:HG2	5:A:503:HOH:O	2.18	0.42
1:E:63:ASN:HD22	1:E:66:LYS:HE2	1.85	0.42
1:A:40:PRO:HB3	2:E:310:GLC:H1	2.02	0.42
1:E:107:PHE:CE2	3:E:312:HEX:H52	2.55	0.41
1:A:139:TYR:HE1	1:A:206:ASN:HD22	1.68	0.41
1:E:142:TRP:CE2	1:E:143:ILE:HD12	2.55	0.41
1:E:63:ASN:ND2	1:E:66:LYS:HB2	2.36	0.41
1:A:164:ALA:O	1:A:167:LYS:HD2	2.21	0.40
1:E:81:TYR:CD2	1:E:88:VAL:HG11	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:150:ALA:C	5:A:500:HOH:O[1_545]	1.97	0.23
1:A:151:ALA:N	5:A:500:HOH:O[1_545]	1.99	0.21
5:A:398:HOH:O	5:A:455:HOH:O[2_656]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/209 (98%)	199 (97%)	6 (3%)	0	100	100
1	E	187/209 (90%)	175 (94%)	10 (5%)	2 (1%)	21	16
All	All	392/418 (94%)	374 (95%)	16 (4%)	2 (0%)	38	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	42	PHE
1	E	82	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/180 (99%)	167 (94%)	11 (6%)	26	27
1	E	166/180 (92%)	156 (94%)	10 (6%)	27	29
All	All	344/360 (96%)	323 (94%)	21 (6%)	32	28

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	50[A]	SER
1	A	50[B]	SER
1	A	57	LYS
1	A	115	GLU
1	A	130	LYS
1	A	138	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	167	LYS
1	A	171	VAL
1	A	201[A]	MET
1	A	201[B]	MET
1	E	41	VAL
1	E	71	GLN
1	E	84	GLU
1	E	113	ASP
1	E	153	TYR
1	E	174	GLU
1	E	182	LEU
1	E	196	GLU
1	E	201[A]	MET
1	E	201[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	72	ASN
1	A	127	ASN
1	A	170	ASN
1	A	206	ASN
1	E	63	ASN
1	E	127	ASN
1	E	206	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	A	300	3	12,12,12	0.65	0	17,17,17	2.11	5 (29%)
3	HEX	A	301	2	5,5,5	0.35	0	4,4,4	0.29	0
4	D10	A	302	-	9,9,9	0.36	0	8,8,8	0.29	0
3	HEX	A	311	2	5,5,5	0.26	0	4,4,4	0.57	0
2	GLC	E	310	3	12,12,12	0.59	0	17,17,17	2.16	5 (29%)
3	HEX	E	312	-	5,5,5	0.30	0	4,4,4	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	300	3	-	0/2/22/22	0/1/1/1
3	HEX	A	301	2	-	0/3/3/3	0/0/0/0
4	D10	A	302	-	-	0/7/7/7	0/0/0/0
3	HEX	A	311	2	-	0/3/3/3	0/0/0/0
2	GLC	E	310	3	-	0/2/22/22	0/1/1/1
3	HEX	E	312	-	-	0/3/3/3	0/0/0/0

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	GLC	C1-O5-C5	5.19	122.69	113.40
2	E	310	GLC	C1-O5-C5	5.08	122.49	113.40
2	E	310	GLC	O5-C1-C2	5.05	117.69	109.86
2	A	300	GLC	O5-C1-C2	4.78	117.27	109.86
2	E	310	GLC	O5-C5-C4	2.94	115.20	109.76
2	A	300	GLC	O1-C1-O5	-2.38	103.90	110.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	GLC	O5-C5-C4	2.22	113.87	109.76
2	A	300	GLC	C1-C2-C3	2.14	113.93	110.53
2	E	310	GLC	O1-C1-O5	-2.03	104.84	110.32
2	E	310	GLC	C1-C2-C3	2.00	113.70	110.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	205/209 (98%)	0.42	11 (5%) 25 25	27, 35, 53, 62	0
1	E	188/209 (89%)	1.83	70 (37%) 1 0	28, 43, 49, 53	0
All	All	393/418 (94%)	1.09	81 (20%) 1 1	27, 40, 51, 62	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	182	LEU	8.5
1	E	164	ALA	6.6
1	E	65	ALA	6.5
1	E	171	VAL	6.1
1	E	200	GLN	5.8
1	E	22	PRO	5.6
1	E	142	TRP	5.4
1	E	172	THR	5.0
1	E	62	THR	4.8
1	E	185	VAL	4.6
1	E	25	GLU	4.6
1	E	202	ASN	4.6
1	A	153	TYR	4.5
1	E	178	GLU	4.4
1	E	60	TYR	4.3
1	E	176	CYS	4.3
1	A	168	GLY	4.2
1	E	38	GLY	4.1
1	E	180	ILE	4.1
1	E	189	ALA	4.1
1	E	68	ARG	4.0
1	E	118	GLU	4.0
1	E	29	HIS	4.0
1	E	80	MET	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	84	GLU	3.9
1	E	165	LEU	3.8
1	E	134[A]	MET	3.7
1	E	66	LYS	3.7
1	E	79	GLU	3.6
1	A	169	GLN	3.6
1	E	83	ALA	3.6
1	E	175	GLU	3.5
1	E	193	VAL	3.4
1	E	19	GLU	3.4
1	E	74	LEU	3.4
1	E	203	ALA	3.3
1	E	75	GLU	3.3
1	A	38	GLY	3.3
1	E	186	ASN	3.3
1	E	196	GLU	3.3
1	E	112	CYS	3.2
1	E	157	TYR	3.1
1	E	67	PHE	3.1
1	E	199	THR	3.1
1	E	206	ASN	3.1
1	E	187	TYR	3.0
1	E	163	LYS	2.9
1	E	64	PRO	2.9
1	A	170	ASN	2.9
1	E	63	ASN	2.9
1	E	36	CYS	2.8
1	E	76	VAL	2.8
1	E	139	TYR	2.8
1	E	205	LEU	2.7
1	E	33	PHE	2.7
1	E	119	ASN	2.7
1	A	174	GLU	2.7
1	E	85	TRP	2.7
1	E	159	SER	2.7
1	E	201[A]	MET	2.7
1	E	190	THR	2.6
1	E	55	LYS	2.6
1	E	40	PRO	2.6
1	E	72	ASN	2.5
1	E	183	PHE	2.4
1	E	153	TYR	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	157	TYR	2.4
1	E	69	THR	2.4
1	E	130	LYS	2.4
1	E	198	TYR	2.3
1	E	47	ALA	2.3
1	A	167	LYS	2.3
1	A	124	ILE	2.3
1	E	21	GLY	2.3
1	E	116	ARG	2.1
1	E	138	LYS	2.1
1	A	165	LEU	2.1
1	E	115	GLU	2.1
1	E	71	GLN	2.1
1	E	113	ASP	2.1
1	A	94	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	D10	A	302	10/10	0.34	4.92	45,46,48,48	0
3	HEX	A	301	6/6	0.31	3.64	51,52,53,53	0
3	HEX	E	312	6/6	0.27	1.11	52,53,53,53	0
3	HEX	A	311	6/6	0.16	0.72	41,42,44,45	0
2	GLC	A	300	12/12	0.16	0.13	40,45,48,49	0
2	GLC	E	310	12/12	0.16	-0.09	34,38,40,40	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.