



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 06:39 AM GMT

PDB ID : 4EYT
Title : Crystal structure of the C-terminal domain of Tetrahymena telomerase protein p65
Authors : Singh, M.; Wang, Z.; Koo, B.-K.; Patel, A.; Cascio, D.; Collins, K.; Feigon, J.
Deposited on : 2012-05-01
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	129	
1	B	129	
1	C	129	
1	D	129	
1	E	129	
1	F	129	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4925 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Telomerase associated protein p65.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	97	Total	C	N	O	S	0	0	0
			801	509	140	148	4			
1	B	99	Total	C	N	O	S	0	0	0
			825	526	144	151	4			
1	C	99	Total	C	N	O	S	0	0	0
			825	525	144	152	4			
1	D	100	Total	C	N	O	S	0	0	0
			833	534	145	150	4			
1	E	98	Total	C	N	O	S	0	0	0
			808	515	142	147	4			
1	F	98	Total	C	N	O	S	0	0	0
			818	521	143	150	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	367	MET	-	EXPRESSION TAG	UNP Q6JXI6
A	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
A	374	SER	-	EXPRESSION TAG	UNP Q6JXI6
B	367	MET	-	EXPRESSION TAG	UNP Q6JXI6
B	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
B	374	SER	-	EXPRESSION TAG	UNP Q6JXI6
C	367	MET	-	EXPRESSION TAG	UNP Q6JXI6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
C	374	SER	-	EXPRESSION TAG	UNP Q6JXI6
D	367	MET	-	EXPRESSION TAG	UNP Q6JXI6
D	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
D	374	SER	-	EXPRESSION TAG	UNP Q6JXI6
E	367	MET	-	EXPRESSION TAG	UNP Q6JXI6
E	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
E	374	SER	-	EXPRESSION TAG	UNP Q6JXI6
F	367	MET	-	EXPRESSION TAG	UNP Q6JXI6
F	368	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	369	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	370	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	371	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	372	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	373	HIS	-	EXPRESSION TAG	UNP Q6JXI6
F	374	SER	-	EXPRESSION TAG	UNP Q6JXI6


- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		
3	B	4	Total	O	0	0
			4	4		
3	D	3	Total	O	0	0
			3	3		



LYS

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.19Å 91.90Å 82.83Å 90.00° 107.36° 90.00°	Depositor
Resolution (Å)	60.55 – 2.50 60.55 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.9 (60.55-2.50) 94.6 (60.55-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.225 , 0.268 0.215 , 0.262	Depositor DCC
R_{free} test set	1845 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	68.4	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 57.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 36716 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4925	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/814	0.57	0/1097
1	B	0.46	0/839	0.61	0/1129
1	C	0.43	0/839	0.56	0/1130
1	D	0.46	0/847	0.62	0/1137
1	E	0.40	0/822	0.57	0/1107
1	F	0.42	0/832	0.59	0/1120
All	All	0.43	0/4993	0.59	0/6720

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	0	1	0
1	B	825	0	0	0	0
1	C	825	0	0	2	0
1	D	833	0	0	2	0
1	E	808	0	0	2	0
1	F	818	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
3	A	3	0	0	0	0
3	B	4	0	0	0	0
3	D	3	0	0	0	0
All	All	4925	0	0	10	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (10) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:F:481:ASN:ND2	1:F:484:ASN:OD1	2.31	0.64
1:E:481:ASN:ND2	1:E:483:ASN:OD1	2.31	0.63
1:D:470:ASP:OD1	1:D:473:ARG:NH2	2.33	0.61
1:C:465:ARG:NH2	1:C:510:TYR:OH	2.36	0.59
1:F:385:ASN:ND2	1:F:497:ASP:OD1	2.43	0.51
1:F:412:ASP:OD1	1:F:460:GLY:N	2.44	0.51
1:D:505:ASP:OD1	1:D:505:ASP:N	2.44	0.49
1:E:505:ASP:OD2	1:E:506:GLU:N	2.47	0.47
1:A:391:LEU:N	1:A:394:GLU:OE2	2.50	0.44
1:C:392:LYS:NZ	1:C:408:CYS:O	2.53	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/129 (74%)	90 (95%)	5 (5%)	0	100	100
1	B	97/129 (75%)	92 (95%)	5 (5%)	0	100	100
1	C	97/129 (75%)	92 (95%)	4 (4%)	1 (1%)	22	38
1	D	96/129 (74%)	94 (98%)	2 (2%)	0	100	100
1	E	96/129 (74%)	86 (90%)	10 (10%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	96/129 (74%)	90 (94%)	6 (6%)	0	100	100
All	All	577/774 (74%)	544 (94%)	32 (6%)	1 (0%)	56	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	411	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/121 (73%)	80 (91%)	8 (9%)	14	24
1	B	90/121 (74%)	84 (93%)	6 (7%)	23	40
1	C	90/121 (74%)	87 (97%)	3 (3%)	50	76
1	D	90/121 (74%)	87 (97%)	3 (3%)	50	76
1	E	87/121 (72%)	83 (95%)	4 (5%)	37	62
1	F	90/121 (74%)	88 (98%)	2 (2%)	64	88
All	All	535/726 (74%)	509 (95%)	26 (5%)	35	59

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	377	GLN
1	A	379	CYS
1	A	392	LYS
1	A	463	MET
1	A	469	SER
1	A	514	ILE
1	A	517	LYS
1	A	518	LYS
1	B	409	ASP
1	B	470	ASP
1	B	485	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	515	LYS
1	B	520	GLU
1	B	523	LYS
1	C	463	MET
1	C	472	GLN
1	C	479	LEU
1	D	410	TYR
1	D	516	PHE
1	D	519	ASN
1	E	378	ASN
1	E	388	GLN
1	E	467	GLN
1	E	483	ASN
1	F	411	ILE
1	F	480	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	601	-	4,4,4	0.07	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	601	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	97/129 (75%)	0.45	1 (1%) 79 81	54, 80, 147, 178	0
1	B	99/129 (76%)	0.50	7 (7%) 16 15	54, 83, 129, 161	0
1	C	99/129 (76%)	0.29	4 (4%) 36 37	53, 79, 153, 181	0
1	D	100/129 (77%)	0.71	13 (13%) 4 4	51, 77, 147, 161	0
1	E	98/129 (75%)	0.53	12 (12%) 5 4	59, 90, 162, 174	0
1	F	98/129 (75%)	0.85	14 (14%) 3 3	59, 101, 148, 191	0
All	All	591/774 (76%)	0.55	51 (8%) 11 10	51, 84, 153, 191	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	522	ARG	6.0
1	F	522	ARG	5.2
1	F	510	TYR	5.0
1	F	514	ILE	4.4
1	D	411	ILE	4.1
1	F	411	ILE	4.1
1	C	519	ASN	4.0
1	F	515	LYS	3.9
1	D	513	TYR	3.9
1	B	518	LYS	3.8
1	F	412	ASP	3.8
1	F	521	PHE	3.8
1	D	517	LYS	3.7
1	E	516	PHE	3.6
1	B	514	ILE	3.6
1	B	511	TRP	3.4
1	D	514	ILE	3.4
1	D	518	LYS	3.4
1	F	517	LYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	516	PHE	3.3
1	A	516	PHE	3.1
1	F	518	LYS	2.9
1	F	511	TRP	2.9
1	B	521	PHE	2.8
1	E	510	TYR	2.8
1	D	511	TRP	2.8
1	E	521	PHE	2.7
1	B	510	TYR	2.6
1	C	518	LYS	2.6
1	B	517	LYS	2.5
1	F	508	LYS	2.4
1	D	510	TYR	2.4
1	C	516	PHE	2.4
1	E	485	ASN	2.3
1	F	520	GLU	2.3
1	E	519	ASN	2.3
1	E	499	ILE	2.3
1	D	515	LYS	2.2
1	D	521	PHE	2.2
1	E	487	LEU	2.2
1	E	482	HIS	2.2
1	E	486	LYS	2.2
1	E	513	TYR	2.1
1	D	516	PHE	2.1
1	D	508	LYS	2.1
1	D	522	ARG	2.1
1	C	484	ASN	2.1
1	F	512	ASN	2.1
1	E	502	ILE	2.1
1	F	516	PHE	2.0
1	D	524	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	B	601	5/5	0.21	1.84	99,116,127,131	0

6.5 Other polymers

There are no such residues in this entry.