



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:17 PM GMT

PDB ID : 4EZC
Title : Crystal Structure of the UT-B Urea Transporter from Bos Taurus
Authors : Cao, Y.; Levin, E.J.; Zhou, M.; New York Consortium on Membrane Protein Structure (NYCOMPS)
Deposited on : 2012-05-02
Resolution : 2.36 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	384	
1	B	384	
1	C	384	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	BGC	B	402	-	X
2	BGC	C	403	-	X
3	BOG	A	403	-	X
3	BOG	A	404	-	X
3	BOG	A	405	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
3	BOG	C	404	-	X
3	BOG	C	405	-	X
3	BOG	C	407	-	X
4	SPL	A	407	-	X
4	SPL	B	406	-	X
4	SPL	C	408	-	X

2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 8470 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urea transporter 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	0	0
			2623	1735	407	462	19			
1	B	346	Total	C	N	O	S	0	0	0
			2634	1741	411	463	19			
1	C	346	Total	C	N	O	S	0	0	0
			2634	1741	411	463	19			

- Molecule 2 is SUGAR (BETA-D-GLUCOSE) (three-letter code: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: C₁₄H₂₈O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			20	14	6		
3	A	1	Total	C	O	0	0
			20	14	6		
3	A	1	Total	C	O	0	0
			20	14	6		
3	A	1	Total	C	O	0	0
			20	14	6		
3	B	1	Total	C	O	0	0
			20	14	6		
3	B	1	Total	C	O	0	0
			20	14	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			20	14	6		
3	C	1	Total	C	O	0	0
			20	14	6		
3	C	1	Total	C	O	0	0
			20	14	6		
3	C	1	Total	C	O	0	0
			20	14	6		

- Molecule 4 is OCTANOIC ACID (2-HYDROXY-1-HYDROXYMETHYL-HEPTADEC-3-ENYL)-AMIDE (three-letter code: SPL) (formula: C₂₆H₅₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			30	26	1	3		
4	B	1	Total	C	N	O	0	0
			30	26	1	3		
4	C	1	Total	C	N	O	0	0
			30	26	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total	O	0	0
			59	59		
5	B	68	Total	O	0	0
			68	68		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	66	Total	O	0	0
			66	66		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Urea transporter 1

Chain A: 




• Molecule 1: Urea transporter 1

Chain B: 



• Molecule 1: Urea transporter 1

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.70Å 105.64Å 105.87Å 90.00° 99.00° 90.00°	Depositor
Resolution (Å)	42.95 – 2.36 42.95 – 2.36	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.95-2.36) 98.9 (42.95-2.36)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.37Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.196 , 0.228 0.194 , 0.225	Depositor DCC
R_{free} test set	3283 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.351	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 66520 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8470	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, SPL, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	8/2699 (0.3%)	0.65	1/3699 (0.0%)
1	B	0.71	6/2710 (0.2%)	0.64	0/3713
1	C	0.70	4/2710 (0.1%)	0.64	0/3713
All	All	0.70	18/8119 (0.2%)	0.64	1/11125 (0.0%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	301	TRP	CD2-CE2	5.96	1.48	1.41
1	B	235	TRP	CD2-CE2	5.84	1.48	1.41
1	A	86	TRP	CD2-CE2	5.84	1.48	1.41
1	B	209	TRP	CD2-CE2	5.72	1.48	1.41
1	C	301	TRP	CD2-CE2	5.53	1.48	1.41
1	B	86	TRP	CD2-CE2	5.46	1.48	1.41
1	A	281	TRP	CD2-CE2	5.43	1.47	1.41
1	B	301	TRP	CD2-CE2	5.40	1.47	1.41
1	B	335	TRP	CD2-CE2	5.37	1.47	1.41
1	C	235	TRP	CD2-CE2	5.37	1.47	1.41
1	A	235	TRP	CD2-CE2	5.37	1.47	1.41
1	B	43	TRP	CD2-CE2	5.28	1.47	1.41
1	A	140	TRP	CD2-CE2	5.27	1.47	1.41
1	A	335	TRP	CD2-CE2	5.26	1.47	1.41
1	A	139	TRP	CD2-CE2	5.25	1.47	1.41
1	C	281	TRP	CD2-CE2	5.18	1.47	1.41
1	C	166	TRP	CD2-CE2	5.18	1.47	1.41
1	A	43	TRP	CD2-CE2	5.03	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	LEU	CA-CB-CG	5.24	127.34	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2623	0	0	2	0
1	B	2634	0	0	5	0
1	C	2634	0	0	1	0
2	A	24	0	24	0	0
2	B	36	0	36	0	0
2	C	36	0	36	0	0
3	A	80	0	112	0	0
3	B	40	0	56	0	0
3	C	80	0	112	0	0
4	A	30	0	51	0	0
4	B	30	0	51	0	0
4	C	30	0	51	0	0
5	A	59	0	0	1	0
5	B	68	0	0	1	0
5	C	66	0	0	1	0
All	All	8470	0	529	8	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (8) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:202:THR:N	5:A:536:HOH:O	2.39	0.55
1:B:151:THR:CG2	1:B:155:PHE:CE2	2.94	0.50
1:C:356:LYS:NZ	5:C:539:HOH:O	2.46	0.47
1:A:302:GLN:OE1	1:A:302:GLN:N	2.48	0.46
1:B:151:THR:CG2	1:B:155:PHE:CD2	3.00	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:173:LEU:N	1:B:174:PRO:CD	2.80	0.44
1:B:350:LYS:NZ	5:B:563:HOH:O	2.54	0.41
1:B:349:THR:CG2	1:B:351:ASN:N	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/384 (89%)	332 (97%)	11 (3%)	0	100	100
1	B	344/384 (90%)	334 (97%)	10 (3%)	0	100	100
1	C	344/384 (90%)	335 (97%)	8 (2%)	1 (0%)	50	62
All	All	1031/1152 (90%)	1001 (97%)	29 (3%)	1 (0%)	59	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	35	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/319 (90%)	282 (98%)	6 (2%)	66	82
1	B	289/319 (91%)	282 (98%)	7 (2%)	61	78
1	C	289/319 (91%)	286 (99%)	3 (1%)	85	93
All	All	866/957 (90%)	850 (98%)	16 (2%)	71	85

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	THR
1	A	38	LYS
1	A	46	ASP
1	A	110	SER
1	A	303	THR
1	A	337	PHE
1	B	34	THR
1	B	46	ASP
1	B	133	ASP
1	B	142	LEU
1	B	151	THR
1	B	303	THR
1	B	322	SER
1	C	34	THR
1	C	109	ARG
1	C	208	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	A	401	-	12,12,12	0.58	0	17,17,17	0.95	0
2	BGC	A	402	-	12,12,12	0.57	0	17,17,17	1.62	2 (11%)
3	BOG	A	403	-	20,20,20	0.58	0	25,25,25	0.83	1 (4%)
3	BOG	A	404	-	20,20,20	0.68	0	25,25,25	0.78	0
3	BOG	A	405	-	20,20,20	0.50	0	25,25,25	0.69	0
3	BOG	A	406	-	20,20,20	0.59	0	25,25,25	1.04	2 (8%)
4	SPL	A	407	-	29,29,29	0.55	0	31,31,31	1.54	5 (16%)
2	BGC	B	401	-	12,12,12	0.71	0	17,17,17	1.68	4 (23%)
2	BGC	B	402	-	12,12,12	0.61	0	17,17,17	0.84	0
2	BGC	B	403	-	12,12,12	0.58	0	17,17,17	1.33	1 (5%)
3	BOG	B	404	-	20,20,20	0.59	0	25,25,25	0.77	0
3	BOG	B	405	-	20,20,20	0.54	0	25,25,25	0.99	2 (8%)
4	SPL	B	406	-	29,29,29	0.52	0	31,31,31	1.18	4 (12%)
2	BGC	C	401	-	12,12,12	0.41	0	17,17,17	0.64	0
2	BGC	C	402	-	12,12,12	0.51	0	17,17,17	1.57	5 (29%)
2	BGC	C	403	-	12,12,12	0.53	0	17,17,17	0.82	1 (5%)
3	BOG	C	404	-	20,20,20	0.67	1 (5%)	25,25,25	0.99	1 (4%)
3	BOG	C	405	-	20,20,20	0.55	0	25,25,25	0.68	0
3	BOG	C	406	-	20,20,20	0.55	0	25,25,25	0.80	0
3	BOG	C	407	-	20,20,20	0.59	0	25,25,25	0.62	0
4	SPL	C	408	-	29,29,29	0.54	0	31,31,31	1.36	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	401	-	-	0/2/22/22	0/1/1/1
2	BGC	A	402	-	-	0/2/22/22	0/1/1/1
3	BOG	A	403	-	-	0/11/31/31	0/1/1/1
3	BOG	A	404	-	-	0/11/31/31	0/1/1/1
3	BOG	A	405	-	-	0/11/31/31	0/1/1/1
3	BOG	A	406	-	-	0/11/31/31	0/1/1/1
4	SPL	A	407	-	-	0/32/32/32	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	401	-	-	0/2/22/22	0/1/1/1
2	BGC	B	402	-	-	0/2/22/22	0/1/1/1
2	BGC	B	403	-	-	0/2/22/22	0/1/1/1
3	BOG	B	404	-	-	0/11/31/31	0/1/1/1
3	BOG	B	405	-	-	0/11/31/31	0/1/1/1
4	SPL	B	406	-	-	0/32/32/32	0/0/0/0
2	BGC	C	401	-	-	0/2/22/22	0/1/1/1
2	BGC	C	402	-	-	0/2/22/22	0/1/1/1
2	BGC	C	403	-	-	0/2/22/22	0/1/1/1
3	BOG	C	404	-	-	0/11/31/31	0/1/1/1
3	BOG	C	405	-	-	0/11/31/31	0/1/1/1
3	BOG	C	406	-	-	0/11/31/31	0/1/1/1
3	BOG	C	407	-	-	0/11/31/31	0/1/1/1
4	SPL	C	408	-	-	0/32/32/32	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	404	BOG	O1-C1	2.13	1.44	1.40

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	407	SPL	C2-C3-C4	5.33	119.08	111.84
4	C	408	SPL	C2-C3-C4	4.45	117.89	111.84
2	B	401	BGC	C1-C2-C3	-4.13	103.99	110.53
2	B	403	BGC	C1-O5-C5	3.59	119.82	113.40
2	A	402	BGC	O3-C3-C2	3.49	118.19	110.35
3	C	404	BOG	O1-C1-C2	3.35	112.44	108.18
2	A	402	BGC	C4-C3-C2	-3.32	104.68	110.82
2	C	402	BGC	O5-C5-C4	3.25	115.78	109.76
2	C	402	BGC	C1-O5-C5	3.16	119.06	113.40
4	A	407	SPL	O3-C3-C4	-3.02	102.21	110.90
4	B	406	SPL	C20-C19-N2	2.99	121.50	115.83
4	A	407	SPL	C1-C2-N2	2.77	114.25	109.35
2	B	401	BGC	O5-C1-C2	-2.67	105.72	109.86
3	A	406	BOG	O5-C5-C4	2.56	114.50	109.76
3	A	406	BOG	C3-C4-C5	2.56	114.77	110.20
4	A	407	SPL	C2-N2-C19	2.55	128.39	122.97
4	B	406	SPL	C2-C3-C4	2.54	115.29	111.84
4	C	408	SPL	O3-C3-C4	-2.48	103.75	110.90
3	B	405	BOG	C4-C3-C2	2.48	115.40	110.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	407	SPL	O19-C19-C20	-2.46	117.02	121.92
4	C	408	SPL	C2-N2-C19	2.46	128.20	122.97
2	B	401	BGC	O2-C2-C1	2.40	115.00	109.89
2	C	402	BGC	C1-C2-C3	-2.39	106.75	110.53
3	A	403	BOG	O1-C1-C2	2.35	111.17	108.18
3	B	405	BOG	C1-C2-C3	2.34	114.56	110.00
2	B	401	BGC	C4-C3-C2	-2.33	106.51	110.82
2	C	402	BGC	C4-C3-C2	-2.31	106.54	110.82
4	C	408	SPL	C1-C2-N2	2.29	113.40	109.35
4	C	408	SPL	O19-C19-C20	-2.28	117.38	121.92
4	B	406	SPL	C1-C2-N2	2.27	113.37	109.35
2	C	403	BGC	O5-C5-C6	2.21	111.78	106.34
4	B	406	SPL	C1-C2-C3	-2.20	107.29	113.12
2	C	402	BGC	C6-C5-C4	-2.14	107.83	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/384 (89%)	0.43	24 (6%) 16 18	38, 54, 97, 132	0
1	B	346/384 (90%)	0.48	37 (10%) 6 8	39, 55, 88, 125	0
1	C	346/384 (90%)	0.40	34 (9%) 8 9	40, 55, 87, 130	0
All	All	1037/1152 (90%)	0.44	95 (9%) 9 10	38, 55, 91, 132	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	46	ASP	6.9
1	B	32	TYR	6.0
1	A	373	LEU	5.9
1	C	33	ILE	5.7
1	B	212	LEU	5.3
1	C	51	LEU	5.1
1	A	39	GLU	4.7
1	A	301	TRP	4.7
1	A	43	TRP	4.7
1	A	40	PHE	4.5
1	B	138	PHE	4.4
1	A	32	TYR	4.4
1	B	43	TRP	4.3
1	C	43	TRP	4.3
1	B	48	PRO	4.3
1	C	373	LEU	4.2
1	C	32	TYR	4.1
1	C	47	LYS	4.0
1	B	47	LYS	4.0
1	B	235	TRP	3.7
1	C	48	PRO	3.6
1	B	51	LEU	3.6
1	C	40	PHE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	31	GLY	3.4
1	A	38	LYS	3.3
1	A	138	PHE	3.3
1	A	42	ASN	3.3
1	C	376	ARG	3.3
1	C	375	SER	3.2
1	A	37	MET	3.2
1	B	44	LEU	3.2
1	A	339	LEU	3.1
1	C	372	TYR	3.1
1	C	46	ASP	3.1
1	A	206	ASN	3.1
1	A	41	ALA	3.1
1	A	49	GLN	3.0
1	B	335	TRP	3.0
1	B	372	TYR	2.8
1	B	373	LEU	2.8
1	C	273	PHE	2.8
1	B	207	VAL	2.8
1	A	338	CYS	2.7
1	C	50	ALA	2.7
1	A	104	LEU	2.7
1	B	177	MET	2.7
1	B	173	LEU	2.7
1	C	159	LEU	2.6
1	C	339	LEU	2.6
1	B	172	THR	2.6
1	C	353	ASN	2.5
1	C	56	TRP	2.5
1	C	109	ARG	2.5
1	C	250	LEU	2.5
1	C	104	LEU	2.5
1	C	173	LEU	2.5
1	B	321	MET	2.5
1	C	41	ALA	2.5
1	B	287	LEU	2.5
1	A	335	TRP	2.5
1	B	41	ALA	2.5
1	C	140	TRP	2.4
1	C	34	THR	2.4
1	C	340	ALA	2.4
1	A	310	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	338	CYS	2.4
1	B	82	VAL	2.4
1	B	39	GLU	2.4
1	B	216	GLN	2.4
1	B	104	LEU	2.3
1	B	301	TRP	2.3
1	C	177	MET	2.3
1	C	371	PHE	2.3
1	C	370	ILE	2.3
1	B	175	PHE	2.3
1	B	81	LEU	2.3
1	C	352	PRO	2.2
1	B	141	LEU	2.2
1	B	176	ASN	2.2
1	B	42	ASN	2.2
1	B	120	ASN	2.2
1	A	56	TRP	2.2
1	B	375	SER	2.1
1	A	374	GLN	2.1
1	A	109	ARG	2.1
1	A	369	ARG	2.1
1	B	56	TRP	2.1
1	A	51	LEU	2.1
1	C	212	LEU	2.1
1	B	210	PRO	2.1
1	C	207	VAL	2.0
1	B	180	SER	2.0
1	A	173	LEU	2.0
1	C	369	ARG	2.0
1	C	53	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	BOG	A	405	20/20	0.31	12.97	55,99,115,117	0
2	BGC	B	402	12/12	0.21	12.00	92,102,107,107	0
3	BOG	C	407	20/20	0.45	6.35	69,134,152,154	0
4	SPL	B	406	30/30	0.53	6.23	61,81,92,103	0
4	SPL	C	408	30/30	0.44	4.94	59,72,88,96	0
3	BOG	A	404	20/20	0.39	4.86	59,81,85,89	0
3	BOG	C	404	20/20	0.39	4.57	71,80,91,96	0
3	BOG	A	403	20/20	0.33	4.48	65,97,125,136	0
4	SPL	A	407	30/30	0.39	4.34	55,72,82,86	0
3	BOG	C	405	20/20	0.30	3.32	63,74,81,82	0
2	BGC	C	403	12/12	0.19	2.71	87,93,104,109	0
3	BOG	A	406	20/20	0.33	1.76	67,110,129,134	0
3	BOG	B	405	20/20	0.37	1.71	66,103,129,130	0
2	BGC	A	402	12/12	0.31	1.60	74,93,103,112	0
3	BOG	B	404	20/20	0.24	1.21	71,110,122,126	0
3	BOG	C	406	20/20	0.20	0.92	70,101,109,110	0
2	BGC	B	403	12/12	0.22	0.01	68,85,93,95	0
2	BGC	C	402	12/12	0.16	-0.27	86,96,103,109	0
2	BGC	B	401	12/12	0.19	-0.63	74,79,81,83	0
2	BGC	A	401	12/12	0.14	-0.89	73,80,96,107	0
2	BGC	C	401	12/12	0.25	-	91,95,101,102	0

6.5 Other polymers ⓘ

There are no such residues in this entry.