



wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 08:30 PM GMT

PDB ID : 2F55
Title : Two hepatitis c virus ns3 helicase domains complexed with the same strand of dna
Authors : Lu, J.Z.; Jordan, J.B.; Sakon, J.
Deposited on : 2005-11-25
Resolution : 3.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1696 (3.40-3.20)
Ramachandran outliers	78287	1664 (3.40-3.20)
Sidechain outliers	78261	1662 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	D	13	
2	E	3	
3	A	435	
3	B	435	
3	C	435	




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	20	Total	O	0	0
			20	20		
5	B	14	Total	O	0	0
			14	14		
5	C	25	Total	O	0	0
			25	25		
5	D	3	Total	O	0	0
			3	3		


• Molecule 3: polyprotein

Chain B:



• Molecule 3: polyprotein

Chain C:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	108.20Å 109.80Å 183.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30	Depositor
% Data completeness (in resolution range)	83.2 (50.00-3.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.247 , 0.273	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10129	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.70	0/272	0.98	1/412 (0.2%)
2	E	0.70	0/62	0.99	0/92
3	A	0.54	3/3331 (0.1%)	0.73	2/4548 (0.0%)
3	B	0.59	4/3331 (0.1%)	0.74	1/4548 (0.0%)
3	C	0.53	5/3331 (0.2%)	0.72	1/4548 (0.0%)
All	All	0.56	12/10327 (0.1%)	0.74	5/14148 (0.0%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	469	ARG	NE-CZ	7.51	1.42	1.33
3	B	469	ARG	NE-CZ	7.47	1.42	1.33
3	A	469	ARG	NE-CZ	7.45	1.42	1.33
3	B	469	ARG	CZ-NH2	7.42	1.42	1.33
3	C	469	ARG	CZ-NH2	7.38	1.42	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	402	PRO	CA-N-CD	-12.76	93.64	111.50
3	B	402	PRO	CA-N-CD	-12.74	93.67	111.50
3	A	402	PRO	CA-N-CD	-11.23	95.78	111.50
3	A	401	PRO	C-N-CD	7.38	143.90	128.40
1	D	17	DU	O4'-C1'-C2'	5.03	109.92	105.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	247	0	131	18	0
2	E	57	0	31	6	0
3	A	3251	0	3216	332	0
3	B	3251	0	3215	434	0
3	C	3251	0	3215	354	0
4	A	5	0	0	0	0
4	C	5	0	0	1	0
5	A	20	0	0	2	0
5	B	14	0	0	0	0
5	C	25	0	0	3	0
5	D	3	0	0	0	0
All	All	10129	0	9808	1118	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 56.

The worst 5 of 1118 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:469:ARG:HB2	3:A:469:ARG:NH1	1.31	1.43
3:C:513:LEU:O	3:C:517:LEU:CD2	1.65	1.42
3:A:360:LYS:NZ	3:A:386:LEU:HD21	1.06	1.36
3:A:360:LYS:NZ	3:A:386:LEU:CD2	1.88	1.35
3:A:339:ILE:CG2	3:A:474:ARG:HG2	1.54	1.35

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	428/435 (98%)	318 (74%)	86 (20%)	24 (6%)	3 25
3	B	428/435 (98%)	282 (66%)	104 (24%)	42 (10%)	1 8
3	C	428/435 (98%)	296 (69%)	92 (22%)	40 (9%)	1 9
All	All	1284/1305 (98%)	896 (70%)	282 (22%)	106 (8%)	1 12

5 of 106 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	246	HIS
3	A	349	PHE
3	A	413	ALA
3	A	583	LYS
3	B	245	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	352/354 (99%)	323 (92%)	29 (8%)	17 57
3	B	352/354 (99%)	311 (88%)	41 (12%)	8 35
3	C	352/354 (99%)	320 (91%)	32 (9%)	14 49
All	All	1056/1062 (99%)	954 (90%)	102 (10%)	12 45

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	393	ARG
3	B	524	VAL
3	C	526	GLN
3	B	402	PRO
3	B	460	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
3	B	564	GLN
3	C	198	GLN
3	C	564	GLN
3	B	580	GLN
3	C	201	HIS

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	101	-	4,4,4	0.31	0	6,6,6	0.20	0
4	SO4	C	102	-	4,4,4	0.20	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	101	-	-	0/0/0/0	0/0/0/0
4	SO4	C	102	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.