



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 12:51 PM GMT

PDB ID : 4F5Y
Title : Crystal structure of human STING CTD complex with C-di-GMP
Authors : Gu, L.; Shang, G.; Zhu, D.; Li, N.; Zhang, J.; Zhu, C.; Lu, D.; Liu, C.; Yu, Q.; Zhao, Y.; Xu, S.
Deposited on : 2012-05-13
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	239	
1	B	239	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3292 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protein 173.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	0	0	0
			1553	974	274	299	6			
1	B	193	Total	C	N	O	S	0	0	0
			1553	974	274	299	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	ARG	HIS	SEE REMARK 999	UNP Q86WV6
A	380	LEU	-	EXPRESSION TAG	UNP Q86WV6
A	381	GLU	-	EXPRESSION TAG	UNP Q86WV6
A	382	HIS	-	EXPRESSION TAG	UNP Q86WV6
A	383	HIS	-	EXPRESSION TAG	UNP Q86WV6
A	384	HIS	-	EXPRESSION TAG	UNP Q86WV6
A	385	HIS	-	EXPRESSION TAG	UNP Q86WV6
A	386	HIS	-	EXPRESSION TAG	UNP Q86WV6
A	387	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	232	ARG	HIS	SEE REMARK 999	UNP Q86WV6
B	380	LEU	-	EXPRESSION TAG	UNP Q86WV6
B	381	GLU	-	EXPRESSION TAG	UNP Q86WV6
B	382	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	383	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	384	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	385	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	386	HIS	-	EXPRESSION TAG	UNP Q86WV6
B	387	HIS	-	EXPRESSION TAG	UNP Q86WV6

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 9,9'-[(2R,3R,3AS,5S,7AR,9R,10R,10AS,12S,14AR)-3,5,10,12-TETRAHYDROXY-5,12-DIOXIDOOCTAHYDRO-2H,7H-DIFURO[3,2-D:3',2'-J][1,3,7,9,2,8]TETRAOXA DIPHOSPHACYCLODODECINE-2,9-DIYL]BIS(2-AMINO-1,9-DIHYDRO-6H-PURIN-6-ONE) (three-letter code: C2E) (formula: C₂₀H₂₄N₁₀O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			46	20	10	14	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	71	Total	O	0	0
			71	71		
4	B	67	Total	O	0	0
			67	67		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Transmembrane protein 173

Chain A: 



• Molecule 1: Transmembrane protein 173

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.14Å 81.44Å 91.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.84 – 2.40 46.75 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.84-2.40) 99.8 (46.75-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.211 , 0.275 0.207 , 0.271	Depositor DCC
R_{free} test set	1126 reflections (5.13%)	DCC
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 21983 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3292	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.73 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.6309e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1582	0.63	2/2144 (0.1%)
1	B	0.43	0/1582	0.58	0/2144
All	All	0.44	0/3164	0.60	2/4288 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	319	ASP	N-CA-C	-5.54	96.05	111.00
1	A	259	LEU	CA-CB-CG	5.29	127.48	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	228	GLN	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1553	0	1517	16	0
1	B	1553	0	1517	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	46	0	22	0	0
4	A	71	0	0	4	0
4	B	67	0	0	2	0
All	All	3292	0	3056	46	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (46) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:238:ARG:HG3	1:B:239:VAL:H	1.25	1.01
1:B:237:ASP:HB3	1:B:240:TYR:CE1	2.13	0.82
1:B:237:ASP:CB	1:B:240:TYR:CE1	2.65	0.80
1:A:224:LYS:NZ	1:A:243:SER:OG	2.22	0.72
1:B:237:ASP:HB3	1:B:240:TYR:HE1	1.55	0.71
1:B:173:PRO:HG2	1:B:236:LYS:HE2	1.73	0.70
1:B:238:ARG:HG3	1:B:239:VAL:N	2.05	0.67
1:A:178:ARG:NH1	1:A:240:TYR:OH	2.28	0.67
1:B:293:ARG:NH2	4:B:629:HOH:O	2.28	0.66
1:A:178:ARG:HE	1:A:227:GLN:HA	1.59	0.66
1:B:324:SER:HB3	1:B:327:GLN:HB2	1.77	0.66
1:B:237:ASP:HB2	1:B:240:TYR:CE1	2.35	0.61
1:A:296:GLU:HG3	1:A:311:LEU:HD12	1.83	0.61
1:B:235:ILE:HG22	1:B:236:LYS:HG3	1.83	0.60
1:A:305:SER:HA	1:A:308:ASN:HB2	1.83	0.59
1:A:169:ARG:HD2	4:A:547:HOH:O	2.05	0.57
1:B:284:ARG:HD2	4:B:636:HOH:O	2.05	0.56
1:A:198:LEU:HD11	1:A:260:GLU:HG2	1.87	0.55
1:B:224:LYS:HZ2	1:B:224:LYS:HA	1.71	0.54
1:B:176:GLN:OE1	1:B:180:ARG:NH1	2.41	0.54
1:B:299:LEU:O	1:B:306:GLN:HB2	2.08	0.54
1:B:174:GLU:HG2	1:B:228:GLN:HG3	1.89	0.53
1:B:199:TYR:HD2	1:B:312:ILE:HD11	1.75	0.51
1:B:216:ASP:HB3	1:B:219:ILE:HG13	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:237:ASP:HB2	1:B:240:TYR:CD1	2.46	0.50
1:B:177:ALA:O	1:B:181:THR:HG23	2.12	0.49
1:A:305:SER:OG	4:A:565:HOH:O	2.20	0.48
1:A:338:LYS:NZ	4:A:566:HOH:O	2.48	0.47
1:B:319:ASP:N	1:B:319:ASP:OD1	2.36	0.47
1:A:253:ARG:HD2	4:A:543:HOH:O	2.16	0.46
1:B:206:CYS:HB2	1:B:284:ARG:HD3	1.98	0.45
1:B:317:PRO:HB2	1:B:321:SER:HB3	1.98	0.45
1:A:317:PRO:O	1:A:319:ASP:O	2.34	0.45
1:B:328:GLU:OE1	1:B:331:ARG:NH2	2.43	0.45
1:B:237:ASP:CB	1:B:240:TYR:CD1	3.00	0.44
1:B:321:SER:OG	1:B:322:SER:N	2.49	0.43
1:A:235:ILE:H	1:A:235:ILE:HD13	1.84	0.43
1:B:238:ARG:O	1:B:239:VAL:HG23	2.18	0.43
1:A:179:ILE:HA	1:A:225:LEU:HD11	1.99	0.43
1:B:199:TYR:CD2	1:B:312:ILE:HD11	2.54	0.42
1:B:304:GLU:HG3	1:B:304:GLU:O	2.15	0.42
1:A:164:TYR:O	1:A:169:ARG:HB2	2.21	0.41
1:A:316:GLU:HA	1:A:317:PRO:HD3	1.95	0.41
1:A:152:ASN:CG	1:B:152:ASN:HB2	2.42	0.40
1:B:330:LEU:O	1:B:334:ARG:HG3	2.21	0.40
1:B:198:LEU:HG	1:B:200:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/239 (80%)	179 (94%)	11 (6%)	1 (0%)	38	53
1	B	191/239 (80%)	178 (93%)	12 (6%)	1 (0%)	38	53
All	All	382/478 (80%)	357 (94%)	23 (6%)	2 (0%)	38	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	GLU
1	B	303	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/211 (80%)	152 (90%)	16 (10%)	12	18
1	B	168/211 (80%)	144 (86%)	24 (14%)	5	5
All	All	336/422 (80%)	296 (88%)	40 (12%)	8	10

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	A	170	LEU
1	A	189	LEU
1	A	211	ASN
1	A	212	LEU
1	A	224	LYS
1	A	231	ASP
1	A	235	ILE
1	A	236	LYS
1	A	237	ASP
1	A	241	SER
1	A	250	ASN
1	A	304	GLU
1	A	307	ASN
1	A	308	ASN
1	A	341	VAL
1	B	169	ARG
1	B	170	LEU
1	B	171	ILE
1	B	174	GLU
1	B	176	GLN
1	B	180	ARG
1	B	189	LEU
1	B	212	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	214	MET
1	B	223	ASP
1	B	224	LYS
1	B	228	GLN
1	B	231	ASP
1	B	232	ARG
1	B	282	GLU
1	B	296	GLU
1	B	304	GLU
1	B	319	ASP
1	B	320	ASP
1	B	321	SER
1	B	325	LEU
1	B	340	GLU
1	B	341	VAL
1	B	343	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	157	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	C2E	A	402	-	52,52,52	2.91	17 (32%)	76,82,82	3.18	17 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C2E	A	402	-	-	0/30/62/62	0/0/7/7

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	C2E	O61-C61	9.00	1.42	1.24
3	A	402	C2E	C2-N2	8.18	1.45	1.32
3	A	402	C2E	C21-N21	7.99	1.44	1.32
3	A	402	C2E	O6-C6	6.83	1.37	1.24
3	A	402	C2E	C21-N11	5.97	1.46	1.36
3	A	402	C2E	C2A-C1A	-3.86	1.48	1.53
3	A	402	C2E	C2-N3	3.63	1.38	1.33
3	A	402	C2E	C61-C51	3.58	1.47	1.41
3	A	402	C2E	C21-N31	3.08	1.37	1.33
3	A	402	C2E	C4-N3	3.03	1.40	1.35
3	A	402	C2E	C6-N1	2.98	1.42	1.37
3	A	402	C2E	C2-N1	2.92	1.41	1.36
3	A	402	C2E	C2'-C3'	-2.65	1.46	1.53
3	A	402	C2E	O4A-C4A	-2.58	1.39	1.45
3	A	402	C2E	O3A-C3A	-2.58	1.35	1.44
3	A	402	C2E	C41-N31	2.40	1.39	1.35
3	A	402	C2E	C5-C4	2.36	1.45	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	C2E	C6-C5-N7	17.38	136.48	134.14
3	A	402	C2E	C61-C51-N71	13.66	135.98	134.14
3	A	402	C2E	O4'-C1'-N9	8.45	116.30	108.44
3	A	402	C2E	C2-N3-C4	5.00	122.11	115.09
3	A	402	C2E	N3-C4-N9	4.39	133.34	126.91
3	A	402	C2E	C5-C4-N3	-4.22	119.83	125.94
3	A	402	C2E	C4-C5-N7	-3.73	106.33	109.52
3	A	402	C2E	N21-C21-N11	3.67	121.90	117.86
3	A	402	C2E	C21-N31-C41	3.60	120.15	115.09
3	A	402	C2E	C41-C51-N71	-3.10	106.86	109.52
3	A	402	C2E	N31-C41-N91	2.69	130.86	126.91
3	A	402	C2E	O4A-C1A-N91	2.67	110.92	108.44
3	A	402	C2E	C51-C41-N31	-2.63	122.13	125.94
3	A	402	C2E	C2A-C1A-N91	-2.48	106.89	113.27
3	A	402	C2E	C8-N9-C4	2.46	108.78	106.90
3	A	402	C2E	C81-N91-C41	2.22	108.59	106.90
3	A	402	C2E	C1'-N9-C4	-2.11	123.00	126.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	193/239 (80%)	0.85	23 (11%) 5 4	25, 41, 97, 115	0
1	B	193/239 (80%)	0.92	20 (10%) 7 6	27, 41, 97, 120	0
All	All	386/478 (80%)	0.88	43 (11%) 6 5	25, 41, 97, 120	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	235	ILE	14.7
1	B	236	LYS	14.2
1	A	235	ILE	11.1
1	B	151	GLY	8.3
1	B	237	ASP	7.1
1	B	234	GLY	6.9
1	A	318	ALA	6.8
1	B	214	MET	6.8
1	A	231	ASP	6.5
1	A	239	VAL	5.4
1	B	233	ALA	5.2
1	A	232	ARG	4.9
1	B	229	THR	4.8
1	A	234	GLY	4.4
1	A	236	LYS	4.2
1	B	215	ALA	4.2
1	B	239	VAL	4.0
1	A	306	GLN	4.0
1	A	230	GLY	3.9
1	B	318	ALA	3.9
1	A	153	PHE	3.8
1	A	238	ARG	3.5
1	A	215	ALA	3.4
1	B	240	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	238	ARG	3.1
1	A	229	THR	3.0
1	A	227	GLN	3.0
1	A	240	TYR	3.0
1	B	231	ASP	3.0
1	B	232	ARG	2.9
1	B	320	ASP	2.9
1	B	322	SER	2.9
1	A	151	GLY	2.8
1	A	214	MET	2.8
1	A	237	ASP	2.8
1	B	228	GLN	2.7
1	B	303	PRO	2.6
1	A	228	GLN	2.5
1	A	189	LEU	2.5
1	A	322	SER	2.4
1	A	319	ASP	2.4
1	A	233	ALA	2.1
1	B	211	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	B	500	1/1	0.20	1.61	61,61,61,61	0
2	CA	A	401	1/1	0.16	0.07	57,57,57,57	0
3	C2E	A	402	46/46	0.15	-0.43	26,36,46,58	0

6.5 Other polymers ⓘ

There are no such residues in this entry.