



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 10:09 PM GMT

PDB ID : 3F8R
Title : Crystal structure of Sulfolobus solfataricus Thioredoxin reductase B3 in complex with two NADP molecules
Authors : Ruggiero, A.; Masullo, M.; Ruocco, M.R.; Arcari, P.; Zagari, A.; Vitagliano, L.
Deposited on : 2008-11-13
Resolution : 1.95 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1321 (1.96-1.96)
Clashscore	79885	1488 (1.96-1.96)
Ramachandran outliers	78287	1475 (1.96-1.96)
Sidechain outliers	78261	1475 (1.96-1.96)
RSRZ outliers	66119	1321 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	323	
1	B	323	
1	C	323	
1	D	323	

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 10058 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin reductase (TrxB-3).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2367	1523	395	443	6			
1	B	309	Total	C	N	O	S	0	0	0
			2375	1527	396	446	6			
1	C	307	Total	C	N	O	S	0	0	0
			2356	1517	391	442	6			
1	D	298	Total	C	N	O	S	0	0	0
			2285	1469	381	429	6			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			30	10	5	12	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total	O	0	0
			88	88		
3	B	82	Total	O	0	0
			82	82		
3	C	89	Total	O	0	0
			89	89		
3	D	101	Total	O	0	0
			101	101		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Thioredoxin reductase (TrxB-3)

Chain A:




- Molecule 1: Thioredoxin reductase (TrxB-3)

Chain B:




- Molecule 1: Thioredoxin reductase (TrxB-3)

Chain C:



● Molecule 1: Thioredoxin reductase (TrxB-3)

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.87Å 123.42Å 127.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.95 19.97 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-1.95) 86.2 (19.97-1.95)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 1.94Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.208 , 0.257 0.210 , 0.244	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.1	EDS
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 80321 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10058	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/2407	0.71	0/3252
1	B	0.55	0/2415	0.72	1/3263 (0.0%)
1	C	0.57	0/2396	0.71	0/3238
1	D	0.54	0/2324	0.71	0/3141
All	All	0.55	0/9542	0.71	1/12894 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	13	GLY	N-CA-C	-5.01	100.57	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2367	0	2428	62	0
1	B	2375	0	2432	78	0
1	C	2356	0	2415	79	0
1	D	2285	0	2330	72	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	79	0	36	7	0
2	B	79	0	36	8	0
2	C	79	0	36	10	0
2	D	78	0	36	6	0
3	A	88	0	0	11	0
3	B	82	0	0	12	0
3	C	89	0	0	3	0
3	D	101	0	0	7	0
All	All	10058	0	9749	281	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:156:VAL:H	1:B:241:ASN:ND2	1.61	0.97
1:A:105:LYS:HG2	1:A:111:GLU:HG2	1.52	0.90
1:C:190:LYS:H	1:C:190:LYS:HD3	1.36	0.89
1:A:219:ASP:CG	1:A:220:LYS:H	1.77	0.89
1:D:214:LYS:HD2	1:D:228:GLU:HB2	1.55	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/323 (94%)	295 (97%)	9 (3%)	0	100	100
1	B	305/323 (94%)	288 (94%)	14 (5%)	3 (1%)	22	9
1	C	303/323 (94%)	293 (97%)	10 (3%)	0	100	100
1	D	294/323 (91%)	278 (95%)	14 (5%)	2 (1%)	30	15
All	All	1206/1292 (93%)	1154 (96%)	47 (4%)	5 (0%)	43	30

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	109	LYS
1	B	14	GLU
1	B	233	GLY
1	D	14	GLU
1	D	230	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/267 (95%)	250 (99%)	3 (1%)	82	79
1	B	254/267 (95%)	248 (98%)	6 (2%)	61	51
1	C	252/267 (94%)	248 (98%)	4 (2%)	75	69
1	D	243/267 (91%)	240 (99%)	3 (1%)	82	79
All	All	1002/1068 (94%)	986 (98%)	16 (2%)	75	69

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	225	VAL
1	B	293	LEU
1	C	265	THR
1	B	190	LYS
1	D	108	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	297	GLN
1	C	97	ASN
1	C	241	ASN
1	B	266	ASN
1	C	192	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAP	A	4001	-	52,52,52	1.47	6 (11%)	80,80,80	1.37	11 (13%)
2	NAP	A	4005	-	33,33,52	2.22	6 (18%)	52,52,80	1.26	9 (17%)
2	NAP	B	4002	-	52,52,52	1.43	8 (15%)	80,80,80	1.45	11 (13%)
2	NAP	B	4006	-	33,33,52	2.00	6 (18%)	52,52,80	1.25	6 (11%)
2	NAP	C	4003	-	52,52,52	1.46	6 (11%)	80,80,80	1.38	10 (12%)
2	NAP	C	4007	-	33,33,52	2.14	6 (18%)	52,52,80	1.39	5 (9%)
2	NAP	D	4004	-	52,52,52	1.50	6 (11%)	80,80,80	1.46	10 (12%)
2	NAP	D	4008	-	30,32,52	1.48	4 (13%)	45,49,80	1.45	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	4001	-	-	0/35/67/67	0/3/5/5
2	NAP	A	4005	-	-	0/21/37/67	0/1/3/5
2	NAP	B	4002	-	-	0/35/67/67	0/3/5/5
2	NAP	B	4006	-	-	0/21/37/67	0/1/3/5
2	NAP	C	4003	-	-	0/35/67/67	0/3/5/5
2	NAP	C	4007	-	-	0/21/37/67	0/1/3/5
2	NAP	D	4004	-	-	0/35/67/67	0/3/5/5
2	NAP	D	4008	-	-	0/18/36/67	0/1/3/5

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	4005	NAP	PN-O5D	8.88	1.66	1.51
2	C	4007	NAP	PN-O5D	7.75	1.65	1.51
2	B	4006	NAP	PN-O5D	7.60	1.64	1.51
2	D	4004	NAP	C3N-C7N	5.47	1.59	1.50
2	A	4001	NAP	C3N-C7N	5.44	1.59	1.50

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4008	NAP	O4B-C1B-C2B	-5.62	101.69	106.95
2	B	4002	NAP	O4D-C1D-N1N	5.41	113.49	107.95
2	C	4007	NAP	O4B-C1B-C2B	-4.92	102.35	106.95
2	C	4003	NAP	C3N-C7N-N7N	4.71	123.13	117.77
2	A	4001	NAP	C3N-C7N-N7N	4.60	123.01	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/323 (95%)	0.21	8 (2%) 53 61	8, 22, 45, 59	0
1	B	309/323 (95%)	0.44	15 (4%) 28 35	9, 25, 48, 70	0
1	C	307/323 (95%)	0.32	19 (6%) 20 25	8, 21, 55, 68	0
1	D	298/323 (92%)	0.42	28 (9%) 9 11	7, 22, 57, 71	0
All	All	1222/1292 (94%)	0.35	70 (5%) 22 29	7, 23, 52, 71	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	194	ILE	6.8
1	D	219	ASP	6.6
1	A	98	ARG	4.5
1	B	98	ARG	4.2
1	D	195	TYR	4.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAP	A	4001	48/48	0.20	1.08	36,48,77,79	0
2	NAP	D	4008	30/48	0.14	0.73	11,18,53,55	0
2	NAP	A	4005	31/48	0.13	0.65	8,16,43,46	0
2	NAP	D	4004	48/48	0.20	0.35	33,54,76,79	0
2	NAP	C	4007	31/48	0.12	0.32	10,14,33,37	0
2	NAP	B	4002	48/48	0.16	0.17	28,39,70,72	0
2	NAP	C	4003	48/48	0.17	-0.04	33,42,74,76	0
2	NAP	B	4006	31/48	0.12	-0.20	13,25,48,50	0

6.5 Other polymers ⓘ

There are no such residues in this entry.