



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 09:22 PM GMT

PDB ID : 4F8H  
Title : X-ray Structure of the Anesthetic Ketamine Bound to the GLIC Pentameric  
Ligand-gated Ion Channel  
Authors : Pan, J.J.; Chen, Q.; Willenbring, D.; Kong, X.P.; Cohen, A.; Xu, Y.; Tang, P.  
Deposited on : 2012-05-17  
Resolution : 2.99 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	317	
1	B	317	
1	C	317	
1	D	317	
1	E	317	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	RKE	A	401	-	X
2	RKE	B	401	-	X
2	RKE	C	401	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
2	RKE	D	401	-	X
2	RKE	E	401	-	X
3	LMD	A	402	-	X
3	LMD	B	402	-	X
3	LMD	C	402	-	X
3	LMD	C	405	-	X
3	LMD	D	402	-	X
3	LMD	E	402	-	X
4	PC1	A	403	-	X
4	PC1	A	404	-	X
4	PC1	B	403	-	X
4	PC1	B	404	-	X
4	PC1	C	403	-	X
4	PC1	C	404	-	X
4	PC1	D	403	-	X
4	PC1	D	404	-	X
4	PC1	E	403	-	X
4	PC1	E	404	-	X

## 2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 13358 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proton-gated ion channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	0	0
			2525	1664	404	453	4			
1	B	311	Total	C	N	O	S	0	0	0
			2525	1664	404	453	4			
1	C	311	Total	C	N	O	S	0	0	0
			2525	1664	404	453	4			
1	D	311	Total	C	N	O	S	0	0	0
			2525	1664	404	453	4			
1	E	311	Total	C	N	O	S	0	0	0
			2525	1664	404	453	4			

- Molecule 2 is (R)-KETAMINE (three-letter code: RKE) (formula: C<sub>13</sub>H<sub>16</sub>ClNO).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			16	13	1	1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	0	0
			16	13	1	1	1		
2	C	1	Total	C	Cl	N	O	0	0
			16	13	1	1	1		
2	D	1	Total	C	Cl	N	O	0	0
			16	13	1	1	1		
2	E	1	Total	C	Cl	N	O	0	0
			16	13	1	1	1		


- Molecule 3 is TETRADECYL 4-O-ALPHA-D-GLUCOPYRANOSYL-BETA-D-GLUCOPYRANOSIDE (three-letter code: LMD) (formula: C<sub>26</sub>H<sub>50</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			37	26	11		
3	B	1	Total	C	O	0	0
			37	26	11		
3	C	1	Total	C	O	0	0
			37	26	11		
3	C	1	Total	C	O	0	0
			37	26	11		
3	D	1	Total	C	O	0	0
			37	26	11		
3	E	1	Total	C	O	0	0
			37	26	11		

- Molecule 4 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1)

(formula: C<sub>44</sub>H<sub>88</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			37	27	1	8	1		
4	A	1	Total	C	N	O	P	0	0
			39	29	1	8	1		
4	B	1	Total	C	N	O	P	0	0
			39	29	1	8	1		
4	B	1	Total	C	N	O	P	0	0
			38	28	1	8	1		
4	C	1	Total	C	N	O	P	0	0
			38	28	1	8	1		
4	C	1	Total	C	N	O	P	0	0
			32	22	1	8	1		
4	D	1	Total	C	N	O	P	0	0
			37	27	1	8	1		
4	D	1	Total	C	N	O	P	0	0
			37	27	1	8	1		
4	E	1	Total	C	N	O	P	0	0
			38	28	1	8	1		
4	E	1	Total	C	N	O	P	0	0
			38	28	1	8	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	10	Total	O	0	0
			10	10		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	13	Total 13	O 13	0	0
5	C	13	Total 13	O 13	0	0
5	D	10	Total 10	O 10	0	0
5	E	12	Total 12	O 12	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proton-gated ion channel

Chain A:




- Molecule 1: Proton-gated ion channel

Chain B:



- Molecule 1: Proton-gated ion channel

Chain C:






GLY  
PHE


- Molecule 1: Proton-gated ion channel

Chain D:



- Molecule 1: Proton-gated ion channel

Chain E:



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	184.07Å 132.74Å 162.08Å 90.00° 103.61° 90.00°	Depositor
Resolution (Å)	24.84 – 2.99 25.00 – 2.99	Depositor EDS
% Data completeness (in resolution range)	89.8 (24.84-2.99) 97.7 (25.00-2.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 2.99Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, $R_{free}$	0.187 , 0.219 0.188 , 0.216	Depositor DCC
$R_{free}$ test set	3744 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.1	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 74569 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13358	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RKE, PC1, LMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/2593	0.57	0/3545
1	B	0.43	0/2593	0.59	0/3545
1	C	0.42	0/2593	0.57	0/3545
1	D	0.42	0/2593	0.57	0/3545
1	E	0.41	0/2593	0.57	0/3545
All	All	0.42	0/12965	0.57	0/17725

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2525	0	2545	80	2
1	B	2525	0	2545	82	0
1	C	2525	0	2545	82	0
1	D	2525	0	2545	74	2
1	E	2525	0	2545	78	0
2	A	16	0	16	6	0
2	B	16	0	16	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	16	0	16	5	0
2	D	16	0	16	5	0
2	E	16	0	16	6	0
3	A	37	0	50	7	0
3	B	37	0	50	6	0
3	C	74	0	100	15	0
3	D	37	0	50	6	0
3	E	37	0	50	6	0
4	A	76	0	100	7	0
4	B	77	0	102	7	0
4	C	70	0	88	7	0
4	D	74	0	96	6	0
4	E	76	0	100	10	0
5	A	10	0	0	2	0
5	B	13	0	0	6	0
5	C	13	0	0	1	0
5	D	10	0	0	0	0
5	E	12	0	0	2	0
All	All	13358	0	13591	426	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 426 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:404:PC1:C22	4:A:404:PC1:H31	1.68	1.13
4:A:404:PC1:H222	4:A:404:PC1:C3	1.80	1.11
4:A:404:PC1:H31	4:A:404:PC1:H222	1.00	0.99
2:D:401:RKE:HAK	2:D:401:RKE:CLAP	2.02	0.97
2:A:401:RKE:HAK	2:A:401:RKE:CLAP	2.01	0.97

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:173:ASN:O	1:D:139:ASN:ND2[4_555]	2.07	0.13
1:A:139:ASN:ND2	1:D:173:ASN:O[4_555]	2.14	0.06

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/317 (98%)	279 (90%)	26 (8%)	4 (1%)	18	62
1	B	309/317 (98%)	280 (91%)	26 (8%)	3 (1%)	22	70
1	C	309/317 (98%)	281 (91%)	24 (8%)	4 (1%)	18	62
1	D	309/317 (98%)	280 (91%)	26 (8%)	3 (1%)	22	70
1	E	309/317 (98%)	277 (90%)	29 (9%)	3 (1%)	22	70
All	All	1545/1585 (98%)	1397 (90%)	131 (8%)	17 (1%)	21	67

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	12	ALA
1	A	155	VAL
1	B	12	ALA
1	B	155	VAL
1	C	12	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/284 (99%)	257 (92%)	23 (8%)	17	52
1	B	280/284 (99%)	256 (91%)	24 (9%)	15	50
1	C	280/284 (99%)	256 (91%)	24 (9%)	15	50
1	D	280/284 (99%)	257 (92%)	23 (8%)	17	52
1	E	280/284 (99%)	257 (92%)	23 (8%)	17	52
All	All	1400/1420 (99%)	1283 (92%)	117 (8%)	16	51

5 of 117 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	63	VAL
1	C	229	VAL
1	E	181	GLU
1	C	93	SER
1	C	130	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	B	101	GLN
1	C	101	GLN
1	D	101	GLN
1	E	101	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RKE	A	401	-	17,17,17	1.44	2 (11%)	24,24,24	1.09	1 (4%)
3	LMD	A	402	-	38,38,38	0.92	1 (2%)	49,49,49	1.39	3 (6%)
4	PC1	A	403	-	36,36,53	3.04	8 (22%)	44,44,61	1.29	6 (13%)
4	PC1	A	404	-	38,38,53	3.00	6 (15%)	46,46,61	1.58	6 (13%)
2	RKE	B	401	-	17,17,17	1.40	2 (11%)	24,24,24	1.19	2 (8%)
3	LMD	B	402	-	38,38,38	0.96	1 (2%)	49,49,49	1.73	8 (16%)
4	PC1	B	403	-	38,38,53	3.04	7 (18%)	46,46,61	1.24	6 (13%)
4	PC1	B	404	-	37,37,53	2.96	5 (13%)	45,45,61	1.25	7 (15%)
2	RKE	C	401	-	17,17,17	1.41	2 (11%)	24,24,24	1.12	1 (4%)
3	LMD	C	402	-	38,38,38	0.96	1 (2%)	49,49,49	1.51	6 (12%)
4	PC1	C	403	-	37,37,53	3.30	7 (18%)	45,45,61	1.23	5 (11%)
4	PC1	C	404	-	31,31,53	3.15	7 (22%)	39,39,61	1.30	4 (10%)
3	LMD	C	405	-	38,38,38	0.95	1 (2%)	49,49,49	1.83	12 (24%)
2	RKE	D	401	-	17,17,17	1.42	2 (11%)	24,24,24	1.14	1 (4%)
3	LMD	D	402	-	38,38,38	0.94	1 (2%)	49,49,49	1.62	8 (16%)
4	PC1	D	403	-	36,36,53	2.99	7 (19%)	44,44,61	1.24	4 (9%)
4	PC1	D	404	-	36,36,53	3.12	7 (19%)	44,44,61	1.57	7 (15%)
2	RKE	E	401	-	17,17,17	1.37	2 (11%)	24,24,24	1.11	2 (8%)
3	LMD	E	402	-	38,38,38	1.00	1 (2%)	49,49,49	1.54	7 (14%)
4	PC1	E	403	-	37,37,53	3.00	8 (21%)	45,45,61	1.17	4 (8%)
4	PC1	E	404	-	37,37,53	3.08	7 (18%)	45,45,61	1.31	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RKE	A	401	-	-	1/8/23/23	0/2/2/2
3	LMD	A	402	-	-	0/23/63/63	0/2/2/2
4	PC1	A	403	-	-	0/40/40/57	0/0/0/0
4	PC1	A	404	-	-	2/42/42/57	0/0/0/0
2	RKE	B	401	-	-	1/8/23/23	0/2/2/2
3	LMD	B	402	-	-	0/23/63/63	0/2/2/2
4	PC1	B	403	-	-	0/42/42/57	0/0/0/0
4	PC1	B	404	-	-	0/41/41/57	0/0/0/0
2	RKE	C	401	-	-	1/8/23/23	0/2/2/2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LMD	C	402	-	-	0/23/63/63	0/2/2/2
4	PC1	C	403	-	-	0/41/41/57	0/0/0/0
4	PC1	C	404	-	-	0/35/35/57	0/0/0/0
3	LMD	C	405	-	-	0/23/63/63	0/2/2/2
2	RKE	D	401	-	-	1/8/23/23	0/2/2/2
3	LMD	D	402	-	-	0/23/63/63	0/2/2/2
4	PC1	D	403	-	-	0/40/40/57	0/0/0/0
4	PC1	D	404	-	-	0/40/40/57	0/0/0/0
2	RKE	E	401	-	-	1/8/23/23	0/2/2/2
3	LMD	E	402	-	-	0/23/63/63	0/2/2/2
4	PC1	E	403	-	-	0/41/41/57	0/0/0/0
4	PC1	E	404	-	-	0/41/41/57	0/0/0/0

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	403	PC1	C39-C38	-13.78	1.49	1.55
4	B	403	PC1	C3A-C39	-13.37	1.49	1.55
4	C	403	PC1	C2B-C2A	-12.85	1.50	1.55
4	D	403	PC1	C38-C37	-12.70	1.50	1.55
4	B	404	PC1	C39-C38	-12.64	1.50	1.55

The worst 5 of 106 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	LMD	OAS-CAM-CAN	7.70	117.98	108.18
4	A	404	PC1	O21-C21-C22	6.83	126.53	111.56
3	C	402	LMD	OAS-CAM-CAN	6.48	116.42	108.18
3	A	402	LMD	OAS-CAM-CAN	6.18	116.05	108.18
3	C	405	LMD	OAS-CAM-CAN	5.92	115.71	108.18

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	PC1	C2-O21-C21-C22
4	A	404	PC1	C2-O21-C21-O22
2	D	401	RKE	CAE-CAL-NAN-CAM
2	B	401	RKE	CAE-CAL-NAN-CAM
2	E	401	RKE	CAE-CAL-NAN-CAM

There are no ring outliers.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/317 (98%)	-0.64	2 (0%) 86 32	44, 65, 111, 145	0
1	B	311/317 (98%)	-0.52	6 (1%) 64 13	42, 65, 108, 141	0
1	C	311/317 (98%)	-0.55	6 (1%) 64 13	45, 65, 105, 143	0
1	D	311/317 (98%)	-0.53	4 (1%) 74 19	42, 65, 110, 139	0
1	E	311/317 (98%)	-0.53	7 (2%) 57 12	47, 65, 105, 142	0
All	All	1555/1585 (98%)	-0.56	25 (1%) 68 16	42, 65, 111, 145	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	57	VAL	6.3
1	E	58	ARG	4.6
1	C	58	ARG	4.3
1	B	136	ASP	3.8
1	D	57	VAL	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	RKE	E	401	16/16	0.34	13.01	42,118,164,200	16
2	RKE	A	401	16/16	0.37	10.35	47,112,178,184	0
2	RKE	B	401	16/16	0.40	9.20	37,117,176,199	0
4	PC1	B	403	39/54	0.34	9.15	45,100,209,235	0
3	LMD	E	402	37/37	0.41	8.66	45,127,203,222	0
3	LMD	A	402	37/37	0.36	8.58	51,113,204,215	0
2	RKE	C	401	16/16	0.34	8.52	37,123,165,203	0
4	PC1	C	403	38/54	0.35	8.39	44,105,192,228	0
2	RKE	D	401	16/16	0.60	8.30	39,117,178,207	16
3	LMD	D	402	37/37	0.41	8.03	42,126,193,218	0
4	PC1	E	404	38/54	0.44	7.45	64,108,207,242	0
4	PC1	D	403	37/54	0.40	6.76	46,108,195,207	0
4	PC1	A	404	39/54	0.37	6.62	43,103,212,243	0
4	PC1	B	404	38/54	0.32	6.35	44,112,183,209	0
3	LMD	C	402	37/37	0.40	6.31	44,128,217,218	0
4	PC1	E	403	38/54	0.33	5.95	43,95,189,215	0
4	PC1	C	404	32/54	0.38	5.94	57,123,201,258	0
3	LMD	B	402	37/37	0.35	5.44	38,141,200,211	0
4	PC1	A	403	37/54	0.32	5.11	29,95,207,223	0
3	LMD	C	405	37/37	0.26	4.65	43,102,189,217	0
4	PC1	D	404	37/54	0.28	3.64	54,100,193,258	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.