



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 09:08 AM GMT

PDB ID : 2F9Q
Title : Crystal Structure of Human Cytochrome P450 2D6
Authors : Rowland, P.
Deposited on : 2005-12-06
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	479	
1	B	479	
1	C	479	
1	D	479	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	HEM	A	600	-	X
3	HEM	C	600	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14421 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 2D6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	454	Total	C	N	O	S	0	0	0
			3569	2281	634	639	15			
1	B	451	Total	C	N	O	S	0	0	0
			3545	2265	630	635	15			
1	C	454	Total	C	N	O	S	0	0	0
			3569	2281	634	639	15			
1	D	451	Total	C	N	O	S	0	0	0
			3545	2265	630	635	15			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	HIS	CLONING ARTIFACT	UNP P10635
A	25	LYS	ARG	CLONING ARTIFACT	UNP P10635
A	26	LYS	ARG	CLONING ARTIFACT	UNP P10635
A	27	THR	GLN	CLONING ARTIFACT	UNP P10635
A	28	SER	ARG	CLONING ARTIFACT	UNP P10635
A	29	SER	TRP	CLONING ARTIFACT	UNP P10635
A	30	LYS	ALA	CLONING ARTIFACT	UNP P10635
A	31	GLY	ALA	CLONING ARTIFACT	UNP P10635
A	32	LYS	ARG	CLONING ARTIFACT	UNP P10635
A	33	LEU	TYR	CLONING ARTIFACT	UNP P10635
A	230	ASP	LEU	ENGINEERED	UNP P10635
A	231	ARG	LEU	ENGINEERED	UNP P10635
A	498	HIS	-	EXPRESSION TAG	UNP P10635
A	499	HIS	-	EXPRESSION TAG	UNP P10635
A	500	HIS	-	EXPRESSION TAG	UNP P10635
A	501	HIS	-	EXPRESSION TAG	UNP P10635
B	24	ALA	HIS	CLONING ARTIFACT	UNP P10635
B	25	LYS	ARG	CLONING ARTIFACT	UNP P10635
B	26	LYS	ARG	CLONING ARTIFACT	UNP P10635
B	27	THR	GLN	CLONING ARTIFACT	UNP P10635
B	28	SER	ARG	CLONING ARTIFACT	UNP P10635

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	29	SER	TRP	CLONING ARTIFACT	UNP P10635
B	30	LYS	ALA	CLONING ARTIFACT	UNP P10635
B	31	GLY	ALA	CLONING ARTIFACT	UNP P10635
B	32	LYS	ARG	CLONING ARTIFACT	UNP P10635
B	33	LEU	TYR	CLONING ARTIFACT	UNP P10635
B	230	ASP	LEU	ENGINEERED	UNP P10635
B	231	ARG	LEU	ENGINEERED	UNP P10635
B	498	HIS	-	EXPRESSION TAG	UNP P10635
B	499	HIS	-	EXPRESSION TAG	UNP P10635
B	500	HIS	-	EXPRESSION TAG	UNP P10635
B	501	HIS	-	EXPRESSION TAG	UNP P10635
C	24	ALA	HIS	CLONING ARTIFACT	UNP P10635
C	25	LYS	ARG	CLONING ARTIFACT	UNP P10635
C	26	LYS	ARG	CLONING ARTIFACT	UNP P10635
C	27	THR	GLN	CLONING ARTIFACT	UNP P10635
C	28	SER	ARG	CLONING ARTIFACT	UNP P10635
C	29	SER	TRP	CLONING ARTIFACT	UNP P10635
C	30	LYS	ALA	CLONING ARTIFACT	UNP P10635
C	31	GLY	ALA	CLONING ARTIFACT	UNP P10635
C	32	LYS	ARG	CLONING ARTIFACT	UNP P10635
C	33	LEU	TYR	CLONING ARTIFACT	UNP P10635
C	230	ASP	LEU	ENGINEERED	UNP P10635
C	231	ARG	LEU	ENGINEERED	UNP P10635
C	498	HIS	-	EXPRESSION TAG	UNP P10635
C	499	HIS	-	EXPRESSION TAG	UNP P10635
C	500	HIS	-	EXPRESSION TAG	UNP P10635
C	501	HIS	-	EXPRESSION TAG	UNP P10635
D	24	ALA	HIS	CLONING ARTIFACT	UNP P10635
D	25	LYS	ARG	CLONING ARTIFACT	UNP P10635
D	26	LYS	ARG	CLONING ARTIFACT	UNP P10635
D	27	THR	GLN	CLONING ARTIFACT	UNP P10635
D	28	SER	ARG	CLONING ARTIFACT	UNP P10635
D	29	SER	TRP	CLONING ARTIFACT	UNP P10635
D	30	LYS	ALA	CLONING ARTIFACT	UNP P10635
D	31	GLY	ALA	CLONING ARTIFACT	UNP P10635
D	32	LYS	ARG	CLONING ARTIFACT	UNP P10635
D	33	LEU	TYR	CLONING ARTIFACT	UNP P10635
D	230	ASP	LEU	ENGINEERED	UNP P10635
D	231	ARG	LEU	ENGINEERED	UNP P10635
D	498	HIS	-	EXPRESSION TAG	UNP P10635
D	499	HIS	-	EXPRESSION TAG	UNP P10635
D	500	HIS	-	EXPRESSION TAG	UNP P10635

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	501	HIS	-	EXPRESSION TAG	UNP P10635

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	3	Total	O	0	0
			3	3		
4	C	2	Total	O	0	0
			2	2		
4	D	2	Total	O	0	0
			2	2		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome P450 2D6

Chain A:




• Molecule 1: Cytochrome P450 2D6

Chain B:




• Molecule 1: Cytochrome P450 2D6

Chain C:



● Molecule 1: Cytochrome P450 2D6

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	145.07Å 155.50Å 95.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	106.00 – 3.00 39.97 – 3.00	Depositor EDS
% Data completeness (in resolution range)	89.6 (106.00-3.00) 89.6 (39.97-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.230 , 0.286 0.230 , 0.282	Depositor DCC
R_{free} test set	1601 reflections (4.23%)	DCC
Wilson B-factor (Å ²)	64.7	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 71.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 39438 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14421	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/3664	0.82	1/4982 (0.0%)
1	B	0.72	1/3639 (0.0%)	0.80	3/4947 (0.1%)
1	C	0.79	1/3664 (0.0%)	0.84	1/4982 (0.0%)
1	D	0.77	2/3639 (0.1%)	0.82	6/4947 (0.1%)
All	All	0.77	4/14606 (0.0%)	0.82	11/19858 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	1
1	C	0	2
1	D	0	2
All	All	0	9

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	269	ARG	CZ-NH1	15.49	1.53	1.33
1	C	57	CYS	CB-SG	-5.78	1.72	1.81
1	D	269	ARG	CZ-NH2	5.50	1.40	1.33
1	B	161	CYS	CB-SG	-5.42	1.73	1.81

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	269	ARG	NE-CZ-NH2	-13.90	113.35	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	441	ARG	NE-CZ-NH2	-7.01	116.79	120.30
1	D	441	ARG	NE-CZ-NH2	-6.71	116.95	120.30
1	D	388	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	D	234	PRO	N-CA-CB	6.54	111.15	103.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	113	GLY	Peptide
1	A	40	LEU	Peptide
1	A	53	ASN	Peptide
1	A	83	GLY	Peptide
1	B	113	GLY	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3569	0	3513	126	0
1	B	3545	0	3487	116	0
1	C	3569	0	3513	139	0
1	D	3545	0	3487	125	0
2	B	10	0	0	2	0
3	A	43	0	30	5	0
3	B	43	0	30	6	0
3	C	43	0	30	5	0
3	D	43	0	30	8	0
4	A	4	0	0	0	0
4	B	3	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
All	All	14421	0	14120	468	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 468 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:227:VAL:HB	1:C:227:VAL:CB	1.75	1.16
1:B:227:VAL:CB	1:C:227:VAL:HB	1.75	1.15
1:A:60:GLN:NE2	1:B:60:GLN:HE21	1.49	1.11
1:C:60:GLN:HE21	1:D:60:GLN:NE2	1.53	1.05
1:A:60:GLN:NE2	1:B:60:GLN:NE2	2.06	1.03

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/479 (94%)	396 (88%)	34 (8%)	20 (4%)	4	22
1	B	447/479 (93%)	397 (89%)	33 (7%)	17 (4%)	5	27
1	C	450/479 (94%)	397 (88%)	38 (8%)	15 (3%)	6	32
1	D	447/479 (93%)	395 (88%)	36 (8%)	16 (4%)	5	29
All	All	1794/1916 (94%)	1585 (88%)	141 (8%)	68 (4%)	5	27

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	PRO
1	A	40	LEU
1	A	114	PRO
1	A	145	GLY
1	A	229	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/409 (93%)	330 (87%)	50 (13%)	6	25
1	B	377/409 (92%)	322 (85%)	55 (15%)	5	21
1	C	380/409 (93%)	328 (86%)	52 (14%)	5	24
1	D	377/409 (92%)	322 (85%)	55 (15%)	5	21
All	All	1514/1636 (92%)	1302 (86%)	212 (14%)	5	23

5 of 212 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	395	LEU
1	C	144	LEU
1	D	343	ARG
1	B	477	HIS
1	C	63	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	477	HIS
1	C	250	GLN
1	D	350	GLN
1	C	52	GLN
1	C	175	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	A	600	1	49,50,50	2.32	17 (34%)	46,82,82	2.07	9 (19%)
2	SO4	B	2000	-	4,4,4	0.34	0	6,6,6	0.94	1 (16%)
2	SO4	B	2001	-	4,4,4	0.43	0	6,6,6	0.57	0
3	HEM	B	600	1	49,50,50	2.77	15 (30%)	46,82,82	1.63	8 (17%)
3	HEM	C	600	1	49,50,50	2.49	16 (32%)	46,82,82	1.53	8 (17%)
3	HEM	D	600	1	49,50,50	2.78	17 (34%)	46,82,82	1.52	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	600	1	-	0/14/114/114	0/0/8/8
2	SO4	B	2000	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2001	-	-	0/0/0/0	0/0/0/0
3	HEM	B	600	1	-	0/14/114/114	0/0/8/8
3	HEM	C	600	1	-	0/14/114/114	0/0/8/8
3	HEM	D	600	1	-	0/14/114/114	0/0/8/8

The worst 5 of 65 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	600	HEM	C2B-C1B	10.19	1.47	1.44
3	B	600	HEM	C2B-C1B	7.97	1.46	1.44
3	C	600	HEM	C2B-C1B	6.51	1.46	1.44
3	B	600	HEM	C2D-C1D	-5.73	1.43	1.44
3	B	600	HEM	C3D-C2D	5.66	1.53	1.43

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	600	HEM	C3B-C4B-NB	-7.60	108.56	114.00
3	A	600	HEM	C4D-ND-C1D	5.91	111.20	105.16
3	B	600	HEM	C4D-ND-C1D	5.37	110.65	105.16
3	D	600	HEM	C3B-C4B-NB	-5.13	110.33	114.00
3	B	600	HEM	C3B-C4B-NB	-4.72	110.62	114.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/479 (94%)	-0.08	0 100 100	50, 57, 63, 77	0
1	B	451/479 (94%)	-0.00	3 (0%) 84 28	46, 57, 63, 71	0
1	C	454/479 (94%)	-0.10	1 (0%) 93 54	49, 57, 63, 75	0
1	D	451/479 (94%)	-0.04	0 100 100	48, 57, 63, 72	0
All	All	1810/1916 (94%)	-0.06	4 (0%) 93 54	46, 57, 63, 77	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	286	PRO	2.4
1	C	228	PRO	2.3
1	B	228	PRO	2.2
1	B	424	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HEM	A	600	43/43	0.31	2.96	48,52,55,56	0
3	HEM	C	600	43/43	0.27	2.14	48,52,55,56	0
3	HEM	B	600	43/43	0.30	1.95	48,52,55,56	0
3	HEM	D	600	43/43	0.26	1.77	48,52,55,56	0
2	SO4	B	2001	5/5	0.28	1.50	52,54,57,58	0
2	SO4	B	2000	5/5	0.14	-2.25	46,51,54,54	0

6.5 Other polymers ⓘ

There are no such residues in this entry.