



wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 07:55 PM GMT

PDB ID : 2FAH
Title : The structure of mitochondrial PEPCK, Complex with Mn and GDP
Authors : Holyoak, T.; Sullivan, S.M.; Nowak, T.
Deposited on : 2005-12-07
Resolution : 2.09 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	608	
1	B	608	
1	C	608	
1	D	608	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	MN	A	2003	-	X
2	MN	C	4002	-	X
2	MN	D	5003	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 20770 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoenolpyruvate carboxykinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	608	Total	C	N	O	S	0	11	0
			4804	3048	865	858	33			
1	B	608	Total	C	N	O	S	0	13	0
			4786	3039	855	859	33			
1	C	608	Total	C	N	O	S	0	6	0
			4762	3021	857	851	33			
1	D	608	Total	C	N	O	S	0	5	0
			4749	3017	848	851	33			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	GLY	-	INSERTION	UNP P21642
A	131	PRO	SER	SEE REMARK 999	UNP P21642
A	233	PRO	ARG	SEE REMARK 999	UNP P21642
A	268	ARG	ALA	SEE REMARK 999	UNP P21642
A	339	GLU	ARG	SEE REMARK 999	UNP P21642
A	502	ARG	SER	SEE REMARK 999	UNP P21642
B	129	GLY	-	INSERTION	UNP P21642
B	131	PRO	SER	SEE REMARK 999	UNP P21642
B	233	PRO	ARG	SEE REMARK 999	UNP P21642
B	268	ARG	ALA	SEE REMARK 999	UNP P21642
B	339	GLU	ARG	SEE REMARK 999	UNP P21642
B	502	ARG	SER	SEE REMARK 999	UNP P21642
C	129	GLY	-	INSERTION	UNP P21642
C	131	PRO	SER	SEE REMARK 999	UNP P21642
C	233	PRO	ARG	SEE REMARK 999	UNP P21642
C	268	ARG	ALA	SEE REMARK 999	UNP P21642
C	339	GLU	ARG	SEE REMARK 999	UNP P21642
C	502	ARG	SER	SEE REMARK 999	UNP P21642
D	129	GLY	-	INSERTION	UNP P21642
D	131	PRO	SER	SEE REMARK 999	UNP P21642
D	233	PRO	ARG	SEE REMARK 999	UNP P21642

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	268	ARG	ALA	SEE REMARK 999	UNP P21642
D	339	GLU	ARG	SEE REMARK 999	UNP P21642
D	502	ARG	SER	SEE REMARK 999	UNP P21642

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mn	0	0
			2	2		
2	A	2	Total	Mn	0	0
			2	2		
2	D	2	Total	Mn	0	0
			2	2		
2	C	2	Total	Mn	0	0
			2	2		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	C	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	3	4		
4	B	1	Total	C	O	0	0
			7	3	4		
4	C	1	Total	C	O	0	0
			7	3	4		
4	D	1	Total	C	O	0	0
			7	3	4		
4	B	1	Total	C	O	0	0
			7	3	4		

- Molecule 5 is BETA-D-FRUCTOFURANOSYL6-O-OCTANOYL-ALPHA-D-GLUCOPYRANOSIDE (three-letter code: 20S) (formula: $C_{20}H_{36}O_{12}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			32	20	12		
5	C	1	Total	C	O	0	0
			32	20	12		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	384	Total	O	0	0
			384	384		
6	B	305	Total	O	0	0
			305	305		
6	C	417	Total	O	0	0
			417	417		
6	D	344	Total	O	0	0
			344	344		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphoenolpyruvate carboxykinase

Chain A:




- Molecule 1: Phosphoenolpyruvate carboxykinase

Chain B:




- Molecule 1: Phosphoenolpyruvate carboxykinase

Chain C:



● Molecule 1: Phosphoenolpyruvate carboxykinase

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	74.59Å 90.85Å 103.34Å 64.23° 73.74° 71.18°	Depositor
Resolution (Å)	32.81 – 2.09 32.80 – 2.09	Depositor EDS
% Data completeness (in resolution range)	94.7 (32.81-2.09) 94.7 (32.80-2.09)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.234 0.186 , 0.232	Depositor DCC
R_{free} test set	6402 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	24.8	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 127383 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20770	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, 20S, MLA, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/4975	0.64	0/6766
1	B	0.49	0/4969	0.62	1/6761 (0.0%)
1	C	0.52	0/4923	0.63	1/6697 (0.0%)
1	D	0.50	0/4905	0.61	1/6677 (0.0%)
All	All	0.51	0/19772	0.62	3/26901 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	502	ARG	NE-CZ-NH2	-5.86	117.37	120.30
1	C	639	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	D	168	PRO	C-N-CD	-5.58	108.33	120.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	169	PRO	Peptide
1	A	170	THR	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4804	0	4708	63	0
1	B	4786	0	4688	61	0
1	C	4762	0	4663	48	0
1	D	4749	0	4629	54	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	28	0	12	0	0
3	B	28	0	12	0	0
3	C	28	0	12	0	0
3	D	28	0	12	1	0
4	A	7	0	2	0	0
4	B	14	0	4	1	0
4	C	7	0	2	1	0
4	D	7	0	2	0	0
5	C	32	0	30	1	0
5	D	32	0	30	2	0
6	A	384	0	0	9	0
6	B	305	0	0	9	0
6	C	417	0	0	13	0
6	D	344	0	0	10	0
All	All	20770	0	18806	224	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 224 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:197:GLY:H	1:A:198:PRO:HD3	1.06	1.06
1:D:209:VAL:HG21	1:D:265:PHE:CE2	1.93	1.02
1:D:170:THR:HG22	6:D:6251:HOH:O	1.60	1.01
1:B:209:VAL:HG21	1:B:265:PHE:CE2	2.00	0.96
1:A:197:GLY:H	1:A:198:PRO:CD	1.81	0.93

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	617/608 (102%)	597 (97%)	14 (2%)	6 (1%)	22	14
1	B	619/608 (102%)	596 (96%)	19 (3%)	4 (1%)	33	28
1	C	612/608 (101%)	597 (98%)	14 (2%)	1 (0%)	56	57
1	D	611/608 (100%)	591 (97%)	17 (3%)	3 (0%)	38	33
All	All	2459/2432 (101%)	2381 (97%)	64 (3%)	14 (1%)	33	28

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	170	THR
1	B	207	ASP
1	D	168	PRO
1	A	255	GLY
1	B	255	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/499 (102%)	493 (97%)	16 (3%)	52	54
1	B	509/499 (102%)	489 (96%)	20 (4%)	43	43
1	C	503/499 (101%)	495 (98%)	8 (2%)	75	79
1	D	500/499 (100%)	488 (98%)	12 (2%)	61	65
All	All	2021/1996 (101%)	1965 (97%)	56 (3%)	60	59

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	203	ARG
1	B	403	THR
1	D	403	THR
1	B	205	ASP
1	B	276[A]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	311	ASN
1	C	135	ASN
1	D	311	ASN
1	B	524	ASN
1	C	202	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 8 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GDP	A	2000	2	30,30,30	1.56	5 (16%)	44,47,47	3.17	9 (20%)
4	MLA	A	2001	2	6,6,6	0.92	0	7,7,7	1.08	0
3	GDP	B	3000	2	30,30,30	1.42	5 (16%)	44,47,47	2.72	7 (15%)
4	MLA	B	3001	2	6,6,6	1.03	0	7,7,7	0.91	0
4	MLA	B	6001	-	6,6,6	1.04	0	7,7,7	0.71	0
3	GDP	C	4000	2	30,30,30	1.56	5 (16%)	44,47,47	3.50	9 (20%)
4	MLA	C	4003	2	6,6,6	1.10	0	7,7,7	1.24	1 (14%)
5	20S	C	6003	-	33,33,33	1.28	3 (9%)	46,46,46	1.95	4 (8%)
3	GDP	D	5000	2	30,30,30	1.63	6 (20%)	44,47,47	3.19	9 (20%)
4	MLA	D	5001	2	6,6,6	0.99	0	7,7,7	0.94	0
5	20S	D	6002	-	33,33,33	1.26	3 (9%)	46,46,46	1.95	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	2000	2	-	0/16/32/32	0/1/3/3
4	MLA	A	2001	2	-	0/4/4/4	0/0/0/0
3	GDP	B	3000	2	-	0/16/32/32	0/1/3/3
4	MLA	B	3001	2	-	0/4/4/4	0/0/0/0
4	MLA	B	6001	-	-	0/4/4/4	0/0/0/0
3	GDP	C	4000	2	-	0/16/32/32	0/1/3/3
4	MLA	C	4003	2	-	0/4/4/4	0/0/0/0
5	20S	C	6003	-	-	0/22/61/61	0/2/2/2
3	GDP	D	5000	2	-	0/16/32/32	0/1/3/3
4	MLA	D	5001	2	-	0/4/4/4	0/0/0/0
5	20S	D	6002	-	-	0/22/61/61	0/2/2/2

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	6003	20S	C11-C10	-5.23	1.19	1.51
5	D	6002	20S	C11-C10	-5.01	1.20	1.51
3	D	5000	GDP	C6-C5	4.75	1.49	1.41
3	C	4000	GDP	C6-C5	4.53	1.48	1.41
3	B	3000	GDP	C6-C5	3.99	1.47	1.41

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	4000	GDP	C6-C5-N7	20.15	136.85	134.14
3	D	5000	GDP	C6-C5-N7	17.81	136.54	134.14
3	A	2000	GDP	C6-C5-N7	17.63	136.51	134.14
3	B	3000	GDP	C6-C5-N7	13.35	135.94	134.14
5	C	6003	20S	C11-C10-C9	10.94	173.82	114.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	608/608 (100%)	0.06	16 (2%) 53 58	9, 15, 19, 25	0
1	B	608/608 (100%)	0.15	23 (3%) 38 43	11, 15, 19, 26	0
1	C	608/608 (100%)	0.08	18 (2%) 48 53	12, 15, 19, 25	0
1	D	608/608 (100%)	0.11	22 (3%) 41 45	11, 15, 19, 24	0
All	All	2432/2432 (100%)	0.10	79 (3%) 44 50	9, 15, 19, 26	0

The worst 5 of 79 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	398	PRO	6.4
1	C	396	THR	5.4
1	A	396	THR	5.3
1	B	400	VAL	5.2
1	B	222	THR	4.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MN	A	2003	1/1	0.35	13.64	21,21,21,21	1
2	MN	C	4002	1/1	0.29	6.84	9,9,9,9	1
2	MN	D	5003	1/1	0.29	3.44	14,14,14,14	1
4	MLA	A	2001	7/7	0.21	1.20	34,36,38,38	0
4	MLA	D	5001	7/7	0.22	0.73	27,30,32,32	0
4	MLA	B	3001	7/7	0.16	0.08	27,27,29,29	0
4	MLA	B	6001	7/7	0.14	-0.08	30,32,34,36	0
5	20S	D	6002	32/32	0.10	-0.15	11,15,19,20	0
4	MLA	C	4003	7/7	0.17	-0.27	21,21,26,30	0
5	20S	C	6003	32/32	0.09	-0.42	8,12,18,18	0
3	GDP	C	4000	28/28	0.11	-0.68	9,11,13,14	0
3	GDP	D	5000	28/28	0.09	-0.73	16,18,23,24	0
3	GDP	B	3000	28/28	0.09	-1.13	15,18,18,19	0
2	MN	B	3003	1/1	0.12	-1.16	23,23,23,23	1
3	GDP	A	2000	28/28	0.07	-1.74	9,11,14,15	0
2	MN	A	2002	1/1	0.08	-1.84	15,15,15,15	1
2	MN	C	4001	1/1	0.10	-1.99	9,9,9,9	0
2	MN	D	5002	1/1	0.10	-2.36	38,38,38,38	1
2	MN	B	3002	1/1	0.03	-4.60	28,28,28,28	1

6.5 Other polymers ⓘ

There are no such residues in this entry.