



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 02:56 PM GMT

PDB ID : 2FAT
Title : An anti-urokinase plasminogen activator receptor (UPAR) antibody: Crystal structure and binding epitope
Authors : Li, Y.; Parry, G.; Shi, X.; Chen, L.; Callahan, J.A.; Mazar, A.P.; Huang, M.
Deposited on : 2005-12-07
Resolution : 1.77 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6152 (1.80-1.76)
Ramachandran outliers	78287	6074 (1.80-1.76)
Sidechain outliers	78261	6073 (1.80-1.76)
RSRZ outliers	66119	4990 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	L	212	
2	H	214	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3439 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB ATN-615, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	212	Total	C	N	O	S	0	0	0
			1635	1023	269	335	8			

- Molecule 2 is a protein called FAB ATN-615, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	214	Total	C	N	O	S	0	0	0
			1626	1039	264	317	6			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	97	Total	O	0	0
			97	97		
3	L	81	Total	O	0	0
			81	81		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FAB ATN-615, light chain

Chain L: 



- Molecule 2: FAB ATN-615, heavy chain

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	37.16Å 84.47Å 134.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.29 – 1.77 40.28 – 1.77	Depositor EDS
% Data completeness (in resolution range)	93.5 (40.29-1.77) 97.7 (40.28-1.77)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.239 , 0.273 0.255 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.195	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 41076 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3439	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	0.32	0/1675	0.83	3/2275 (0.1%)
2	H	0.40	0/1677	0.90	11/2295 (0.5%)
All	All	0.36	0/3352	0.87	14/4570 (0.3%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	139	TYR	C-N-CD	-20.24	76.08	120.60
1	L	139	TYR	C-N-CA	13.73	179.66	122.00
2	H	130	MET	N-CA-C	-11.59	79.71	111.00
2	H	127	GLY	N-CA-C	11.01	140.64	113.10
2	H	126	PRO	N-CA-C	9.80	137.59	112.10
2	H	128	ASN	N-CA-C	8.77	134.69	111.00
2	H	56	ASN	N-CA-C	-7.67	90.29	111.00
2	H	53	GLY	N-CA-C	-6.36	97.20	113.10
1	L	140	PRO	CA-N-CD	-5.57	103.70	111.50
2	H	125	ALA	C-N-CD	-5.29	108.95	120.60
2	H	172	LEU	CA-CB-CG	5.23	127.32	115.30
2	H	135	CYS	CA-CB-SG	5.20	123.36	114.00
2	H	125	ALA	C-N-CA	5.09	143.39	122.00
2	H	60	ASN	N-CA-C	-5.01	97.46	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1635	0	1559	39	0
2	H	1626	0	1565	64	0
3	H	97	0	0	1	0
3	L	81	0	0	0	0
All	All	3439	0	3124	100	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

All (100) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:L:80:ALA:HA	1:L:106:ILE:HG21	1.53	0.90
1:L:159:LEU:HD11	2:H:166:GLN:HG3	1.57	0.84
1:L:194:GLU:HG2	1:L:205:VAL:HG22	1.60	0.84
2:H:12:VAL:HG21	2:H:82(C):LEU:HD12	1.62	0.80
1:L:154:ARG:NH1	1:L:180:LEU:HD21	1.98	0.77
2:H:24:ALA:HB1	2:H:27:TYR:CE2	2.21	0.76
2:H:126:PRO:HG3	2:H:133:LEU:HD23	1.68	0.75
1:L:139:TYR:CD2	1:L:140:PRO:HD3	2.22	0.74
1:L:141:LYS:NZ	1:L:162:TRP:HB3	2.03	0.74
2:H:30:THR:HA	2:H:52(A):HIS:CG	2.24	0.73
2:H:98:HIS:HB2	2:H:100:TYR:CE1	2.25	0.72
2:H:12:VAL:HG23	2:H:16:ALA:HB3	1.71	0.71
2:H:28:SER:O	2:H:30:THR:N	2.24	0.70
1:L:141:LYS:HZ1	1:L:162:TRP:HB3	1.57	0.70
2:H:6:GLN:H	2:H:105:GLN:NE2	1.90	0.69
2:H:105:GLN:CD	2:H:105:GLN:H	1.95	0.69
2:H:94:ARG:HB3	2:H:102:VAL:HG12	1.75	0.69
2:H:29:PHE:CD2	2:H:76:SER:HA	2.30	0.66
2:H:136:LEU:HD21	2:H:138:LYS:NZ	2.12	0.65
2:H:40:ARG:NH1	2:H:85:GLU:HG3	2.12	0.65
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.32	0.64
1:L:154:ARG:HH12	1:L:180:LEU:CD2	2.10	0.64
2:H:28:SER:HB2	2:H:31:ASN:OD1	1.98	0.63
1:L:154:ARG:HH12	1:L:180:LEU:HD21	1.61	0.63
2:H:27:TYR:CE1	2:H:29:PHE:HA	2.34	0.63
2:H:32:PHE:O	2:H:52(A):HIS:CE1	2.52	0.63
1:L:179:THR:O	1:L:180:LEU:HD23	1.99	0.63
2:H:97:PRO:HG2	2:H:98:HIS:ND1	2.14	0.62
1:L:16:GLY:HA2	1:L:77:SER:OG	1.98	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:L:166:ASP:OD1	1:L:168:LYS:HB2	2.00	0.62
1:L:46:PRO:HG3	2:H:100(A):PHE:O	1.99	0.61
1:L:34:HIS:HD2	1:L:50:GLU:H	1.48	0.61
2:H:98:HIS:HB2	2:H:100:TYR:HE1	1.65	0.61
1:L:149:ILE:HD11	1:L:178:LEU:HD21	1.83	0.61
1:L:146:LYS:HE2	1:L:153:GLU:CD	2.22	0.59
2:H:30:THR:HA	2:H:52(A):HIS:CD2	2.38	0.58
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.85	0.58
2:H:32:PHE:O	2:H:52(A):HIS:HE1	1.84	0.57
2:H:105:GLN:NE2	2:H:105:GLN:H	2.02	0.57
2:H:24:ALA:HB1	2:H:27:TYR:CZ	2.39	0.57
1:L:182:LYS:O	1:L:186:GLU:HG3	2.04	0.57
1:L:34:HIS:CD2	1:L:50:GLU:H	2.25	0.55
1:L:189:ASN:OD1	1:L:211:ASN:ND2	2.41	0.53
1:L:135:LEU:HD12	1:L:135:LEU:N	2.24	0.53
1:L:154:ARG:CZ	1:L:180:LEU:HD21	2.40	0.52
2:H:64:LYS:O	2:H:65:ASP:OD1	2.28	0.51
1:L:49:PHE:HD1	1:L:50:GLU:HG3	1.75	0.51
2:H:100(A):PHE:N	2:H:100(A):PHE:CD1	2.78	0.51
2:H:197:SER:O	2:H:199:THR:HG23	2.11	0.51
1:L:146:LYS:HE2	1:L:153:GLU:OE1	2.11	0.51
2:H:207:ALA:O	2:H:208:ALA:CB	2.59	0.51
2:H:6:GLN:HE21	2:H:104:GLY:HA3	1.76	0.51
1:L:154:ARG:HD3	1:L:178:LEU:HD11	1.92	0.50
1:L:197:HIS:HD2	1:L:199:THR:OG1	1.94	0.50
2:H:121:VAL:HG21	2:H:201:VAL:HG11	1.94	0.49
2:H:64:LYS:O	2:H:65:ASP:CB	2.60	0.49
2:H:136:LEU:HD21	2:H:138:LYS:HZ3	1.78	0.48
2:H:100(A):PHE:N	2:H:100(A):PHE:HD1	2.10	0.48
2:H:6:GLN:NE2	2:H:106:GLY:H	2.12	0.48
2:H:6:GLN:HE22	2:H:91:PHE:HA	1.78	0.47
1:L:141:LYS:HD3	1:L:172:TYR:CD1	2.48	0.47
2:H:13:LYS:H	2:H:13:LYS:HD2	1.79	0.47
1:L:148:LYS:HD2	1:L:194:GLU:OE2	2.15	0.47
1:L:180:LEU:HB3	1:L:184:GLU:HB2	1.97	0.47
1:L:198:LYS:HG3	1:L:199:THR:N	2.30	0.47
1:L:146:LYS:HE2	1:L:153:GLU:OE2	2.15	0.46
2:H:126:PRO:CG	2:H:133:LEU:HD23	2.42	0.46
2:H:172:LEU:C	2:H:172:LEU:HD23	2.35	0.46
2:H:29:PHE:CD2	2:H:52(A):HIS:CD2	3.04	0.46
1:L:141:LYS:HZ3	1:L:162:TRP:HB3	1.80	0.45
2:H:40:ARG:NH1	2:H:85:GLU:HA	2.31	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:121:VAL:HG21	2:H:201:VAL:CG1	2.45	0.45
2:H:119:PRO:HB3	2:H:140:TYR:HB3	1.97	0.45
2:H:12:VAL:CG2	2:H:82(C):LEU:HD12	2.41	0.45
1:L:114:VAL:HG12	1:L:206:LYS:HG3	1.99	0.44
2:H:136:LEU:HD21	2:H:138:LYS:HZ1	1.81	0.44
2:H:53:GLY:O	2:H:55:ASP:N	2.51	0.44
2:H:52(A):HIS:N	2:H:52(A):HIS:ND1	2.66	0.44
2:H:98:HIS:HB2	2:H:100:TYR:CD1	2.53	0.44
2:H:97:PRO:HG2	2:H:98:HIS:CE1	2.53	0.43
1:L:211:ASN:HB2	1:L:212:GLU:H	1.59	0.43
2:H:128:ASN:ND2	2:H:128:ASN:C	2.72	0.43
2:H:20:ILE:HG22	2:H:21:SER:N	2.33	0.43
1:L:192:THR:CG2	1:L:205:VAL:HG13	2.48	0.43
2:H:149:TRP:CZ3	2:H:190:CYS:HB3	2.53	0.43
2:H:66:LYS:O	2:H:82:LEU:HA	2.18	0.42
2:H:187:THR:HG23	2:H:204:LYS:HE3	2.02	0.42
2:H:50:TRP:CE2	2:H:58:GLU:HB3	2.54	0.42
2:H:94:ARG:HB3	2:H:102:VAL:CG1	2.48	0.42
2:H:11:VAL:HG21	2:H:142:PRO:HG3	2.03	0.41
1:L:198:LYS:C	1:L:200:SER:H	2.24	0.41
2:H:74:SER:HB2	3:H:290:HOH:O	2.21	0.41
2:H:47:TRP:CH2	2:H:49:GLY:HA2	2.56	0.41
1:L:161:SER:OG	2:H:162:PRO:HD2	2.21	0.41
1:L:208:PHE:HB2	1:L:213:CYS:CB	2.50	0.41
2:H:126:PRO:HB2	2:H:127:GLY:H	1.31	0.41
1:L:192:THR:HG23	1:L:207:SER:OG	2.21	0.40
2:H:201:VAL:HG22	2:H:202:ASP:N	2.36	0.40
1:L:209:ASN:H	1:L:213:CYS:HB2	1.86	0.40
2:H:125:ALA:HA	2:H:126:PRO:HD3	1.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	210/212 (99%)	199 (95%)	7 (3%)	4 (2%)	12	2
2	H	212/214 (99%)	194 (92%)	11 (5%)	7 (3%)	6	0
All	All	422/426 (99%)	393 (93%)	18 (4%)	11 (3%)	8	1

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	140	PRO
1	L	212	GLU
2	H	29	PHE
2	H	208	ALA
1	L	211	ASN
2	H	54	SER
2	H	126	PRO
2	H	127	GLY
1	L	167	SER
2	H	52(A)	HIS
2	H	28	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	186/186 (100%)	184 (99%)	2 (1%)	84	75
2	H	182/182 (100%)	173 (95%)	9 (5%)	35	13
All	All	368/368 (100%)	357 (97%)	11 (3%)	53	32

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	162	TRP
1	L	212	GLU
2	H	12	VAL
2	H	13	LYS
2	H	28	SER
2	H	65	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	105	GLN
2	H	126	PRO
2	H	144	PRO
2	H	172	LEU
2	H	191	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	34	HIS
1	L	137	ASN
1	L	197	HIS
1	L	211	ASN
2	H	6	GLN
2	H	56	ASN
2	H	105	GLN
2	H	128	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	212/212 (100%)	0.55	13 (6%) 21 18	16, 29, 56, 89	0
2	H	214/214 (100%)	0.85	26 (12%) 5 5	13, 26, 72, 110	0
All	All	426/426 (100%)	0.70	39 (9%) 9 8	13, 28, 59, 110	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	127	GLY	29.4
1	L	213	CYS	9.1
2	H	55	ASP	8.6
2	H	52(A)	HIS	8.4
2	H	128	ASN	7.9
2	H	29	PHE	7.3
2	H	129	SER	7.0
2	H	126	PRO	6.9
2	H	209	GLY	6.3
2	H	54	SER	5.4
2	H	27	TYR	4.8
2	H	53	GLY	4.2
2	H	28	SER	4.0
2	H	1	GLY	3.8
1	L	162	TRP	3.6
1	L	211	ASN	3.4
2	H	30	THR	3.2
2	H	2	VAL	3.2
1	L	156	ASN	3.2
1	L	154	ARG	2.9
2	H	208	ALA	2.9
2	H	99	TRP	2.7
1	L	159	LEU	2.6
2	H	31	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	202	SER	2.5
2	H	98	HIS	2.5
1	L	180	LEU	2.5
2	H	56	ASN	2.4
2	H	32	PHE	2.4
1	L	201	THR	2.3
1	L	155	GLN	2.3
2	H	125	ALA	2.3
2	H	76	SER	2.2
2	H	207	ALA	2.1
1	L	212	GLU	2.1
1	L	168	LYS	2.1
2	H	167	SER	2.1
2	H	71	ALA	2.0
1	L	140	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.