



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 12:57 AM GMT

PDB ID : 1FPB
Title : CRYSTAL STRUCTURE OF THE NEUTRAL FORM OF FRUCTOSE
1,6-BISPHOSPHATASE COMPLEXED WITH REGULATORY INHIBITOR
FRUCTOSE 2,6-BISPHOSPHATE AT 2.6-ANGSTROMS RESOLUTION
Authors : Liang, J.Y.; Huang, S.; Zhang, Y.; Ke, H.; Lipscomb, W.N.
Deposited on : 1992-02-03
Resolution : 2.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : NOT EXECUTED
EDS : NOT EXECUTED
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	335	
1	B	335	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 5924 atoms, of which 1066 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FRUCTOSE 1,6-BISPHOSPHATASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	314	Total	C	H	N	O	S	0	0	1
			2926	1523	530	404	454	15			
1	B	314	Total	C	H	N	O	S	0	0	1
			2932	1529	530	404	454	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLN	GLU	CONFLICT	UNP P00636
A	96	THR	SER	CONFLICT	UNP P00636
A	199	ASN	ASP	CONFLICT	UNP P00636
B	20	GLN	GLU	CONFLICT	UNP P00636
B	96	THR	SER	CONFLICT	UNP P00636
B	199	ASN	ASP	CONFLICT	UNP P00636

- Molecule 2 is SUGAR (FRUCTOSE-2,6-DIPHOSPHATE) (three-letter code: FDP) (formula: C₆H₁₄O₁₂P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	P	0	0
			22	6	2	12	2		
2	A	1	Total	C	H	O	P	0	0
			22	6	2	12	2		
2	B	1	Total	C	H	O	P	0	0
			22	6	2	12	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE

Chain A:



• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE

Chain B:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	131.40Å 131.40Å 68.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.60	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.174 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5924	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2435	1.29	12/3293 (0.4%)
1	B	0.67	0/2442	1.37	16/3302 (0.5%)
All	All	0.68	0/4877	1.33	28/6595 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	5
All	All	0	8

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	TYR	CB-CG-CD2	-8.75	115.75	121.00
1	B	313	ARG	NE-CZ-NH1	7.96	124.28	120.30
1	B	243	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	A	217	LYS	CA-CB-CG	7.54	129.99	113.40
1	B	313	ARG	NE-CZ-NH2	-7.01	116.80	120.30
1	B	333	LYS	CA-CB-CG	6.83	128.42	113.40
1	A	48	VAL	CG1-CB-CG2	-6.46	100.56	110.90
1	B	249	VAL	CG1-CB-CG2	-6.44	100.60	110.90
1	B	263	MET	CA-CB-CG	6.39	124.17	113.30
1	A	15	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	B	157	ARG	NE-CZ-NH1	6.37	123.48	120.30
1	B	27	THR	CA-C-N	-6.36	103.49	116.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	49	ARG	NE-CZ-NH1	6.21	123.41	120.30
1	A	49	ARG	NE-CZ-NH1	6.21	123.40	120.30
1	B	139	TYR	CB-CG-CD2	-6.00	117.40	121.00
1	A	244	TYR	CB-CG-CD1	-5.86	117.48	121.00
1	B	236	ASN	CA-C-N	-5.74	104.58	117.20
1	B	27	THR	N-CA-C	5.54	125.96	111.00
1	A	172	MET	CA-CB-CG	5.43	122.54	113.30
1	A	139	TYR	CB-CG-CD2	-5.42	117.75	121.00
1	B	48	VAL	CG1-CB-CG2	-5.27	102.47	110.90
1	A	157	ARG	NE-CZ-NH1	5.20	122.90	120.30
1	B	209	TYR	CB-CG-CD2	-5.19	117.89	121.00
1	A	287	VAL	CG1-CB-CG2	-5.17	102.63	110.90
1	A	76	LEU	CA-CB-CG	5.16	127.17	115.30
1	B	333	LYS	CB-CG-CD	-5.15	98.20	111.60
1	A	243	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	286	TYR	CB-CG-CD1	-5.03	117.98	121.00

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	139	TYR	Sidechain
1	A	240	TYR	Sidechain
1	A	264	TYR	Sidechain
1	B	139	TYR	Sidechain
1	B	167	TYR	Sidechain
1	B	244	TYR	Sidechain
1	B	258	TYR	Sidechain
1	B	264	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2396	530	1920	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2402	530	1927	36	0
2	A	40	4	16	0	0
2	B	20	2	8	0	0
All	All	4858	1066	3871	64	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (64) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:78:ASN:OD1	1:B:96:THR:HG21	1.82	0.79
1:B:182:ASN:HD21	1:B:198:ARG:HA	1.53	0.73
1:A:96:THR:HG22	1:A:98:GLU:H	1.56	0.70
1:B:126:ILE:HG12	1:B:132:ILE:HD13	1.73	0.69
1:B:182:ASN:ND2	1:B:198:ARG:HA	2.09	0.68
1:A:217:LYS:HB3	1:B:232:PHE:CE1	2.33	0.64
1:B:294:LEU:O	1:B:318:LEU:HA	1.99	0.62
1:A:155:PRO:HG3	1:A:306:VAL:HG22	1.79	0.62
1:A:297:THR:HB	1:A:315:PRO:HG2	1.82	0.61
1:B:43:ALA:HB3	1:B:80:LEU:HD21	1.82	0.60
1:A:73:LEU:HD23	1:A:126:ILE:HD13	1.83	0.60
1:A:31:THR:O	1:A:35:ASN:HB2	2.04	0.58
1:B:40:ALA:HB2	1:B:84:VAL:HG21	1.86	0.58
1:A:166:LEU:HD13	1:A:249:VAL:HG12	1.86	0.58
1:B:31:THR:O	1:B:35:ASN:HB2	2.04	0.57
1:B:282:ASN:ND2	1:B:302:VAL:HG12	2.19	0.56
1:A:166:LEU:HD23	1:B:129:LEU:HD12	1.87	0.55
1:B:96:THR:HG22	1:B:98:GLU:H	1.73	0.54
1:A:13:LEU:HD22	1:A:38:CYS:SG	2.48	0.54
1:B:138:ILE:O	1:B:160:VAL:HG22	2.09	0.53
1:B:297:THR:HG21	1:B:305:ILE:HD11	1.91	0.53
1:B:252:VAL:HG11	1:B:284:MET:SD	2.49	0.52
1:B:114:VAL:HG11	1:B:152:ALA:HA	1.90	0.52
1:B:178:VAL:HA	1:B:290:LYS:HE2	1.92	0.52
1:A:78:ASN:OD1	1:A:96:THR:HG21	2.11	0.50
1:B:150:LYS:HD3	1:B:153:LEU:HD12	1.94	0.49
1:A:296:THR:HG21	1:A:328:LEU:HD21	1.94	0.48
1:A:252:VAL:HG22	1:A:318:LEU:HD11	1.96	0.47
1:B:310:ILE:HG13	1:B:311:HIS:CD2	2.49	0.47
1:A:283:PRO:O	1:A:287:VAL:HG23	2.15	0.46
1:A:96:THR:HG22	1:A:97:GLU:N	2.31	0.46
1:B:182:ASN:HD22	1:B:198:ARG:HG2	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:302:VAL:HG21	1:A:316:ILE:HD13	1.97	0.46
1:B:96:THR:HG22	1:B:97:GLU:N	2.30	0.46
1:A:93:VAL:HB	1:A:114:VAL:HG13	1.97	0.46
1:B:276:ARG:HD3	1:B:313:ARG:HG2	1.96	0.46
1:B:182:ASN:HD22	1:B:199:ASN:H	1.63	0.45
1:B:73:LEU:HD23	1:B:126:ILE:HD13	1.98	0.45
1:A:288:MET:HG3	1:A:318:LEU:HB2	1.97	0.45
1:B:114:VAL:HB	1:B:139:TYR:HB2	1.98	0.44
1:B:211:ILE:HD13	1:B:227:ILE:HD11	1.99	0.44
1:B:267:ASN:O	1:B:271:PRO:HA	2.17	0.44
1:A:82:ILE:O	1:A:86:LYS:HG3	2.18	0.44
1:A:166:LEU:O	1:A:171:THR:HA	2.18	0.43
1:B:306:VAL:HA	1:B:307:PRO:HD3	1.79	0.43
1:A:40:ALA:HB2	1:A:84:VAL:HG21	2.00	0.43
1:A:317:ILE:HG21	1:A:327:LEU:HD23	2.00	0.43
1:A:288:MET:HE2	1:A:288:MET:HB3	1.83	0.42
1:B:182:ASN:ND2	1:B:199:ASN:H	2.17	0.42
1:A:131:SER:HB2	1:A:250:ALA:HB2	1.99	0.42
1:A:172:MET:SD	1:A:183:CYS:HB3	2.59	0.42
1:A:89:PHE:HD1	1:A:109:LYS:HA	1.84	0.42
1:A:233:PRO:HA	1:A:234:PRO:HD2	1.76	0.42
1:B:296:THR:HG21	1:B:328:LEU:HD21	2.00	0.42
1:B:27:THR:HB	1:B:30:MET:H	1.84	0.42
1:B:27:THR:HG21	1:B:113:TYR:OH	2.20	0.41
1:B:93:VAL:HB	1:B:114:VAL:HG22	2.02	0.41
1:A:13:LEU:CD2	1:A:38:CYS:SG	3.08	0.41
1:A:214:GLY:HA3	1:B:240:TYR:O	2.21	0.41
1:A:174:VAL:HG22	1:A:183:CYS:SG	2.60	0.41
1:A:333:LYS:HG3	1:A:334:HIS:N	2.35	0.41
1:A:96:THR:HG22	1:A:98:GLU:N	2.30	0.41
1:B:85:LEU:HA	1:B:85:LEU:HD23	1.79	0.41
1:B:106:GLU:O	1:B:110:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/335 (92%)	298 (96%)	11 (4%)	1 (0%)	50	77
1	B	310/335 (92%)	289 (93%)	19 (6%)	2 (1%)	33	63
All	All	620/670 (92%)	587 (95%)	30 (5%)	3 (0%)	38	67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	ASN
1	B	236	ASN
1	B	199	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/278 (94%)	252 (97%)	9 (3%)	49	78
1	B	262/278 (94%)	253 (97%)	9 (3%)	49	78
All	All	523/556 (94%)	505 (97%)	18 (3%)	49	78

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ILE
1	A	25	ARG
1	A	125	ASN
1	A	197	ASP
1	A	230	LYS
1	A	265	PRO
1	A	268	LYS
1	A	276	ARG
1	A	300	GLU
1	B	6	PHE
1	B	19	GLU
1	B	127	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	146	GLU
1	B	199	ASN
1	B	235	ASP
1	B	265	PRO
1	B	268	LYS
1	B	308	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	154	GLN
1	A	158	ASN
1	A	182	ASN
1	A	282	ASN
1	B	154	GLN
1	B	182	ASN
1	B	199	ASN
1	B	267	ASN
1	B	282	ASN
1	B	334	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FDP	A	336	-	20,20,20	0.52	0	32,32,32	0.82	0
2	FDP	A	337	-	20,20,20	0.54	0	32,32,32	0.92	0
2	FDP	B	336	-	20,20,20	0.46	0	32,32,32	1.04	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDP	A	336	-	-	0/12/34/34	0/1/1/1
2	FDP	A	337	-	-	0/12/34/34	0/1/1/1
2	FDP	B	336	-	-	0/12/34/34	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	336	FDP	O3P-P1-O2	2.74	114.99	107.09
2	B	336	FDP	O5P-P2-O6	2.05	112.31	106.65
2	B	336	FDP	O1-C1-C2	2.03	118.37	111.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.