



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 12:56 AM GMT

PDB ID : 1FPT  
Title : THREE-DIMENSIONAL STRUCTURE OF THE COMPLEX BETWEEN  
THE FAB FRAGMENT OF AN NEUTRALIZING ANTIBODY FOR TYPE  
1 POLIOVIRUS AND ITS VIRAL EPITOPE  
Authors : Wien, M.W.; Hogle, J.M.  
Deposited on : 1995-01-26  
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	P	18	
2	L	219	
3	H	220	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3465 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB FRAGMENT OF AN NEUTRALIZING ANTIBODY FOR TYPE 1 POLIOVIRUS.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	P	11	83	47	15	21	0	0	0

- Molecule 2 is a protein called IGG2A-KAPPA C3 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	219	1691	1061	283	341	6	0	0	0

- Molecule 3 is a protein called IGG2A-KAPPA C3 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	H	220	1652	1045	269	332	6	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	20	Total 20	O 20	0	0
4	L	17	Total 17	O 17	0	0
4	P	2	Total 2	O 2	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: FAB FRAGMENT OF AN NEUTRALIZING ANTIBODY FOR TYPE 1 POLIOVIRUS

Chain P: 




- Molecule 2: IGG2A-KAPPA C3 FAB (LIGHT CHAIN)

Chain L: 



- Molecule 3: IGG2A-KAPPA C3 FAB (HEAVY CHAIN)

Chain H: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	129.78Å 129.78Å 143.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-3.00)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.230 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3465	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	P	1.15	0/83	2.61	9/110 (8.2%)
2	L	1.11	1/1731 (0.1%)	1.96	48/2351 (2.0%)
3	H	1.10	2/1693 (0.1%)	2.07	56/2314 (2.4%)
All	All	1.11	3/3507 (0.1%)	2.03	113/4775 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	P	0	1
2	L	0	1
3	H	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	186	SER	CA-CB	-5.72	1.44	1.52
2	L	35	TRP	CG-CD2	-5.70	1.33	1.43
3	H	196	TRP	CG-CD2	-5.02	1.35	1.43

The worst 5 of 113 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	142	CYS	CA-CB-SG	-11.87	92.64	114.00
3	H	99	TYR	CB-CG-CD1	-11.30	114.22	121.00
2	L	211	ARG	NE-CZ-NH1	10.43	125.51	120.30
3	H	47	TRP	CD1-CG-CD2	10.11	114.38	106.30
3	H	36	TRP	CD1-CG-CD2	9.13	113.61	106.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	40	ARG	Peptide
3	H	59	TYR	Sidechain
2	L	32	TYR	Sidechain
1	P	94	ASN	Mainchain

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	83	0	76	1	0
2	L	1691	0	1636	57	0
3	H	1652	0	1609	49	0
4	H	20	0	0	0	0
4	L	17	0	0	0	0
4	P	2	0	0	0	0
All	All	3465	0	3321	101	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:L:27(D):HIS:HB2	2:L:92:THR:HG23	1.38	1.04
2:L:147:LYS:HE3	2:L:154:GLU:HG2	1.51	0.93
2:L:113:PRO:HG3	2:L:144:ILE:HD11	1.61	0.80
2:L:118:PHE:CZ	3:H:140:LEU:HA	2.22	0.73
2:L:112:ALA:HB2	2:L:200:THR:HG21	1.70	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone i

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	9/18 (50%)	6 (67%)	1 (11%)	2 (22%)	0	0
2	L	217/219 (99%)	189 (87%)	20 (9%)	8 (4%)	5	28
3	H	218/220 (99%)	189 (87%)	20 (9%)	9 (4%)	4	24
All	All	444/457 (97%)	384 (86%)	41 (9%)	19 (4%)	4	23

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	27(E)	SER
2	L	68	GLY
2	L	211	ARG
3	H	135	GLY
3	H	193	SER

### 5.3.2 Protein sidechains i

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	10/17 (59%)	8 (80%)	2 (20%)	2	10
2	L	196/197 (100%)	179 (91%)	17 (9%)	15	49
3	H	187/187 (100%)	155 (83%)	32 (17%)	3	15
All	All	393/401 (98%)	342 (87%)	51 (13%)	6	26

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	50	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	H	101	ASP
3	H	191	VAL
3	H	68	THR
3	H	105	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	210	ASN
3	H	212	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

### 5.7 Other polymers ⓘ

There are no such residues in this entry.

### 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers

EDS was not executed - this section will therefore be empty.