



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 02:03 PM GMT

PDB ID : 3FP3  
Title : Crystal structure of Tom71  
Authors : Li, J.; Qian, X.; Hu, J.; Sha, B.  
Deposited on : 2009-01-03  
Resolution : 1.98 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	6577 (2.00-1.96)
Clashscore	79885	8091 (2.00-1.96)
Ramachandran outliers	78287	7989 (2.00-1.96)
Sidechain outliers	78261	7987 (2.00-1.96)
RSRZ outliers	66119	6578 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	537	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	A	3	-	X
2	SO4	A	4	-	X
3	CL	A	1	-	X
3	CL	A	640	-	X
3	CL	A	641	-	X
3	CL	A	642	-	X

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4669 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TPR repeat-containing protein YHR117W.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	498	4108	2615	671	808	14	0	19	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	GLY	-	EXPRESSION TAG	UNP P38825
A	104	SER	-	EXPRESSION TAG	UNP P38825
A	105	HIS	-	EXPRESSION TAG	UNP P38825
A	106	MET	-	EXPRESSION TAG	UNP P38825

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Cl	0	0
			4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	542	Total	O	0	0
			542	542		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TPR repeat-containing protein YHR117W



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.90Å 83.42Å 109.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.63 – 1.98 19.63 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.63-1.98) 99.9 (19.63-1.98)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.79 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.187 , 0.227 0.242 , 0.273	Depositor DCC
$R_{free}$ test set	2508 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 29.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 49516 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4669	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	0/4184	0.75	7/5656 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	MET	CG-SD-CE	-9.59	84.86	100.20
1	A	201	ARG	NE-CZ-NH2	-8.96	115.82	120.30
1	A	201	ARG	NE-CZ-NH1	8.68	124.64	120.30
1	A	337	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	238	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	319	LEU	CB-CG-CD1	5.22	119.87	111.00
1	A	235	MET	CB-CG-SD	5.04	127.52	112.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4108	0	4043	59	0
2	A	15	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	0	5	0
4	A	542	0	0	13	0
All	All	4669	0	4043	60	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (60) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:578:GLU:O	1:A:578:GLU:HG3	1.68	0.94
1:A:245[B]:MET:HG3	3:A:641:CL:CL	2.16	0.81
1:A:135:ASN:O	1:A:138:PHE:HB2	1.82	0.80
1:A:215[A]:MET:HE3	1:A:236:LEU:O	1.81	0.79
1:A:201:ARG:HD2	1:A:217:ASP:OD2	1.85	0.77
1:A:365:GLN:HG3	1:A:366:VAL:N	2.00	0.76
1:A:201:ARG:CD	1:A:217:ASP:OD2	2.37	0.72
1:A:533:GLN:O	1:A:534:SER:HB2	1.93	0.67
1:A:391:LYS:H	1:A:391:LYS:HE2	1.60	0.65
1:A:112:ILE:HG13	1:A:152:TYR:CD1	2.32	0.64
1:A:383:PHE:O	1:A:387[B]:THR:HG23	1.97	0.64
1:A:420:GLN:HG2	4:A:910:HOH:O	2.00	0.62
1:A:139:THR:HA	4:A:834:HOH:O	1.99	0.61
1:A:109:GLU:N	4:A:825:HOH:O	2.34	0.61
1:A:138:PHE:O	1:A:139:THR:C	2.39	0.60
1:A:147:ILE:HG23	1:A:167[B]:ILE:HD11	1.85	0.58
1:A:399:LYS:HE3	4:A:899:HOH:O	2.05	0.56
1:A:391:LYS:N	1:A:391:LYS:HE2	2.21	0.55
1:A:115:LEU:HD11	3:A:1:CL:CL	2.44	0.54
1:A:150:TYR:CB	1:A:167[A]:ILE:HG13	2.39	0.53
1:A:284:SER:HB3	4:A:872:HOH:O	2.07	0.53
1:A:177:LEU:HD23	4:A:842:HOH:O	2.07	0.53
1:A:201:ARG:HD3	1:A:217:ASP:OD2	2.07	0.52
1:A:365:GLN:HB3	4:A:883:HOH:O	2.09	0.52
1:A:205:ASN:HD22	1:A:210:ASN:HD22	1.56	0.52
1:A:136:HIS:C	1:A:138:PHE:H	2.13	0.51
1:A:215[A]:MET:CE	1:A:240:LEU:HB2	2.41	0.51
1:A:565[B]:GLN:HE21	1:A:565[B]:GLN:H	1.59	0.51
1:A:365:GLN:CB	4:A:883:HOH:O	2.59	0.50
1:A:475:LYS:HE2	4:A:746:HOH:O	2.11	0.50
1:A:365:GLN:HG3	1:A:366:VAL:H	1.75	0.50
1:A:266:ASN:ND2	1:A:562:ARG:HH22	2.11	0.49
1:A:157:ASP:O	3:A:640:CL:CL	2.68	0.49

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:335:THR:HG22	4:A:763:HOH:O	2.11	0.49
1:A:215[A]:MET:HE1	1:A:240:LEU:HB2	1.94	0.49
1:A:508:ARG:N	1:A:508:ARG:HD2	2.29	0.47
1:A:297:LEU:HD11	1:A:319:LEU:HG	1.95	0.47
1:A:136:HIS:C	1:A:138:PHE:N	2.68	0.47
1:A:578:GLU:HB2	4:A:1073:HOH:O	2.14	0.47
1:A:150:TYR:HB2	1:A:167[A]:ILE:HG13	1.98	0.45
1:A:136:HIS:O	1:A:138:PHE:N	2.50	0.45
1:A:545:LYS:HB3	1:A:545:LYS:HE3	1.69	0.44
1:A:365:GLN:HE21	1:A:365:GLN:C	2.21	0.44
3:A:642:CL:CL	4:A:784:HOH:O	2.59	0.44
1:A:154:ILE:HD12	1:A:167[A]:ILE:HD12	2.00	0.43
1:A:211:PHE:O	1:A:215[B]:MET:HG3	2.19	0.43
1:A:391:LYS:H	1:A:391:LYS:CE	2.29	0.43
1:A:196:LYS:HE2	1:A:196:LYS:HB2	1.76	0.43
1:A:504:ASP:O	1:A:508:ARG:CD	2.66	0.43
1:A:565[B]:GLN:H	1:A:565[B]:GLN:NE2	2.16	0.43
1:A:417:HIS:CE1	4:A:667:HOH:O	2.71	0.43
1:A:215[A]:MET:HE1	2:A:4:SO4:O1	2.18	0.43
1:A:215[A]:MET:CE	2:A:4:SO4:O1	2.67	0.43
1:A:578:GLU:O	1:A:578:GLU:CG	2.50	0.42
1:A:601:LEU:HD11	3:A:641:CL:CL	2.57	0.42
1:A:394:SER:HG	1:A:398:PHE:HE2	1.63	0.41
1:A:337:ASP:HB3	1:A:339:PRO:HD2	2.02	0.41
1:A:572[A]:GLN:HA	1:A:572[A]:GLN:OE1	2.21	0.41
1:A:504:ASP:O	1:A:508:ARG:HD2	2.21	0.41
1:A:205:ASN:ND2	1:A:210:ASN:HD22	2.18	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/537 (95%)	504 (99%)	5 (1%)	1 (0%)	56	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	137	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	452/463 (98%)	432 (96%)	20 (4%)	39	30

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	121	SER
1	A	131	LYS
1	A	138	PHE
1	A	139	THR
1	A	201	ARG
1	A	205	ASN
1	A	235	MET
1	A	322	LYS
1	A	334	ASN
1	A	337	ASP
1	A	365	GLN
1	A	391	LYS
1	A	393	ASN
1	A	394	SER
1	A	423	PHE
1	A	508	ARG
1	A	532	ARG
1	A	544	GLU
1	A	545	LYS
1	A	595	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	249	ASN
1	A	266	ASN
1	A	365	GLN
1	A	375	HIS
1	A	393	ASN
1	A	417	HIS
1	A	430	ASN
1	A	502	GLN
1	A	547	ASN
1	A	613	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	2	-	4,4,4	0.17	0	6,6,6	0.12	0
2	SO4	A	3	-	4,4,4	0.15	0	6,6,6	0.09	0
2	SO4	A	4	-	4,4,4	0.20	0	6,6,6	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	2	-	-	0/0/0/0	0/0/0/0
2	SO4	A	3	-	-	0/0/0/0	0/0/0/0
2	SO4	A	4	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	498/537 (92%)	0.79	38 (7%) 14 14	15, 33, 49, 64	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	PHE	7.2
1	A	136	HIS	6.6
1	A	334	ASN	6.2
1	A	139	THR	4.7
1	A	392	GLU	4.6
1	A	137	PHE	4.2
1	A	120	PRO	3.7
1	A	335	THR	3.7
1	A	452	LEU	3.2
1	A	391	LYS	2.7
1	A	578	GLU	2.6
1	A	109	GLU	2.6
1	A	261	GLN	2.6
1	A	112	ILE	2.6
1	A	254	LYS	2.5
1	A	544	GLU	2.5
1	A	414	THR	2.5
1	A	305	LEU	2.5
1	A	340[A]	LEU	2.5
1	A	435	PHE	2.5
1	A	520	ILE	2.5
1	A	382	ILE	2.4
1	A	385	ALA	2.4
1	A	149	TYR	2.3
1	A	497	ASP	2.3
1	A	349	CYS	2.3
1	A	448	PRO	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	532	ARG	2.2
1	A	123	ARG	2.2
1	A	116	LYS	2.2
1	A	453	ALA	2.1
1	A	167[A]	ILE	2.1
1	A	627	GLU	2.1
1	A	345	ALA	2.1
1	A	413	PRO	2.1
1	A	291	TYR	2.1
1	A	343	ASN	2.1
1	A	529	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
2	SO4	A	3	5/5	0.65	24.54	161,161,161,161	0
3	CL	A	640	1/1	0.32	9.37	53,53,53,53	0
3	CL	A	641	1/1	0.42	8.29	44,44,44,44	0
3	CL	A	1	1/1	0.29	6.87	58,58,58,58	0
2	SO4	A	4	5/5	0.34	5.83	62,65,66,66	0
3	CL	A	642	1/1	0.27	2.62	51,51,51,51	0
2	SO4	A	2	5/5	0.21	0.94	59,60,61,61	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.