



wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 04:35 PM GMT

PDB ID : 3FVQ
Title : Crystal structure of the nucleotide binding domain FbpC complexed with ATP
Authors : Newstead, S.; Bilton, P.; Carpenter, E.P.; Campopiano, D.; Iwata, S.
Deposited on : 2009-01-16
Resolution : 1.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	359	
1	B	359	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5963 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fe(3+) ions import ATP-binding protein fbpC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	0	10	0
			2653	1677	477	492	7			
1	B	350	Total	C	N	O	S	0	13	0
			2723	1719	494	504	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	ALA	VAL	SEE REMARK 999	UNP Q5FA19
A	283	TYR	PHE	SEE REMARK 999	UNP Q5FA19
A	353	GLU	-	EXPRESSION TAG	UNP Q5FA19
A	354	HIS	-	EXPRESSION TAG	UNP Q5FA19
A	355	HIS	-	EXPRESSION TAG	UNP Q5FA19
A	356	HIS	-	EXPRESSION TAG	UNP Q5FA19
A	357	HIS	-	EXPRESSION TAG	UNP Q5FA19
A	358	HIS	-	EXPRESSION TAG	UNP Q5FA19
A	359	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	147	ALA	VAL	SEE REMARK 999	UNP Q5FA19
B	283	TYR	PHE	SEE REMARK 999	UNP Q5FA19
B	353	GLU	-	EXPRESSION TAG	UNP Q5FA19
B	354	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	355	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	356	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	357	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	358	HIS	-	EXPRESSION TAG	UNP Q5FA19
B	359	HIS	-	EXPRESSION TAG	UNP Q5FA19

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	243	Total	O	0	0
			243	243		
4	B	279	Total	O	0	0
			279	279		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fe(3+) ions import ATP-binding protein fbpC

Chain A: 



- Molecule 1: Fe(3+) ions import ATP-binding protein fbpC

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.86Å 89.09Å 149.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.27 – 1.90 47.33 – 1.90	Depositor EDS
% Data completeness (in resolution range)	87.4 (52.27-1.90) 87.4 (47.33-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.195 , 0.256 0.186 , 0.234	Depositor DCC
R_{free} test set	2624 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 51946 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5963	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.69	0/2731	0.78	1/3702 (0.0%)
1	B	0.78	2/2813 (0.1%)	0.81	2/3819 (0.1%)
All	All	0.74	2/5544 (0.0%)	0.79	3/7521 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	38	ALA	CA-CB	5.72	1.64	1.52
1	B	195	VAL	CB-CG1	5.31	1.64	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	232	ASP	CB-CG-OD1	-5.54	113.31	118.30
1	B	232	ASP	CB-CA-C	-5.19	100.02	110.40
1	A	48	ARG	NE-CZ-NH2	-5.11	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2653	0	2711	99	0
1	B	2723	0	2796	110	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	243	0	0	21	0
4	B	279	0	0	13	0
All	All	5963	0	5531	195	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

The worst 5 of 195 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:276:LEU:HB3	4:A:448:HOH:O	1.39	1.22
1:A:242:GLY:HA3	1:A:277:LEU:HD11	1.30	1.07
1:B:178:ARG:HD2	4:B:448:HOH:O	1.58	1.03
1:B:5:LEU:HD11	1:B:50[B]:LEU:HD11	1.38	1.01
1:B:145:GLN:HE22	1:B:168:ALA:H	1.10	0.98

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/359 (97%)	335 (96%)	11 (3%)	3 (1%)	25	10
1	B	361/359 (101%)	348 (96%)	11 (3%)	2 (1%)	33	19
All	All	710/718 (99%)	683 (96%)	22 (3%)	5 (1%)	30	15

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	350	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	90	VAL
1	A	268	ALA
1	A	333	GLY
1	B	331	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/288 (98%)	267 (95%)	15 (5%)	32	18
1	B	291/288 (101%)	278 (96%)	13 (4%)	38	24
All	All	573/576 (100%)	545 (95%)	28 (5%)	35	21

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	302	THR
1	B	15	GLN
1	B	320	LEU
1	A	318	THR
1	A	334	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	HIS
1	B	15	GLN
1	B	219	GLN
1	A	219	GLN
1	B	286	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ATP	A	401	3	33,33,33	1.36	4 (12%)	52,52,52	1.93	12 (23%)
2	ATP	B	360	3	33,33,33	1.32	4 (12%)	52,52,52	1.90	9 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	401	3	-	0/22/38/38	0/1/3/3
2	ATP	B	360	3	-	0/22/38/38	0/1/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	ATP	C4-N9	-3.57	1.32	1.37
2	B	360	ATP	PG-O3B	3.32	1.66	1.60
2	B	360	ATP	C4-N9	-3.27	1.33	1.37
2	B	360	ATP	PB-O3A	3.14	1.65	1.59
2	A	401	ATP	PB-O3A	2.89	1.65	1.59

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	360	ATP	N3-C2-N1	-8.18	121.87	128.71
2	A	401	ATP	N3-C2-N1	-8.07	121.96	128.71
2	B	360	ATP	N3-C4-N9	4.24	133.10	125.43
2	B	360	ATP	C8-N9-C4	4.09	110.02	106.90
2	A	401	ATP	N3-C4-N9	3.88	132.44	125.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/359 (95%)	0.13	16 (4%) 30 30	3, 21, 39, 47	3 (0%)
1	B	350/359 (97%)	0.03	8 (2%) 57 59	4, 16, 34, 53	0
All	All	693/718 (96%)	0.08	24 (3%) 43 43	3, 18, 37, 53	3 (0%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	351	THR	8.9
1	A	15	GLN	4.3
1	B	350	ASN	4.1
1	A	273	ARG	3.7
1	A	267	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CA	A	360	1/1	0.11	1.51	9,9,9,9	0
3	CA	B	361	1/1	0.08	0.01	5,5,5,5	0
2	ATP	A	401	31/31	0.09	-0.36	5,14,21,22	0
2	ATP	B	360	31/31	0.08	-0.37	2,9,13,16	0
3	CA	A	361	1/1	0.08	-1.79	46,46,46,46	0

6.5 Other polymers ⓘ

There are no such residues in this entry.