



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 02:53 AM GMT

PDB ID : 1FYZ
Title : METHANE MONOOXYGENASE HYDROXYLASE, FORM II REDUCED
BY SOAKING
Authors : Whittington, D.A.; Lippard, S.J.
Deposited on : 2000-10-03
Resolution : 2.15 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1094 (2.18-2.14)
Clashscore	79885	1299 (2.18-2.14)
Ramachandran outliers	78287	1272 (2.18-2.14)
Sidechain outliers	78261	1272 (2.18-2.14)
RSRZ outliers	66119	1094 (2.18-2.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	527	
1	B	527	
2	C	389	
2	D	389	
3	E	170	
3	F	170	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 18742 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METHANE MONOOXYGENASE COMPONENT A, ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	511	Total	C	N	O	S	0	0	0
			4185	2677	721	769	18			
1	B	511	Total	C	N	O	S	0	0	0
			4185	2677	721	769	18			

- Molecule 2 is a protein called METHANE MONOOXYGENASE COMPONENT A, BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	388	Total	C	N	O	S	0	0	0
			3193	2054	551	580	8			
2	D	387	Total	C	N	O	S	0	0	0
			3183	2048	549	578	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	370	ARG	ALA	CONFLICT	UNP P18798
D	370	ARG	ALA	CONFLICT	UNP P18798

- Molecule 3 is a protein called METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	167	Total	C	N	O	S	0	0	0
			1375	872	247	251	5			
3	F	168	Total	C	N	O	S	0	0	0
			1386	878	250	253	5			

- Molecule 4 is FE (II) ION (three-letter code: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Fe 2	0	0
4	A	2	Total 2	Fe 2	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Ca 1	0	0
5	C	2	Total 2	Ca 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	281	Total 281	O 281	0	0
6	B	263	Total 263	O 263	0	0
6	C	298	Total 298	O 298	0	0
6	D	171	Total 171	O 171	0	0
6	E	166	Total 166	O 166	0	0
6	F	49	Total 49	O 49	0	0




• Molecule 2: METHANE MONOOXYGENASE COMPONENT A, BETA CHAIN

Chain D:




• Molecule 3: METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN

Chain E:



• Molecule 3: METHANE MONOOXYGENASE COMPONENT A, GAMMA CHAIN

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.27Å 171.66Å 221.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.79 – 2.15 29.79 – 2.15	Depositor EDS
% Data completeness (in resolution range)	93.5 (29.79-2.15) 93.6 (29.79-2.15)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.04 (at 2.16Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.189 , 0.230 0.195 , 0.235	Depositor DCC
R_{free} test set	4903 reflections (3.54%)	DCC
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 145230 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18742	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/4310	0.55	0/5853
1	B	0.32	0/4310	0.55	0/5853
2	C	0.36	0/3289	0.56	0/4464
2	D	0.32	0/3279	0.52	0/4453
3	E	0.33	0/1404	0.58	0/1892
3	F	0.29	0/1416	0.49	0/1907
All	All	0.33	0/18008	0.55	0/24422

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4185	0	3981	104	0
1	B	4185	0	3981	124	0
2	C	3193	0	3042	63	0
2	D	3183	0	3029	88	0
3	E	1375	0	1370	17	0
3	F	1386	0	1377	50	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	1	0	0	0	0
5	C	2	0	0	0	0
6	A	281	0	0	5	0
6	B	263	0	0	6	0
6	C	298	0	0	9	0
6	D	171	0	0	2	0
6	E	166	0	0	0	0
6	F	49	0	0	1	0
All	All	18742	0	16780	391	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

The worst 5 of 391 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:F:41:THR:HG23	3:F:43:PHE:H	1.27	0.99
1:A:44:THR:HG22	1:A:46:TYR:H	1.32	0.95
1:B:78:GLN:HE22	1:B:150:GLN:HE21	1.02	0.94
1:A:352:ALA:HA	1:A:404:PRO:HB2	1.50	0.93
1:A:78:GLN:HE22	1:A:150:GLN:HE21	1.01	0.91

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	509/527 (97%)	487 (96%)	22 (4%)	0	100	100
1	B	509/527 (97%)	483 (95%)	24 (5%)	2 (0%)	43	39
2	C	386/389 (99%)	377 (98%)	8 (2%)	1 (0%)	50	47
2	D	385/389 (99%)	368 (96%)	14 (4%)	3 (1%)	27	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	165/170 (97%)	163 (99%)	2 (1%)	0	100	100
3	F	166/170 (98%)	161 (97%)	5 (3%)	0	100	100
All	All	2120/2172 (98%)	2039 (96%)	75 (4%)	6 (0%)	50	47

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	40	LYS
2	D	64	ALA
2	D	205	PRO
2	D	251	VAL
1	B	284	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/442 (98%)	421 (97%)	12 (3%)	56	58
1	B	433/442 (98%)	426 (98%)	7 (2%)	75	81
2	C	322/323 (100%)	314 (98%)	8 (2%)	60	64
2	D	321/323 (99%)	314 (98%)	7 (2%)	64	70
3	E	145/147 (99%)	141 (97%)	4 (3%)	56	58
3	F	146/147 (99%)	142 (97%)	4 (3%)	57	60
All	All	1800/1824 (99%)	1758 (98%)	42 (2%)	63	68

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	310	TYR
2	C	168	ARG
3	F	11	THR
2	C	33	ASN
2	C	80	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	214	ASN
1	B	343	HIS
3	E	165	HIS
1	B	227	ASN
1	B	268	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	511/527 (96%)	-0.19	5 (0%) 79 83	18, 28, 49, 67	0
1	B	511/527 (96%)	-0.24	9 (1%) 65 71	19, 29, 50, 70	0
2	C	388/389 (99%)	-0.44	6 (1%) 70 76	15, 22, 39, 57	0
2	D	387/389 (99%)	0.22	17 (4%) 33 36	21, 38, 61, 70	0
3	E	167/170 (98%)	-0.40	2 (1%) 75 80	17, 25, 40, 68	0
3	F	168/170 (98%)	0.57	14 (8%) 11 13	31, 48, 66, 74	0
All	All	2132/2172 (98%)	-0.13	53 (2%) 54 59	15, 30, 56, 74	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	17	ASN	4.7
3	E	4	LEU	4.6
2	D	44	LYS	3.7
2	D	380	ILE	3.4
3	F	83	PHE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CA	C	5007	1/1	0.07	-0.72	38,38,38,38	0
5	CA	C	5006	1/1	0.05	-1.04	55,55,55,55	0
4	FE2	A	5001	1/1	0.06	-1.47	30,30,30,30	0
4	FE2	B	5004	1/1	0.05	-2.41	43,43,43,43	0
4	FE2	B	5003	1/1	0.05	-2.60	29,29,29,29	0
4	FE2	A	5002	1/1	0.03	-3.32	39,39,39,39	0
5	CA	A	5005	1/1	0.03	-3.68	34,34,34,34	0

6.5 Other polymers

There are no such residues in this entry.