



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 01:01 PM GMT

PDB ID : 3G07
Title : Methyltransferase domain of human Bicoid-interacting protein 3 homolog (Drosophila)
Authors : Wu, H.; Dombrowski, L.; Tempel, W.; McCarthy, A.; Loppnau, P.; Bountra, C.; Weigelt, J.; Arrowsmith, C.H.; Edwards, A.M.; Bochkarev, A.; Min, J.; Plotnikov, A.N.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2009-01-27
Resolution : 2.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2232 (2.70-2.62)
Clashscore	79885	2700 (2.70-2.62)
Ramachandran outliers	78287	2657 (2.70-2.62)
Sidechain outliers	78261	2657 (2.70-2.62)
RSRZ outliers	66119	2234 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	
1	C	292	
1	D	292	
1	E	292	
1	F	292	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8567 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 7SK snRNA methylphosphate capping enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	Se	0	0	0
			1415	916	239	255	3	2			
1	B	193	Total	C	N	O	S	Se	0	0	0
			1422	919	241	257	3	2			
1	C	190	Total	C	N	O	S	Se	0	0	0
			1386	897	235	249	3	2			
1	D	193	Total	C	N	O	S	Se	0	0	0
			1426	919	242	260	3	2			
1	E	193	Total	C	N	O	S	Se	0	0	0
			1410	911	238	256	3	2			
1	F	193	Total	C	N	O	S	Se	0	0	0
			1346	859	235	247	3	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
A	399	SER	-	EXPRESSION TAG	UNP Q7L2J0
B	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
B	399	SER	-	EXPRESSION TAG	UNP Q7L2J0
C	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
C	399	SER	-	EXPRESSION TAG	UNP Q7L2J0
D	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
D	399	SER	-	EXPRESSION TAG	UNP Q7L2J0
E	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
E	399	SER	-	EXPRESSION TAG	UNP Q7L2J0
F	398	GLY	-	EXPRESSION TAG	UNP Q7L2J0
F	399	SER	-	EXPRESSION TAG	UNP Q7L2J0


- Molecule 2 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	C	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	D	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	E	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	F	1	Total	C	N	O	S	0	0
			27	15	6	5	1		


● Molecule 1: 7SK snRNA methylphosphate capping enzyme

Chain D: 




● Molecule 1: 7SK snRNA methylphosphate capping enzyme

Chain E: 



● Molecule 1: 7SK snRNA methylphosphate capping enzyme

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	182.99Å 182.99Å 151.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.70 – 2.65 29.70 – 2.62	Depositor EDS
% Data completeness (in resolution range)	50.0 (29.70-2.65) 99.9 (29.70-2.62)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.61Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.265 , 0.282 0.260 , 0.273	Depositor DCC
R_{free} test set	1532 reflections (2.70%)	DCC
Wilson B-factor (Å ²)	66.1	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 69.3	EDS
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56777 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8567	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1449	0.46	0/1984
1	B	0.31	0/1456	0.47	0/1992
1	C	0.31	0/1419	0.46	0/1942
1	D	0.33	0/1460	0.47	0/1996
1	E	0.31	0/1445	0.47	0/1979
1	F	0.31	0/1374	0.46	0/1883
All	All	0.31	0/8603	0.47	0/11776

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1415	0	0	2	0
1	B	1422	0	0	1	0
1	C	1386	0	0	2	0
1	D	1426	0	0	2	0
1	E	1410	0	0	2	0
1	F	1346	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	27	0	0	0	0
2	B	27	0	0	0	0
2	C	27	0	0	0	0
2	D	27	0	0	0	0
2	E	27	0	0	1	0
2	F	27	0	0	1	0
All	All	8567	0	0	11	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (11) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:F:446:ASP:OD2	1:F:608:ARG:NH1	2.45	0.49
1:E:446:ASP:OD2	1:E:608:ARG:NH1	2.48	0.47
1:F:581:LEU:O	2:F:701:SAM:N	2.50	0.45
1:A:446:ASP:OD2	1:A:608:ARG:NH1	2.50	0.45
1:C:446:ASP:OD2	1:C:608:ARG:NH1	2.50	0.44
1:B:446:ASP:OD2	1:B:608:ARG:NH1	2.51	0.44
1:D:446:ASP:OD2	1:D:608:ARG:NH1	2.51	0.43
1:A:574:GLU:OE1	1:D:546:THR:OG1	2.38	0.41
1:E:451:GLY:O	2:E:701:SAM:N	2.54	0.41
1:F:588:HIS:CE1	1:F:642:LEU:O	2.74	0.40
1:C:588:HIS:CE1	1:C:642:LEU:O	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/292 (64%)	177 (95%)	10 (5%)	0	100	100
1	B	187/292 (64%)	177 (95%)	10 (5%)	0	100	100
1	C	184/292 (63%)	175 (95%)	9 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	187/292 (64%)	177 (95%)	10 (5%)	0	100	100
1	E	187/292 (64%)	178 (95%)	9 (5%)	0	100	100
1	F	187/292 (64%)	177 (95%)	10 (5%)	0	100	100
All	All	1119/1752 (64%)	1061 (95%)	58 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/253 (52%)	129 (98%)	3 (2%)	63	88
1	B	134/253 (53%)	131 (98%)	3 (2%)	64	89
1	C	127/253 (50%)	125 (98%)	2 (2%)	75	93
1	D	135/253 (53%)	133 (98%)	2 (2%)	76	94
1	E	128/253 (51%)	125 (98%)	3 (2%)	63	88
1	F	114/253 (45%)	112 (98%)	2 (2%)	71	92
All	All	770/1518 (51%)	755 (98%)	15 (2%)	69	91

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	482	SER
1	A	581	LEU
1	A	621	SER
1	B	482	SER
1	B	581	LEU
1	B	621	SER
1	C	482	SER
1	C	581	LEU
1	D	581	LEU
1	D	621	SER
1	E	482	SER
1	E	581	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	621	SER
1	F	581	LEU
1	F	621	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SAM	A	701	-	26,29,29	0.92	2 (7%)	38,42,42	2.00	5 (13%)
2	SAM	B	701	-	26,29,29	0.93	2 (7%)	38,42,42	2.14	6 (15%)
2	SAM	C	701	-	26,29,29	0.92	2 (7%)	38,42,42	2.09	6 (15%)
2	SAM	D	701	-	26,29,29	0.95	2 (7%)	38,42,42	2.14	7 (18%)
2	SAM	E	701	-	26,29,29	0.96	2 (7%)	38,42,42	2.00	4 (10%)
2	SAM	F	701	-	26,29,29	0.92	2 (7%)	38,42,42	2.01	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAM	A	701	-	-	0/13/33/33	0/1/3/3
2	SAM	B	701	-	-	0/13/33/33	0/1/3/3
2	SAM	C	701	-	-	0/13/33/33	0/1/3/3
2	SAM	D	701	-	-	0/13/33/33	0/1/3/3
2	SAM	E	701	-	-	0/13/33/33	0/1/3/3
2	SAM	F	701	-	-	0/13/33/33	0/1/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	701	SAM	C2-N3	3.26	1.38	1.32
2	C	701	SAM	C2-N3	3.07	1.38	1.32
2	A	701	SAM	C2-N3	3.04	1.38	1.32
2	D	701	SAM	C2-N3	3.03	1.38	1.32
2	F	701	SAM	C2-N3	2.96	1.38	1.32
2	B	701	SAM	C2-N3	2.52	1.37	1.32
2	A	701	SAM	C2-N1	2.34	1.38	1.33
2	C	701	SAM	C2-N1	2.29	1.38	1.33
2	F	701	SAM	C2-N1	2.28	1.38	1.33
2	E	701	SAM	C2-N1	2.17	1.38	1.33
2	B	701	SAM	C2-N1	2.10	1.38	1.33
2	D	701	SAM	C2-N1	2.08	1.38	1.33

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	SAM	N3-C2-N1	-10.70	119.76	128.71
2	C	701	SAM	N3-C2-N1	-10.38	120.03	128.71
2	D	701	SAM	N3-C2-N1	-10.21	120.17	128.71
2	F	701	SAM	N3-C2-N1	-9.86	120.46	128.71
2	E	701	SAM	N3-C2-N1	-9.78	120.53	128.71
2	A	701	SAM	N3-C2-N1	-9.45	120.81	128.71
2	A	701	SAM	N3-C4-N9	4.12	132.87	125.43
2	B	701	SAM	N3-C4-N9	3.91	132.49	125.43
2	D	701	SAM	N3-C4-N9	3.83	132.34	125.43
2	E	701	SAM	N3-C4-N9	3.62	131.96	125.43
2	F	701	SAM	N3-C4-N9	3.24	131.28	125.43
2	C	701	SAM	N3-C4-N9	3.15	131.12	125.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	701	SAM	CB-CA-C	-3.11	106.74	111.44
2	F	701	SAM	C4-C5-N7	-2.51	107.37	109.52
2	A	701	SAM	C5-C4-N3	-2.47	120.33	125.70
2	A	701	SAM	CG-CB-CA	2.43	115.80	112.22
2	D	701	SAM	C5-C4-N3	-2.35	120.59	125.70
2	E	701	SAM	N7-C8-N9	-2.27	107.94	114.36
2	F	701	SAM	N7-C8-N9	-2.24	108.03	114.36
2	C	701	SAM	C4-C5-N7	-2.21	107.63	109.52
2	D	701	SAM	C2-N3-C4	2.21	120.30	114.01
2	B	701	SAM	C8-N9-C4	2.19	108.57	106.90
2	B	701	SAM	N7-C8-N9	-2.19	108.17	114.36
2	B	701	SAM	C2-N3-C4	2.19	120.25	114.01
2	B	701	SAM	C5-C4-N3	-2.18	120.96	125.70
2	D	701	SAM	N7-C8-N9	-2.16	108.25	114.36
2	E	701	SAM	C5-C4-N3	-2.15	121.02	125.70
2	F	701	SAM	C5-C4-N3	-2.13	121.06	125.70
2	C	701	SAM	C2-N3-C4	2.07	119.90	114.01
2	F	701	SAM	C2-N3-C4	2.07	119.90	114.01
2	C	701	SAM	N7-C8-N9	-2.06	108.54	114.36
2	A	701	SAM	C2-N3-C4	2.05	119.85	114.01
2	D	701	SAM	C4-C5-N7	-2.04	107.77	109.52
2	C	701	SAM	C5-C4-N3	-2.01	121.32	125.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	193/292 (66%)	0.22	9 (4%)	30 32	54, 85, 121, 136	0
1	B	193/292 (66%)	0.13	5 (2%)	53 56	48, 80, 118, 133	0
1	C	190/292 (65%)	0.20	13 (6%)	17 18	55, 87, 125, 138	0
1	D	193/292 (66%)	-0.04	4 (2%)	60 64	52, 84, 121, 134	0
1	E	193/292 (66%)	-0.06	7 (3%)	41 44	56, 83, 121, 137	0
1	F	193/292 (66%)	0.03	8 (4%)	35 38	58, 92, 128, 139	0
All	All	1155/1752 (65%)	0.08	46 (3%)	36 39	48, 85, 123, 139	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	628	THR	4.2
1	C	642	LEU	4.2
1	B	580	CYS	4.1
1	C	580	CYS	3.8
1	A	684	ALA	3.6
1	F	629	LEU	3.5
1	A	432	GLY	3.5
1	C	486	ASN	3.4
1	A	580	CYS	3.2
1	A	615	LEU	3.2
1	C	619	PRO	3.1
1	E	539	VAL	3.0
1	D	432	GLY	3.0
1	F	650	TYR	3.0
1	E	579	LEU	2.9
1	A	579	LEU	2.9
1	C	678	VAL	2.8
1	D	684	ALA	2.8
1	F	654	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	623	TYR	2.6
1	F	658	PHE	2.6
1	E	578	VAL	2.6
1	D	479	LEU	2.6
1	B	450	LEU	2.5
1	B	579	LEU	2.5
1	E	642	LEU	2.5
1	D	540	PRO	2.5
1	F	684	ALA	2.4
1	A	659	SER	2.4
1	F	642	LEU	2.3
1	C	567	LEU	2.3
1	C	664	VAL	2.3
1	A	563	ASP	2.2
1	B	448	LEU	2.2
1	C	663	LEU	2.2
1	C	579	LEU	2.2
1	A	471	VAL	2.1
1	C	448	LEU	2.1
1	A	570	ALA	2.1
1	B	578	VAL	2.1
1	C	614	VAL	2.1
1	F	580	CYS	2.1
1	C	636	ASN	2.1
1	F	448	LEU	2.0
1	E	640	ILE	2.0
1	E	540	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SAM	E	701	27/27	0.18	-0.24	61,70,89,159	0
2	SAM	B	701	27/27	0.16	-0.53	48,62,93,154	0
2	SAM	D	701	27/27	0.15	-0.55	55,69,88,158	0
2	SAM	F	701	27/27	0.13	-0.77	67,84,116,128	0
2	SAM	A	701	27/27	0.15	-0.89	58,70,95,168	0
2	SAM	C	701	27/27	0.12	-1.27	60,72,118,155	0

6.5 Other polymers

There are no such residues in this entry.