



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 08:33 PM GMT

PDB ID : 1G2D  
Title : STRUCTURE OF A CYS2HIS2 ZINC FINGER/TATA BOX COMPLEX  
(CLONE #2)  
Authors : Wolfe, S.A.; Grant, R.A.; Elrod-Erickson, M.; Pabo, C.O.  
Deposited on : 2000-10-18  
Resolution : 2.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	16	
1	D	16	
2	B	16	
2	E	16	
3	C	90	
3	F	90	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2997 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*GP\*AP\*CP\*GP\*CP\*TP\*AP\*TP\*AP\*AP\*AP\*AP\*GP\*GP\*AP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	16	Total	C	N	O	P	0	0	0
			332	158	70	89	15			
1	D	16	Total	C	N	O	P	0	0	0
			332	158	70	89	15			

- Molecule 2 is a DNA chain called 5'-D(\*TP\*CP\*CP\*TP\*TP\*TP\*TP\*AP\*TP\*AP\*GP\*CP\*GP\*TP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	P	0	0	0
			318	155	49	99	15			
2	E	16	Total	C	N	O	P	0	0	0
			318	155	49	99	15			

- Molecule 3 is a protein called TATA BOX ZINC FINGER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	89	Total	C	N	O	S	0	0	0
			738	452	153	125	8			
3	F	88	Total	C	N	O	S	0	0	0
			729	446	151	124	8			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	3	Total	Zn	0	0
			3	3		
4	F	3	Total	Zn	0	0
			3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total 28	O 28	0	0
5	B	34	Total 34	O 34	0	0
5	C	50	Total 50	O 50	0	0
5	D	26	Total 26	O 26	0	0
5	E	35	Total 35	O 35	0	0
5	F	51	Total 51	O 51	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-D(\*GP\*AP\*CP\*GP\*CP\*TP\*AP\*TP\*AP\*AP\*AP\*AP\*GP\*GP\*AP\*G)-3',

Chain A: 




- Molecule 1: 5'-D(\*GP\*AP\*CP\*GP\*CP\*TP\*AP\*TP\*AP\*AP\*AP\*AP\*GP\*GP\*AP\*G)-3',

Chain D: 



- Molecule 2: 5'-D(\*TP\*CP\*CP\*TP\*TP\*TP\*TP\*AP\*TP\*AP\*GP\*CP\*GP\*TP\*CP\*C)-3',

Chain B: 




- Molecule 2: 5'-D(\*TP\*CP\*CP\*TP\*TP\*TP\*TP\*AP\*TP\*AP\*GP\*CP\*GP\*TP\*CP\*C)-3',

Chain E: 



- Molecule 3: TATA BOX ZINC FINGER PROTEIN

Chain C: 



- Molecule 3: TATA BOX ZINC FINGER PROTEIN

Chain F: 

M201	E202	R203	P204	Y205		S211	C212	D213		K219	T220	N221	L222	D223	T224	H225	I226	R227	T228		P234	F235		C240	H241	R242		H247		C265		H281	T282	K283	I284		H287	Q288	LYS	ASP
------	------	------	------	------	--	------	------	------	--	------	------	------	------	------	------	------	------	------	------	--	------	------	--	------	------	------	--	------	--	------	--	------	------	------	------	--	------	------	-----	-----

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.10Å 105.10Å 104.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.20	Depositor
% Data completeness (in resolution range)	94.6 (20.00-2.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	6.60	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.225 , 0.262	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2997	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/375	0.78	0/578
1	D	0.41	0/375	0.71	0/578
2	B	0.39	0/353	0.75	0/542
2	E	0.42	0/353	0.76	0/542
3	C	0.34	0/755	0.59	0/1011
3	F	0.36	0/746	0.62	0/1000
All	All	0.38	0/2957	0.69	0/4251

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	332	0	180	11	0
1	D	332	0	180	6	0
2	B	318	0	185	15	0
2	E	318	0	185	8	0
3	C	738	0	724	16	0
3	F	729	0	711	15	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	3	0	0	0	0
4	F	3	0	0	0	0
5	A	28	0	0	0	0
5	B	34	0	0	1	0
5	C	50	0	0	1	0
5	D	26	0	0	0	0
5	E	35	0	0	3	0
5	F	51	0	0	1	0
All	All	2997	0	2165	66	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:2:DG:H2''	1:A:3:DA:H5'	1.17	1.11
1:A:2:DG:H2''	1:A:3:DA:C5'	1.86	1.04
2:B:27:DA:H2''	2:B:28:DG:H5''	1.43	0.98
1:D:57:DT:H2''	1:D:58:DA:H5'	1.53	0.90
1:A:5:DG:H2''	1:A:6:DC:H5'	1.50	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	87/90 (97%)	85 (98%)	2 (2%)	0	100	100
3	F	86/90 (96%)	84 (98%)	2 (2%)	0	100	100
All	All	173/180 (96%)	169 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	82/83 (99%)	79 (96%)	3 (4%)	45	54
3	F	81/83 (98%)	74 (91%)	7 (9%)	15	14
All	All	163/166 (98%)	153 (94%)	10 (6%)	26	28

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	203	ARG
3	F	211	SER
3	F	241	MET
3	F	201	MET
3	F	213	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	F	247	HIS
3	F	251	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.