



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 03:17 PM GMT

PDB ID : 1G3X
Title : INTERCALATION OF AN 9ACRIDINE-PEPTIDE DRUG IN A DNA DO-
DECAMER
Authors : Malinina, L.; Soler-Lopez, M.; Aymami, J.; Subirana, J.A.
Deposited on : 2000-10-25
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	12	
1	B	12	
1	C	12	
1	D	12	
1	E	12	
1	F	12	
1	G	12	
1	H	12	
1	I	12	
1	J	12	
1	K	12	
1	L	12	
2	M	5	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3055 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	B	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	C	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	D	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	E	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	F	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	G	11	Total	C	N	O	P	0	0	0
			224	107	43	64	10			
1	H	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	I	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	J	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	K	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	L	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			

- Molecule 2 is a protein called N(ALPHA)-(9-ACRIDINOYL)-TETRAARGININE-AMIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	M	5	Total	C	N	O	0	0	0
			48	35	8	5			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total Mg 1 1	0	0
3	J	1	Total Mg 1 1	0	0
3	K	2	Total Mg 2 2	0	0
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	L	1	Total Mg 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	11	Total O 11 11	0	0
4	B	16	Total O 16 16	1	0
4	C	3	Total O 3 3	0	0
4	D	5	Total O 5 5	0	0
4	E	8	Total O 8 8	0	0
4	F	3	Total O 3 3	0	0
4	G	9	Total O 9 9	0	0
4	H	2	Total O 2 2	0	0
4	I	7	Total O 7 7	0	0
4	J	10	Total O 10 10	0	0
4	K	11	Total O 11 11	1	0
4	L	17	Total O 17 17	1	0
4	M	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain A: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain B: 




- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain C: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain D: 




- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain E: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain F: 




- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain G: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain H: 




- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain I: 




- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain J: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain K: 



- Molecule 1: 5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3'

Chain L: 



- Molecule 2: N(ALPHA)-(9-ACRIDINOYL)-TETRAARGININE-AMIDE

Chain M: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.47Å 68.36Å 77.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70	Depositor
% Data completeness (in resolution range)	89.0 (20.00-2.70)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.220 , 0.266	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3055	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 9AC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.23	0/272	1.89	10/418 (2.4%)
1	B	1.23	0/272	2.29	22/418 (5.3%)
1	C	1.31	0/272	2.09	13/418 (3.1%)
1	D	1.25	0/272	2.01	13/418 (3.1%)
1	E	1.29	0/272	2.10	15/418 (3.6%)
1	F	1.14	0/272	2.05	13/418 (3.1%)
1	G	1.35	2/251 (0.8%)	2.20	14/386 (3.6%)
1	H	1.27	0/272	1.98	12/418 (2.9%)
1	I	1.19	0/272	2.15	21/418 (5.0%)
1	J	1.13	0/272	2.14	13/418 (3.1%)
1	K	1.27	0/272	2.59	25/418 (6.0%)
1	L	1.14	0/272	2.09	11/418 (2.6%)
2	M	0.76	0/31	0.97	0/38
All	All	1.23	2/3274 (0.1%)	2.13	182/5022 (3.6%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	403	DC	N1-C6	-5.85	1.33	1.37
1	G	410	DG	C3'-O3'	-5.14	1.37	1.44

The worst 5 of 182 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	DC	O4'-C1'-N1	13.03	117.12	108.00
1	L	617	DA	O4'-C1'-N9	-12.15	99.50	108.00
1	K	603	DC	O4'-C1'-N1	-11.43	100.00	108.00
1	K	606	DA	O4'-C1'-C2'	-10.96	97.13	105.90
1	F	324	DG	O4'-C4'-C3'	-10.27	99.84	106.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	243	0	136	2	0
1	B	243	0	136	2	0
1	C	243	0	136	2	0
1	D	243	0	136	0	0
1	E	243	0	136	1	0
1	F	243	0	136	0	0
1	G	224	0	125	1	0
1	H	243	0	136	1	0
1	I	243	0	136	3	0
1	J	243	0	136	0	0
1	K	243	0	136	1	0
1	L	243	0	136	0	0
2	M	48	0	38	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	G	1	0	0	0	0
3	J	1	0	0	0	0
3	K	2	0	0	0	0
3	L	1	0	0	0	0
4	A	11	0	0	0	0
4	B	16	0	0	0	0
4	C	3	0	0	0	0
4	D	5	0	0	0	0
4	E	8	0	0	0	0
4	F	3	0	0	0	0
4	G	9	0	0	0	0
4	H	2	0	0	0	0
4	I	7	0	0	0	0
4	J	10	0	0	0	0
4	K	11	0	0	0	0
4	L	17	0	0	0	0
4	M	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3055	0	1659	14	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:G:403:DC:H2'	1:G:404:DG:C8	2.27	0.69
1:H:415:DC:H2'	1:H:416:DG:C8	2.35	0.61
1:I:503:DC:H2'	1:I:504:DG:C8	2.43	0.54
1:B:115:DC:H2'	1:B:116:DG:C8	2.43	0.53
1:K:611:DC:H2''	1:K:612:DG:H5''	1.96	0.47

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	M	2/5 (40%)	0	1 (50%)	1 (50%)	0 0

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	M	702	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	M	1/4 (25%)	0	1 (100%)	0 0

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	703	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.