



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 04:10 AM GMT

PDB ID : 3G8C
Title : Crystal Structure of Biotin Carboxylase in Complex with Biotin, Bicarbonate, ADP and Mg Ion
Authors : Chou, C.Y.; Yu, L.P.; Tong, L.
Deposited on : 2009-02-12
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	444	
1	B	444	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	BTN	B	1005	-	X

2 Entry composition i


There are 6 unique types of molecules in this entry. The entry contains 7794 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biotin carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3421	2155	611	633	22			
1	B	444	Total	C	N	O	S	0	0	0
			3421	2155	611	633	22			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is BIOTIN (three-letter code: BTN) (formula: $C_{10}H_{16}N_2O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
4	B	1	Total	C	N	O	S	0	0
			16	10	2	3	1		

- Molecule 5 is BICARBONATE ION (three-letter code: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	1	3		
5	B	1	Total	C	O	0	0
			4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	396	Total	O	0	0
			396	396		
6	B	460	Total	O	0	0
			460	460		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Biotin carboxylase



- Molecule 1: Biotin carboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.32Å 106.17Å 121.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.04 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-2.00) 99.2 (29.04-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.182 , 0.216 0.182 , 0.215	Depositor DCC
R_{free} test set	3678 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 72986 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7794	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BCT, MG, BTN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3484	0.53	0/4702
1	B	0.43	0/3484	0.53	0/4702
All	All	0.43	0/6968	0.53	0/9404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3421	0	3445	25	0
1	B	3421	0	3445	26	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	15	0	0
4	B	16	0	15	0	0
5	A	4	0	0	0	0
5	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	396	0	0	3	4
6	B	460	0	0	9	4
All	All	7794	0	6944	51	4

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (51) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:106:LYS:HE2	1:B:270:ARG:HH12	1.34	0.91
1:A:172:VAL:HG11	1:A:178:LEU:HD13	1.55	0.89
1:B:106:LYS:HE2	1:B:270:ARG:NH1	1.86	0.88
1:A:1:MET:HE3	6:A:646:HOH:O	1.82	0.79
1:B:428:GLU:H	1:B:428:GLU:CD	1.90	0.75
1:A:370:HIS:CD2	1:A:370:HIS:H	2.13	0.64
1:B:106:LYS:CE	1:B:270:ARG:HH12	2.10	0.64
1:B:407:GLN:HG2	6:B:881:HOH:O	1.98	0.64
1:A:178:LEU:HD12	1:A:181:SER:HB2	1.81	0.62
1:B:67:ILE:CD1	1:B:94:GLN:HG2	2.29	0.62
1:B:370:HIS:CD2	1:B:370:HIS:H	2.20	0.59
1:A:436:ASN:C	1:A:436:ASN:HD22	2.05	0.59
1:B:278:LEU:HD22	1:B:287:ILE:HG21	1.85	0.58
1:A:155:PRO:HB3	1:A:173:ARG:HG2	1.85	0.57
1:A:311:GLU:OE2	1:A:314:ARG:NH1	2.38	0.56
1:A:172:VAL:HG11	1:A:178:LEU:CD1	2.32	0.55
1:B:370:HIS:HE1	6:B:652:HOH:O	1.90	0.55
1:A:206:ASN:N	1:A:207:PRO:HD3	2.24	0.53
1:B:252:LEU:HD12	6:B:764:HOH:O	2.09	0.52
1:B:298:PRO:O	1:B:302:MET:HG2	2.09	0.52
1:B:131:VAL:HG22	1:B:285:TYR:HB3	1.93	0.50
1:B:314:ARG:NH1	6:B:879:HOH:O	2.44	0.50
1:A:131:VAL:HG22	1:A:285:TYR:HB3	1.94	0.50
1:A:34:SER:HB2	1:A:37:ARG:NH2	2.28	0.49
1:B:323:ILE:HG12	1:B:327:GLU:HB2	1.94	0.48
1:B:314:ARG:HD3	6:B:496:HOH:O	2.14	0.47
1:B:108:GLU:HG3	6:B:696:HOH:O	2.13	0.47
1:B:105:PRO:HG2	1:B:291:THR:HB	1.95	0.47
1:A:167:ARG:HD2	1:A:192:ALA:HB2	1.97	0.47
1:A:158:ILE:HB	1:A:170:ARG:HB3	1.97	0.47
1:B:423:ARG:HD3	6:B:767:HOH:O	2.15	0.46
1:A:298:PRO:O	1:A:302:MET:HG2	2.15	0.46
1:B:146:ARG:NH1	6:B:683:HOH:O	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:370:HIS:HE1	6:A:473:HOH:O	1.99	0.45
1:B:159:LYS:HG2	1:B:169:MET:HG2	1.99	0.45
1:A:436:ASN:ND2	1:A:438:HIS:H	2.15	0.45
1:A:160:ALA:HA	1:A:198:VAL:HG12	1.98	0.44
1:A:278:LEU:HD22	1:A:287:ILE:HG21	1.99	0.44
1:B:157:ILE:HD11	1:B:169:MET:HE3	1.99	0.44
1:A:133:GLY:HA2	1:A:152:ILE:HD13	1.99	0.44
1:B:407:GLN:HG3	6:B:801:HOH:O	2.18	0.43
1:A:442:LYS:NZ	6:A:687:HOH:O	2.49	0.43
1:B:206:ASN:CG	1:B:206:ASN:O	2.57	0.42
1:B:369:SER:OG	1:B:371:ILE:HG12	2.20	0.42
1:B:251:GLU:H	1:B:251:GLU:CD	2.22	0.42
1:A:18:LEU:HD21	1:A:22:LYS:HE2	2.02	0.41
1:A:142:MET:SD	1:A:179:ALA:HA	2.61	0.41
1:A:157:ILE:HB	1:A:203:TYR:HD1	1.86	0.41
1:A:140:ASP:O	1:A:142:MET:HG2	2.21	0.41
1:B:170:ARG:CZ	1:B:184:MET:HG2	2.51	0.40
1:A:301:GLU:HG2	1:A:306:VAL:O	2.22	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
6:A:821:HOH:O	6:B:660:HOH:O[2_655]	2.02	0.18
6:A:861:HOH:O	6:B:707:HOH:O[2_655]	2.03	0.17
6:A:831:HOH:O	6:B:790:HOH:O[2_655]	2.06	0.14
6:A:835:HOH:O	6:B:852:HOH:O[2_655]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/444 (100%)	425 (96%)	17 (4%)	0	100	100
1	B	442/444 (100%)	430 (97%)	12 (3%)	0	100	100
All	All	884/888 (100%)	855 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/357 (100%)	349 (98%)	8 (2%)	64	65
1	B	357/357 (100%)	351 (98%)	6 (2%)	73	75
All	All	714/714 (100%)	700 (98%)	14 (2%)	68	69

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	LYS
1	A	177	GLU
1	A	180	GLN
1	A	193	PHE
1	A	251	GLU
1	A	278	LEU
1	A	436	ASN
1	A	443	LYS
1	B	138	LEU
1	B	151	ARG
1	B	205	GLU
1	B	234	ARG
1	B	278	LEU
1	B	444	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	A	370	HIS
1	A	404	ASN
1	A	426	ASN
1	A	431	GLN
1	A	436	ASN
1	A	438	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	145	ASN
1	B	319	GLN
1	B	370	HIS
1	B	404	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	A	1000	3	29,29,29	1.08	2 (6%)	45,45,45	1.78	6 (13%)
4	BTN	A	1004	-	17,17,17	2.81	3 (17%)	23,23,23	1.36	3 (13%)
5	BCT	A	1006	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ADP	B	1001	3	29,29,29	1.10	2 (6%)	45,45,45	1.81	7 (15%)
4	BTN	B	1005	-	17,17,17	2.83	3 (17%)	23,23,23	1.54	5 (21%)
5	BCT	B	1007	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1000	3	-	0/16/32/32	0/1/3/3
4	BTN	A	1004	-	-	0/7/28/28	0/0/2/2
5	BCT	A	1006	-	-	0/0/0/0	0/0/0/0
2	ADP	B	1001	3	-	0/16/32/32	0/1/3/3
4	BTN	B	1005	-	-	0/7/28/28	0/0/2/2
5	BCT	B	1007	-	-	0/0/0/0	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1005	BTN	C2-S1	-8.02	1.68	1.82
4	A	1004	BTN	C2-S1	-7.89	1.68	1.82
4	B	1005	BTN	O3-C3	7.78	1.41	1.23
4	A	1004	BTN	O3-C3	7.76	1.41	1.23
2	A	1000	ADP	C5-C4	3.39	1.48	1.40
2	B	1001	ADP	C5-C4	3.33	1.48	1.40
4	A	1004	BTN	C5-C4	-2.31	1.52	1.56
4	B	1005	BTN	C5-C4	-2.27	1.52	1.56
2	B	1001	ADP	C4-N9	-2.23	1.34	1.37
2	A	1000	ADP	C4-N9	-2.14	1.34	1.37

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	ADP	N3-C2-N1	-6.61	123.18	128.71
2	A	1000	ADP	N3-C2-N1	-6.25	123.48	128.71
2	A	1000	ADP	N3-C4-N9	5.52	135.40	125.43
2	B	1001	ADP	N3-C4-N9	5.17	134.77	125.43
2	B	1001	ADP	C4-C5-N7	-3.50	106.52	109.52
4	B	1005	BTN	C5-C6-S1	3.45	108.79	106.04
2	B	1001	ADP	C5-C4-N3	-3.39	118.32	125.70
2	A	1000	ADP	C5-C4-N3	-3.37	118.36	125.70
2	B	1001	ADP	O4'-C1'-N9	3.32	111.52	108.44
2	A	1000	ADP	C4-C5-N7	-3.28	106.71	109.52
2	A	1000	ADP	O4'-C1'-N9	3.26	111.47	108.44
4	B	1005	BTN	C4-C2-S1	3.15	109.23	105.29
4	A	1004	BTN	N2-C3-N1	2.96	110.77	108.99
4	A	1004	BTN	C4-C2-S1	2.88	108.90	105.29
4	B	1005	BTN	N2-C3-N1	2.62	110.56	108.99
2	B	1001	ADP	C2-N3-C4	2.58	121.35	114.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	ADP	C2-N3-C4	2.57	121.33	114.01
2	B	1001	ADP	C3'-C2'-C1'	2.26	104.45	100.91
4	B	1005	BTN	C4-N2-C3	-2.21	110.65	112.54
4	A	1004	BTN	O12-C11-C10	2.05	121.46	114.22
4	B	1005	BTN	O12-C11-C10	2.05	121.45	114.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/444 (100%)	0.72	69 (15%) 3 3	15, 26, 76, 80	0
1	B	444/444 (100%)	0.02	12 (2%) 52 52	17, 26, 50, 69	0
All	All	888/888 (100%)	0.37	81 (9%) 9 9	15, 26, 70, 80	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	166	GLY	10.7
1	A	179	ALA	10.2
1	A	189	ALA	9.8
1	A	193	PHE	9.8
1	A	139	GLY	9.3
1	A	187	ALA	8.5
1	A	188	GLU	8.3
1	A	174	GLY	8.2
1	A	191	ALA	8.2
1	A	192	ALA	8.0
1	A	171	VAL	7.9
1	A	185	THR	7.9
1	A	196	ASP	7.5
1	A	152	ILE	7.4
1	A	184	MET	7.2
1	A	146	ARG	7.2
1	A	178	LEU	7.2
1	A	167	ARG	7.0
1	A	170	ARG	7.0
1	A	138	LEU	6.7
1	A	154	TYR	6.5
1	A	195	ASN	6.3
1	A	176	ALA	6.2
1	A	148	ILE	6.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	181	SER	5.9
1	B	166	GLY	5.9
1	A	180	GLN	5.9
1	A	151	ARG	5.8
1	A	144	LYS	5.6
1	A	169	MET	5.6
1	A	183	SER	5.5
1	A	150	LYS	5.5
1	A	141	ASP	5.5
1	B	167	ARG	5.4
1	A	197	MET	5.2
1	A	137	PRO	5.2
1	A	143	ASP	5.2
1	A	190	LYS	5.2
1	A	194	SER	5.2
1	B	164	GLY	5.1
1	A	164	GLY	5.1
1	A	165	GLY	5.1
1	A	157	ILE	5.0
1	A	158	ILE	4.9
1	A	182	ILE	4.7
1	A	155	PRO	4.6
1	A	140	ASP	4.4
1	A	198	VAL	4.4
1	A	186	ARG	4.3
1	A	149	ALA	4.2
1	A	145	ASN	4.2
1	A	147	ALA	4.1
1	A	142	MET	4.0
1	A	200	MET	4.0
1	A	172	VAL	3.9
1	B	168	GLY	3.9
1	A	173	ARG	3.9
1	A	168	GLY	3.8
1	A	1	MET	3.3
1	A	177	GLU	3.3
1	A	175	ASP	3.3
1	B	163	GLY	3.2
1	A	133	GLY	3.0
1	B	235	ARG	3.0
1	A	135	ASP	2.9
1	B	194	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	254	ARG	2.8
1	B	193	PHE	2.7
1	A	203	TYR	2.6
1	A	134	SER	2.4
1	A	156	VAL	2.4
1	A	442	LYS	2.4
1	A	162	GLY	2.4
1	B	281	ASN	2.3
1	B	165	GLY	2.3
1	A	160	ALA	2.3
1	A	251	GLU	2.3
1	A	199	TYR	2.2
1	A	136	GLY	2.1
1	B	1	MET	2.0
1	B	176	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	BTN	B	1005	16/16	0.20	2.08	41,43,45,45	0
4	BTN	A	1004	16/16	0.15	0.89	32,35,39,40	0
2	ADP	B	1001	27/27	0.14	-0.43	39,44,50,51	0
2	ADP	A	1000	27/27	0.15	-0.47	48,50,51,52	0
3	MG	B	1003	1/1	0.13	-0.70	66,66,66,66	0
5	BCT	B	1007	4/4	0.10	-1.80	22,23,23,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	BCT	A	1006	4/4	0.07	-3.01	19,20,20,20	0
3	MG	A	1002	1/1	0.07	-5.88	46,46,46,46	0

6.5 Other polymers ⓘ

There are no such residues in this entry.