



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 04:48 PM GMT

PDB ID : 1GCJ
Title : N-TERMINAL FRAGMENT OF IMPORTIN-BETA
Authors : Lee, S.J.; Imamoto, N.; Sakai, H.; Nakagawa, A.; Kose, S.; Koike, M.; Yamamoto, M.; Kumasaka, T.; Yoneda, Y.; Tsukihara, T.
Deposited on : 2000-07-31
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7389 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMPORTIN BETA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	460	Total	C	N	O	S	Se	0	0	0
			3589	2260	597	708	11	13			
1	B	460	Total	C	N	O	S	Se	0	0	0
			3589	2260	597	708	11	13			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	146	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	181	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	219	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	245	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	252	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	256	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	268	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	291	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	353	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	388	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	410	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	417	MSE	MET	MODIFIED RESIDUE	UNP P70168
A	450	VAL	-	CLONING ARTIFACT	UNP P70168
A	451	PRO	-	CLONING ARTIFACT	UNP P70168
A	452	ASN	-	CLONING ARTIFACT	UNP P70168
A	453	SER	-	CLONING ARTIFACT	UNP P70168
A	454	SER	-	CLONING ARTIFACT	UNP P70168
A	500	PRO	-	CLONING ARTIFACT	UNP P70168
A	501	ASP	-	CLONING ARTIFACT	UNP P70168
A	502	TYR	-	CLONING ARTIFACT	UNP P70168
A	503	ALA	-	CLONING ARTIFACT	UNP P70168
A	504	SER	-	CLONING ARTIFACT	UNP P70168
A	505	GLY	-	CLONING ARTIFACT	UNP P70168
B	1	MSE	MET	MODIFIED RESIDUE	UNP P70168

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	146	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	181	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	219	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	245	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	252	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	256	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	268	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	291	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	353	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	388	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	410	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	417	MSE	MET	MODIFIED RESIDUE	UNP P70168
B	450	VAL	-	CLONING ARTIFACT	UNP P70168
B	451	PRO	-	CLONING ARTIFACT	UNP P70168
B	452	ASN	-	CLONING ARTIFACT	UNP P70168
B	453	SER	-	CLONING ARTIFACT	UNP P70168
B	454	SER	-	CLONING ARTIFACT	UNP P70168
B	500	PRO	-	CLONING ARTIFACT	UNP P70168
B	501	ASP	-	CLONING ARTIFACT	UNP P70168
B	502	TYR	-	CLONING ARTIFACT	UNP P70168
B	503	ALA	-	CLONING ARTIFACT	UNP P70168
B	504	SER	-	CLONING ARTIFACT	UNP P70168
B	505	GLY	-	CLONING ARTIFACT	UNP P70168

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	115	Total O 115 115	0	0
2	B	96	Total O 96 96	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: IMPORTIN BETA



• Molecule 1: IMPORTIN BETA



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.83Å 103.78Å 126.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.18 – 2.60	Depositor
% Data completeness (in resolution range)	97.9 (39.18-2.60)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.215 , 0.270	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7389	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/3643	0.56	0/4938
1	B	0.35	1/3643 (0.0%)	0.54	0/4938
All	All	0.34	1/7286 (0.0%)	0.55	0/9876

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	436	CYS	CB-SG	6.80	1.93	1.82

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3589	0	3563	140	0
1	B	3589	0	3563	141	0
2	A	115	0	0	15	0
2	B	96	0	0	7	0
All	All	7389	0	7126	275	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 19.

The worst 5 of 275 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:105:ARG:HB3	1:A:106:PRO:HD3	1.44	0.99
1:B:105:ARG:HB3	1:B:106:PRO:HD3	1.45	0.98
1:B:188:ASN:HD22	1:B:188:ASN:H	1.12	0.95
1:A:170:SER:HB2	1:A:208:ASN:HD21	1.30	0.94
1:A:177:ILE:HG22	1:A:181:MSE:HE2	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/460 (100%)	436 (95%)	16 (4%)	6 (1%)	18	35
1	B	458/460 (100%)	439 (96%)	15 (3%)	4 (1%)	25	49
All	All	916/920 (100%)	875 (96%)	31 (3%)	10 (1%)	21	42

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	337	ASP
1	B	447	VAL
1	A	339	ASP
1	A	361	ASP
1	A	212	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/386 (103%)	389 (98%)	10 (2%)	60	86
1	B	399/386 (103%)	386 (97%)	13 (3%)	50	79
All	All	798/772 (103%)	775 (97%)	23 (3%)	55	83

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	LEU
1	B	32	ASN
1	B	436	CYS
1	B	23	LYS
1	B	46	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	ASN
1	B	46	ASN
1	B	220	GLN
1	A	366	HIS
1	A	401	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.