



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 03:02 PM GMT

PDB ID : 3GCU  
Title : Human P38 MAP kinase in complex with RL48  
Authors : Gruetter, C.; Simard, J.R.; Getlik, M.; Rauh, D.  
Deposited on : 2009-02-22  
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	360	
1	B	360	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	BOG	A	403	-	X
3	BOG	B	403	-	X
4	MES	A	404	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5897 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	1	0
			2720	1745	463	500	12			
1	B	337	Total	C	N	O	S	0	2	0
			2715	1743	463	496	13			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	EXPRESSION TAG	UNP Q16539
B	1	GLY	-	EXPRESSION TAG	UNP Q16539

- Molecule 2 is 1-{3-[(6-AMINOQUINAZOLIN-4-YL)AMINO]PHENYL}-3-[3-TERT-BUTYL-1-(4-METHYLPHENYL)-1H-PYRAZOL-5-YL]UREA (three-letter code: R48) (formula: C<sub>29</sub>H<sub>30</sub>N<sub>8</sub>O).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			38	29	8	1		
2	B	1	Total	C	N	O	0	0
			38	29	8	1		

- Molecule 3 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			20	14	6		
3	A	1	Total	C	O	0	0
			20	14	6		
3	B	1	Total	C	O	0	0
			20	14	6		
3	B	1	Total	C	O	0	0
			20	14	6		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONICACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	161	Total	O	0	0
			161	161		
5	B	133	Total	O	0	0
			133	133		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitogen-activated protein kinase 14

Chain A:



- Molecule 1: Mitogen-activated protein kinase 14

Chain B:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.47Å 74.09Å 71.71Å 90.00° 93.76° 90.00°	Depositor
Resolution (Å)	32.41 – 2.10 32.41 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (32.41-2.10) 99.8 (32.41-2.10)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.26 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.189 , 0.235 0.189 , 0.234	Depositor DCC
$R_{free}$ test set	1077 reflections (2.56%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.7	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.0	EDS
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 43071 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5897	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: R48, MES, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/2786	0.72	2/3787 (0.1%)
1	B	0.67	0/2783	0.75	3/3779 (0.1%)
All	All	0.66	0/5569	0.74	5/7566 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	LEU	CA-CB-CG	6.19	129.53	115.30
1	B	268	MET	CG-SD-CE	5.57	109.11	100.20
1	B	48	LEU	CA-CB-CG	5.54	128.05	115.30
1	B	220	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	330	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2704	33	0
1	B	2715	0	2718	47	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	38	0	30	3	0
2	B	38	0	30	0	0
3	A	40	0	56	1	0
3	B	40	0	56	3	0
4	A	12	0	12	0	0
5	A	161	0	0	6	0
5	B	133	0	0	1	0
All	All	5897	0	5606	79	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (79) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:109[B]:MET:SD	1:B:169:PHE:CZ	2.78	0.77
1:A:148:HIS:HD2	1:A:150:ASP:H	1.30	0.77
1:B:342:TYR:CE2	1:B:346:ILE:HD11	2.22	0.75
1:B:202:GLN:NE2	5:B:587:HOH:O	2.21	0.73
1:B:203:THR:OG1	1:B:293:SER:HB2	1.92	0.69
1:B:109[B]:MET:HE2	1:B:157:ALA:HA	1.75	0.68
1:B:109[B]:MET:SD	1:B:169:PHE:HZ	2.18	0.66
1:B:109[B]:MET:CE	1:B:169:PHE:CE2	2.78	0.66
1:B:305:HIS:HD2	1:B:307:TYR:H	1.43	0.65
1:A:74:LEU:HD23	2:A:401:R48:H1N	1.80	0.64
1:B:283:ASP:OD2	1:B:305:HIS:HE1	1.80	0.64
1:B:109[B]:MET:CE	1:B:169:PHE:CZ	2.81	0.63
1:B:239:VAL:HG21	1:B:291:LEU:HD13	1.80	0.63
1:A:185:THR:HG21	1:A:189:ARG:NH1	2.13	0.63
1:B:18:TRP:HZ2	1:B:37:SER:HB2	1.62	0.63
2:A:401:R48:N1E	5:A:467:HOH:O	2.27	0.62
1:A:186:ARG:HB2	1:A:194:MET:HE1	1.82	0.61
1:B:148:HIS:HD2	1:B:150:ASP:H	1.45	0.61
1:A:241:THR:HG23	1:A:265:MET:H	1.66	0.60
1:A:202:GLN:NE2	5:A:461:HOH:O	2.33	0.60
1:B:305:HIS:CD2	1:B:307:TYR:H	2.20	0.59
1:A:7:THR:O	1:A:22:GLU:HG3	2.02	0.59
1:A:239:VAL:HG21	1:A:291:LEU:HD13	1.85	0.58
1:A:202:GLN:HG2	5:A:611:HOH:O	2.04	0.56
1:A:148:HIS:CD2	1:A:150:ASP:H	2.19	0.56
1:A:48:LEU:HD13	1:A:107:HIS:CE1	2.41	0.56
1:A:155:ASN:ND2	1:A:169:PHE:H	2.03	0.56
1:B:155:ASN:ND2	1:B:169:PHE:H	2.04	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:323:TYR:CE2	1:A:325:GLN:HG3	2.40	0.55
1:B:155:ASN:HA	1:B:169:PHE:HD2	1.73	0.54
1:B:109[B]:MET:HE3	1:B:169:PHE:CE2	2.43	0.54
1:B:109[B]:MET:HE1	1:B:169:PHE:CE2	2.44	0.52
1:A:186:ARG:HB2	1:A:194:MET:CE	2.40	0.51
1:B:167:LEU:HD13	1:B:169:PHE:CE2	2.45	0.51
1:A:269:ASN:HD22	1:A:269:ASN:C	2.13	0.51
1:A:148:HIS:HD2	1:A:150:ASP:N	2.05	0.50
1:A:128:GLN:NE2	5:A:554:HOH:O	2.43	0.50
1:A:148:HIS:HE1	1:A:167:LEU:O	1.95	0.50
1:B:310:GLN:HG3	1:B:311:TYR:CD1	2.47	0.49
1:A:203:THR:OG1	1:A:293:SER:HB2	2.12	0.49
1:B:247:LEU:HD13	1:B:250:ILE:HD12	1.95	0.49
1:B:18:TRP:CZ2	1:B:37:SER:HB2	2.45	0.48
1:B:195:LEU:HD22	3:B:403:BOG:H8'3	1.95	0.47
1:B:148:HIS:HE1	1:B:167:LEU:O	1.97	0.47
1:B:282:VAL:O	1:B:286:GLU:HG3	2.15	0.47
1:B:186:ARG:HB2	1:B:194:MET:HE1	1.97	0.46
1:B:49:ARG:HG3	1:B:49:ARG:HH11	1.79	0.46
1:B:109[B]:MET:HE3	1:B:157:ALA:HB2	1.96	0.46
1:A:35:TYR:HE1	5:A:519:HOH:O	1.99	0.46
1:A:295:LYS:HD2	1:A:295:LYS:HA	1.81	0.46
1:A:167:LEU:HA	2:A:401:R48:H1BA	1.98	0.46
1:B:229:ILE:HG21	3:B:403:BOG:H2	1.97	0.46
1:B:310:GLN:HG3	1:B:311:TYR:CE1	2.53	0.44
1:A:202:GLN:CG	5:A:611:HOH:O	2.62	0.44
1:B:16:THR:HG23	1:B:18:TRP:HE1	1.82	0.44
1:B:46:THR:OG1	1:B:48:LEU:HD12	2.18	0.44
1:A:249:LYS:O	3:A:403:BOG:H61	2.18	0.44
1:B:155:ASN:ND2	1:B:169:PHE:HB2	2.33	0.44
1:B:109[B]:MET:HE3	1:B:169:PHE:CZ	2.52	0.44
1:B:109[B]:MET:HG3	1:B:157:ALA:CB	2.48	0.43
1:A:117:VAL:HG22	1:A:216:LEU:CD2	2.49	0.43
1:B:195:LEU:HD12	3:B:402:BOG:H5'2	2.00	0.43
1:A:55:LEU:HD12	1:A:68:THR:HG23	2.00	0.43
1:A:109:MET:HB3	1:A:157:ALA:HB1	2.00	0.43
1:A:63:ILE:HG21	1:B:160:GLU:HB3	2.01	0.43
1:B:342:TYR:CZ	1:B:346:ILE:HD11	2.54	0.42
1:B:13:LEU:O	1:B:16:THR:HG22	2.20	0.42
1:B:87:LEU:HD11	1:B:107[A]:HIS:NE2	2.35	0.41
1:A:114:ASN:OD1	1:A:183:VAL:HG21	2.20	0.41
1:B:63:ILE:HD11	1:B:67:ARG:NH2	2.36	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:233:LYS:O	1:B:237:ARG:HD3	2.21	0.41
1:A:84:ILE:HB	1:A:167:LEU:HB3	2.02	0.41
1:B:84:ILE:HD11	1:B:106:THR:HB	2.02	0.41
1:B:321:ASP:C	1:B:321:ASP:OD2	2.59	0.41
1:B:109[B]:MET:HE2	1:B:157:ALA:CA	2.49	0.40
1:A:48:LEU:HD13	1:A:107:HIS:HE1	1.85	0.40
1:B:270:PHE:CD1	1:B:286:GLU:HG2	2.56	0.40
1:A:69:TYR:CE2	1:A:340:LEU:HB3	2.56	0.40
1:A:229:ILE:HG12	1:B:45:LYS:HA	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/360 (93%)	327 (97%)	9 (3%)	0	100	100
1	B	335/360 (93%)	324 (97%)	10 (3%)	1 (0%)	50	49
All	All	671/720 (93%)	651 (97%)	19 (3%)	1 (0%)	59	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	334	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/319 (93%)	281 (95%)	16 (5%)	31	27
1	B	298/319 (93%)	279 (94%)	19 (6%)	25	20
All	All	595/638 (93%)	560 (94%)	35 (6%)	29	23

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	SER
1	A	74	LEU
1	A	86	LEU
1	A	87	LEU
1	A	97	GLU
1	A	114	ASN
1	A	138	LEU
1	A	160	GLU
1	A	167	LEU
1	A	189	ARG
1	A	195	LEU
1	A	245	GLU
1	A	253	GLU
1	A	269	ASN
1	A	325	GLN
1	A	333	LEU
1	B	23	ARG
1	B	37	SER
1	B	48	LEU
1	B	63	ILE
1	B	72	LEU
1	B	86	LEU
1	B	87	LEU
1	B	94	ARG
1	B	102	VAL
1	B	109[A]	MET
1	B	109[B]	MET
1	B	119	CYS
1	B	122	LEU
1	B	130	LEU
1	B	160	GLU
1	B	198	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	246	LEU
1	B	291	LEU
1	B	332	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	128	GLN
1	A	148	HIS
1	A	155	ASN
1	A	257	ASN
1	A	264	GLN
1	A	269	ASN
1	B	114	ASN
1	B	120	GLN
1	B	128	GLN
1	B	148	HIS
1	B	155	ASN
1	B	305	HIS
1	B	310	GLN
1	B	325	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	R48	A	401	-	42,42,42	1.86	8 (19%)	61,61,61	2.89	20 (32%)
3	BOG	A	402	-	20,20,20	0.69	0	25,25,25	0.91	1 (4%)
3	BOG	A	403	-	20,20,20	0.94	2 (10%)	25,25,25	1.02	1 (4%)
4	MES	A	404	-	12,12,12	1.98	3 (25%)	16,16,16	2.45	5 (31%)
2	R48	B	401	-	42,42,42	1.73	8 (19%)	61,61,61	2.79	16 (26%)
3	BOG	B	402	-	20,20,20	0.84	1 (5%)	25,25,25	1.70	4 (16%)
3	BOG	B	403	-	20,20,20	0.92	1 (5%)	25,25,25	1.76	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R48	A	401	-	-	0/20/22/22	0/3/5/5
3	BOG	A	402	-	-	0/11/31/31	0/1/1/1
3	BOG	A	403	-	-	0/11/31/31	0/1/1/1
4	MES	A	404	-	-	0/6/14/14	0/1/1/1
2	R48	B	401	-	-	0/20/22/22	0/3/5/5
3	BOG	B	402	-	-	0/11/31/31	0/1/1/1
3	BOG	B	403	-	-	0/11/31/31	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	R48	N1V-N2K	-7.13	1.26	1.39
2	B	401	R48	N1V-N2K	-7.00	1.26	1.39
4	A	404	MES	C8-S	-5.21	1.69	1.78
2	A	401	R48	C2E-N2K	-4.05	1.35	1.43
2	B	401	R48	C2E-N2K	-3.92	1.35	1.43
2	A	401	R48	C2C-N1W	-3.36	1.35	1.41
4	A	404	MES	O1S-S	3.12	1.52	1.45
2	B	401	R48	C6-C5	-2.93	1.41	1.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	BOG	O1-C1	2.87	1.45	1.40
2	A	401	R48	C2G-N2K	-2.85	1.34	1.36
2	A	401	R48	C6-C5	-2.79	1.41	1.45
2	B	401	R48	C1P-C1M	2.75	1.42	1.36
3	A	403	BOG	O1-C1	2.50	1.44	1.40
2	B	401	R48	C2D-N1Y	-2.48	1.35	1.40
2	A	401	R48	C2L-C2F	2.46	1.56	1.52
2	A	401	R48	C1R-C2F	-2.43	1.36	1.39
2	B	401	R48	C2C-N1W	-2.21	1.37	1.41
2	A	401	R48	C1P-C1M	2.18	1.41	1.36
4	A	404	MES	O3S-S	2.09	1.50	1.45
3	A	403	BOG	O5-C1	2.07	1.47	1.41
3	B	402	BOG	O5-C5	2.06	1.49	1.44
2	B	401	R48	C1O-C1L	2.05	1.42	1.38
2	B	401	R48	C1R-C2F	-2.03	1.36	1.39

All (54) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	R48	N3-C2-N1	-9.34	120.90	128.71
2	B	401	R48	C2-N1-C6	8.59	122.09	116.69
2	B	401	R48	N3-C2-N1	-8.58	121.54	128.71
2	A	401	R48	C2-N1-C6	7.57	121.44	116.69
4	A	404	MES	C5-N4-C3	7.29	124.22	109.75
2	A	401	R48	C2-N3-C4	7.11	120.88	115.37
2	A	401	R48	C1R-C2G-N2K	-6.83	105.64	106.68
2	B	401	R48	C2-N3-C4	6.57	120.45	115.37
2	B	401	R48	N1X-C2G-N2K	6.45	127.59	120.99
2	B	401	R48	C1S-C5-C6	-6.35	121.57	124.98
2	A	401	R48	N1X-C2G-N2K	5.89	127.02	120.99
2	A	401	R48	C2E-N2K-N1V	5.71	123.26	118.85
3	B	402	BOG	O1-C1-C2	5.63	115.35	108.18
2	A	401	R48	C1S-C5-C6	-5.56	121.99	124.98
2	B	401	R48	C2E-N2K-N1V	5.53	123.13	118.85
2	A	401	R48	C2L-C2F-N1V	5.37	126.27	120.64
2	B	401	R48	C5-C6-N1	-5.01	117.88	121.31
2	B	401	R48	C6-C5-C4	5.01	118.48	115.63
2	A	401	R48	C6-C5-C4	4.34	118.10	115.63
2	A	401	R48	C5-C6-N1	-4.18	118.45	121.31
3	B	403	BOG	O1-C1-C2	3.94	113.20	108.18
2	B	401	R48	C2F-N1V-N2K	3.70	109.95	105.66
3	B	403	BOG	O5-C5-C4	3.66	116.54	109.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	R48	C2L-C2F-N1V	3.66	124.48	120.64
2	A	401	R48	C1O-C2E-N2K	3.59	124.71	119.62
4	A	404	MES	C7-N4-C3	3.53	120.89	111.66
3	B	403	BOG	O5-C1-C2	3.49	117.46	110.31
4	A	404	MES	C6-C5-N4	3.47	114.42	109.96
3	B	402	BOG	C6-C5-C4	3.40	121.23	113.00
2	B	401	R48	C1O-C2E-N2K	3.34	124.36	119.62
2	B	401	R48	N1Y-C6-N1	3.30	122.28	118.92
2	A	401	R48	C1D-C2L-C2F	3.27	116.81	109.88
3	B	402	BOG	C1'-O1-C1	-3.24	108.13	113.96
2	B	401	R48	C5-C4-N3	-3.23	119.54	122.91
2	B	401	R48	N1X-C1Z-N1W	2.80	117.03	112.57
2	A	401	R48	C2F-N1V-N2K	2.80	108.91	105.66
2	A	401	R48	C1N-C2E-N2K	-2.80	115.64	119.62
2	A	401	R48	C1R-C2F-N1V	-2.77	107.35	111.48
4	A	404	MES	O3S-S-O1S	-2.66	105.63	112.48
3	A	403	BOG	O5-C5-C4	2.62	114.61	109.76
2	B	401	R48	C1R-C2F-N1V	-2.60	107.60	111.48
3	B	402	BOG	O5-C1-O1	-2.59	103.89	109.98
2	A	401	R48	N1Y-C6-N1	2.59	121.56	118.92
2	A	401	R48	C5-C4-N3	-2.57	120.22	122.91
2	B	401	R48	C1B-C2L-C2F	2.55	115.28	109.88
3	B	403	BOG	C1-C2-C3	2.52	114.90	110.00
3	B	403	BOG	C4-C3-C2	2.34	115.13	110.82
4	A	404	MES	O2S-S-C8	2.30	113.45	106.36
3	B	403	BOG	C1-O5-C5	2.25	118.10	113.73
2	A	401	R48	C2E-N2K-C2G	-2.24	125.45	128.50
2	A	401	R48	C2D-N1Y-C6	-2.19	124.24	128.67
3	A	402	BOG	C1'-O1-C1	-2.16	110.07	113.96
3	B	403	BOG	C1'-O1-C1	-2.15	110.09	113.96
2	A	401	R48	C1R-C2F-C2L	-2.10	126.36	128.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/360 (94%)	-0.10	10 (2%) 49 54	20, 28, 47, 56	0
1	B	337/360 (93%)	0.14	16 (4%) 30 33	19, 29, 49, 58	0
All	All	676/720 (93%)	0.02	26 (3%) 38 43	19, 28, 48, 58	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	183	VAL	7.4
1	A	172	ALA	7.0
1	B	329	SER	4.5
1	B	34	ALA	4.4
1	A	119	CYS	4.2
1	B	198	MET	3.8
1	B	5	ARG	3.6
1	B	14	ASN	3.6
1	B	169	PHE	3.2
1	B	119	CYS	3.0
1	B	110	GLY	3.0
1	A	198	MET	2.9
1	B	352	PRO	2.9
1	B	15	LYS	2.8
1	B	331	ASP	2.8
1	B	171	LEU	2.8
1	A	35	TYR	2.5
1	A	110	GLY	2.4
1	B	97	GLU	2.4
1	A	199	HIS	2.4
1	B	98	GLU	2.3
1	A	118	LYS	2.2
1	A	352	PRO	2.2
1	B	62	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	245	GLU	2.0
1	B	170	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	BOG	B	403	20/20	0.27	4.85	51,65,69,70	0
3	BOG	A	403	20/20	0.31	2.62	59,64,67,68	0
4	MES	A	404	12/12	0.16	2.54	26,33,35,37	12
2	R48	A	401	38/38	0.14	-0.10	18,21,27,27	0
3	BOG	B	402	20/20	0.10	-0.29	25,28,34,35	0
3	BOG	A	402	20/20	0.10	-0.43	30,34,39,41	0
2	R48	B	401	38/38	0.12	-0.58	17,22,30,31	0

## 6.5 Other polymers

There are no such residues in this entry.