



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 03:29 AM GMT

PDB ID : 1GD8
Title : THE CRYSTAL STRUCTURE OF BACTERIA-SPECIFIC L17 RIBOSOMAL PROTEIN.
Authors : Vassilyev, D.G.; Shirouzu, M.; Wada, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2000-09-22
Resolution : 2.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	118	
1	B	118	
1	C	118	
1	D	118	
1	E	118	
1	F	118	
1	G	118	
1	H	118	
1	I	118	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7852 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S RIBOSOMAL PROTEIN L17.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	B	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	C	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	D	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	E	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	F	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	G	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	H	105	Total	C	N	O	0	0	0
			855	536	174	145			
1	I	105	Total	C	N	O	0	0	0
			855	536	174	145			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total	O	0	0
			36	36		
2	B	15	Total	O	0	0
			15	15		
2	C	16	Total	O	0	0
			16	16		
2	D	27	Total	O	0	0
			27	27		
2	E	13	Total	O	0	0
			13	13		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	F	4	Total 4	O 4	0	0
2	G	19	Total 19	O 19	0	0
2	H	13	Total 13	O 13	0	0
2	I	14	Total 14	O 14	0	0


i

• Molecule 1: 50S RIBOSOMAL PROTEIN L17




● Molecule 1: 50S RIBOSOMAL PROTEIN L17

Chain E:




● Molecule 1: 50S RIBOSOMAL PROTEIN L17

Chain F:




● Molecule 1: 50S RIBOSOMAL PROTEIN L17

Chain G:




● Molecule 1: 50S RIBOSOMAL PROTEIN L17

Chain H:



● Molecule 1: 50S RIBOSOMAL PROTEIN L17

Chain I:



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	109.24Å 109.24Å 128.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 39.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.2 (50.00-2.30) 97.2 (39.97-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11276.44 (at 2.29Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.210 , 0.254 0.211 , 0.252	Depositor DCC
R_{free} test set	3065 reflections (4.94%)	DCC
Wilson B-factor (Å ²)	49.7	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.0	EDS
Estimated twinning fraction	0.337 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtriage
Outliers	0 of 65159 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7852	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/867	0.70	0/1162
1	B	0.41	0/867	0.72	0/1162
1	C	0.38	0/867	0.72	0/1162
1	D	0.42	0/867	0.73	0/1162
1	E	0.39	0/867	0.65	0/1162
1	F	0.39	0/867	0.64	0/1162
1	G	0.42	0/867	0.69	0/1162
1	H	0.39	0/867	0.66	0/1162
1	I	0.38	0/867	0.62	0/1162
All	All	0.40	0/7803	0.68	0/10458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	906	62	0
1	B	855	0	906	57	0
1	C	855	0	906	67	0
1	D	855	0	906	55	0
1	E	855	0	906	77	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	855	0	906	65	0
1	G	855	0	906	101	0
1	H	855	0	906	110	0
1	I	855	0	906	71	0
2	A	36	0	0	5	0
2	B	15	0	0	1	0
2	C	16	0	0	0	0
2	D	27	0	0	3	0
2	E	13	0	0	5	0
2	F	4	0	0	0	0
2	G	19	0	0	5	0
2	H	13	0	0	1	0
2	I	14	0	0	0	0
All	All	7852	0	8154	603	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 38.

The worst 5 of 603 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:G:68:ARG:HD2	1:H:68:ARG:HD3	1.36	1.05
1:F:74:LYS:HD3	1:F:74:LYS:H	1.17	1.05
1:C:78:LYS:HE2	1:C:83:ILE:HD11	1.34	1.04
1:G:26:LYS:HG3	1:G:75:LEU:HD11	1.35	1.03
1:E:60:LEU:HD23	1:E:60:LEU:H	1.23	1.03

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/118 (87%)	99 (96%)	3 (3%)	1 (1%)	22	23
1	B	103/118 (87%)	98 (95%)	4 (4%)	1 (1%)	22	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	103/118 (87%)	96 (93%)	5 (5%)	2 (2%)	12	9
1	D	103/118 (87%)	94 (91%)	8 (8%)	1 (1%)	22	23
1	E	103/118 (87%)	97 (94%)	4 (4%)	2 (2%)	12	9
1	F	103/118 (87%)	98 (95%)	3 (3%)	2 (2%)	12	9
1	G	103/118 (87%)	98 (95%)	5 (5%)	0	100	100
1	H	103/118 (87%)	92 (89%)	10 (10%)	1 (1%)	22	23
1	I	103/118 (87%)	96 (93%)	5 (5%)	2 (2%)	12	9
All	All	927/1062 (87%)	868 (94%)	47 (5%)	12 (1%)	18	17

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	105	ARG
1	F	58	GLY
1	I	104	ARG
1	I	58	GLY
1	H	104	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/101 (88%)	69 (78%)	20 (22%)	1	1
1	B	89/101 (88%)	75 (84%)	14 (16%)	4	3
1	C	89/101 (88%)	72 (81%)	17 (19%)	2	2
1	D	89/101 (88%)	78 (88%)	11 (12%)	7	7
1	E	89/101 (88%)	72 (81%)	17 (19%)	2	2
1	F	89/101 (88%)	75 (84%)	14 (16%)	4	3
1	G	89/101 (88%)	72 (81%)	17 (19%)	2	2
1	H	89/101 (88%)	70 (79%)	19 (21%)	1	1
1	I	89/101 (88%)	75 (84%)	14 (16%)	4	3
All	All	801/909 (88%)	658 (82%)	143 (18%)	2	2

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	20	LEU
1	F	20	LEU
1	I	34	ILE
1	E	50	HIS
1	E	73	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	24	GLN
1	F	53	HIS
1	G	91	GLN
1	D	53	HIS
1	H	53	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/118 (88%)	1.05	8 (7%) 14 20	28, 44, 59, 71	0
1	B	105/118 (88%)	1.22	21 (20%) 2 3	36, 50, 63, 67	0
1	C	105/118 (88%)	1.40	19 (18%) 2 3	34, 51, 63, 71	0
1	D	105/118 (88%)	1.23	15 (14%) 3 5	25, 42, 54, 66	0
1	E	105/118 (88%)	1.03	12 (11%) 6 9	37, 52, 63, 70	0
1	F	105/118 (88%)	1.16	13 (12%) 5 7	44, 55, 63, 68	0
1	G	105/118 (88%)	1.37	19 (18%) 2 3	37, 55, 65, 67	0
1	H	105/118 (88%)	1.83	40 (38%) 1 1	37, 58, 68, 70	0
1	I	105/118 (88%)	1.48	27 (25%) 1 2	37, 54, 65, 76	0
All	All	945/1062 (88%)	1.31	174 (18%) 2 3	25, 52, 65, 76	0

The worst 5 of 174 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	16	HIS	8.0
1	C	16	HIS	7.1
1	C	107	ASP	7.0
1	I	60	LEU	6.7
1	B	16	HIS	6.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.