



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 01:06 AM GMT

PDB ID : 3GD5  
Title : Crystal structure of ornithine carbamoyltransferase from *Gloeobacter violaceus*  
Authors : Fedorov, A.A.; Fedorov, E.V.; Toro, R.; Ramagopal, U.A.; Sauder, J.M.; Bursley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-02-23  
Resolution : 2.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	323	
1	B	323	
1	C	323	
1	D	323	
1	E	323	
1	F	323	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 13303 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	2195	1383	399	405	8	0	0	0
1	B	285	2195	1383	399	405	8	0	0	0
1	C	285	2195	1383	399	405	8	0	0	0
1	D	285	2195	1383	399	405	8	0	0	0
1	E	285	2195	1383	399	405	8	0	0	0
1	F	285	2195	1383	399	405	8	0	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
A	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
A	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
A	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7
A	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
A	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7
A	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
A	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
A	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
A	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
A	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7
B	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
B	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
B	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
B	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7
B	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
B	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
B	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
B	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
B	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
B	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
C	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
C	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
C	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7
C	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
C	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
C	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
D	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
D	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
D	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7
D	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
D	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
D	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
E	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
E	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
E	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7
E	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
E	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
E	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	1	MET	-	EXPRESSION TAG	UNP Q7NGR7
F	2	SER	-	EXPRESSION TAG	UNP Q7NGR7
F	3	LEU	-	EXPRESSION TAG	UNP Q7NGR7
F	316	GLU	-	EXPRESSION TAG	UNP Q7NGR7

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
F	317	GLY	-	EXPRESSION TAG	UNP Q7NGR7
F	318	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	319	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	320	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	321	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	322	HIS	-	EXPRESSION TAG	UNP Q7NGR7
F	323	HIS	-	EXPRESSION TAG	UNP Q7NGR7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	17	Total O 17 17	0	0
2	B	37	Total O 37 37	0	0
2	C	34	Total O 34 34	0	0
2	D	8	Total O 8 8	0	0
2	E	23	Total O 23 23	0	0
2	F	14	Total O 14 14	0	0







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.56Å 142.42Å 149.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.94 – 2.10 39.77 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (24.94-2.10) 99.8 (39.77-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.239 , 0.271 0.239 , 0.271	Depositor DCC
$R_{free}$ test set	6192 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.8	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.2	EDS
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 123458 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13303	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/2233	0.60	1/3031 (0.0%)
1	B	0.39	0/2233	0.64	1/3031 (0.0%)
1	C	0.38	0/2233	0.62	1/3031 (0.0%)
1	D	0.33	0/2233	0.59	0/3031
1	E	0.34	0/2233	0.59	0/3031
1	F	0.35	0/2233	0.61	0/3031
All	All	0.35	0/13398	0.61	3/18186 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	160	LEU	CA-CB-CG	5.65	128.30	115.30
1	A	160	LEU	CA-CB-CG	5.22	127.30	115.30
1	B	160	LEU	CA-CB-CG	5.08	126.98	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2195	0	2198	96	0
1	B	2195	0	2198	114	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2195	0	2198	85	0
1	D	2195	0	2198	101	0
1	E	2195	0	2198	87	0
1	F	2195	0	2198	85	0
2	A	17	0	0	3	0
2	B	37	0	0	4	0
2	C	34	0	0	1	0
2	D	8	0	0	3	0
2	E	23	0	0	1	0
2	F	14	0	0	1	0
All	All	13303	0	13188	548	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 21.

The worst 5 of 548 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:54:VAL:HG13	1:D:109:ILE:HB	1.27	1.12
1:A:54:VAL:HG13	1:A:109:ILE:HB	1.30	1.10
1:C:54:VAL:HG13	1:C:109:ILE:HB	1.34	1.06
1:F:56:LEU:HD22	1:F:111:THR:HG23	1.40	1.03
1:B:145:LEU:HD11	1:B:182:MET:CE	1.93	0.97

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/323 (86%)	266 (95%)	11 (4%)	2 (1%)	30 23
1	B	279/323 (86%)	267 (96%)	11 (4%)	1 (0%)	43 39
1	C	279/323 (86%)	268 (96%)	10 (4%)	1 (0%)	43 39
1	D	279/323 (86%)	259 (93%)	18 (6%)	2 (1%)	30 23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	279/323 (86%)	266 (95%)	11 (4%)	2 (1%)	30 23
1	F	279/323 (86%)	268 (96%)	10 (4%)	1 (0%)	43 39
All	All	1674/1938 (86%)	1594 (95%)	71 (4%)	9 (0%)	38 33

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	132	LEU
1	C	132	LEU
1	D	132	LEU
1	E	132	LEU
1	F	132	LEU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	225/254 (89%)	210 (93%)	15 (7%)	23 18
1	B	225/254 (89%)	208 (92%)	17 (8%)	19 14
1	C	225/254 (89%)	209 (93%)	16 (7%)	21 16
1	D	225/254 (89%)	205 (91%)	20 (9%)	14 9
1	E	225/254 (89%)	211 (94%)	14 (6%)	26 21
1	F	225/254 (89%)	202 (90%)	23 (10%)	11 6
All	All	1350/1524 (89%)	1245 (92%)	105 (8%)	18 13

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	266	GLU
1	D	145	LEU
1	F	160	LEU
1	D	17	LEU
1	D	96	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	227	HIS
1	D	137	HIS
1	F	151	ASN
1	C	276	HIS
1	D	151	ASN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/323 (88%)	0.26	14 (4%) 28 31	33, 45, 63, 77	0
1	B	285/323 (88%)	-0.04	6 (2%) 60 65	26, 36, 54, 67	0
1	C	285/323 (88%)	-0.14	1 (0%) 90 92	26, 36, 51, 63	0
1	D	285/323 (88%)	0.48	19 (6%) 17 19	32, 50, 67, 73	0
1	E	285/323 (88%)	0.35	15 (5%) 25 28	30, 41, 63, 82	0
1	F	285/323 (88%)	0.15	12 (4%) 35 39	29, 41, 61, 74	0
All	All	1710/1938 (88%)	0.18	67 (3%) 37 42	26, 41, 62, 82	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	248	GLN	5.0
1	D	11	ARG	4.7
1	B	182	MET	4.2
1	D	122	HIS	3.6
1	E	11	ARG	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.