



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 01:35 PM GMT

PDB ID : 4GDK  
Title : Crystal Structure of Human Atg12 Atg5 Conjugate in Complex with an N-terminal Fragment of Atg16L1  
Authors : Otomo, C.; Metlagel, Z.; Takaesu, G.; Otomo, T.  
Deposited on : 2012-07-31  
Resolution : 2.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	91	
1	D	91	
2	B	275	
2	E	275	
3	C	36	
3	F	36	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6547 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-like protein ATG12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	88	Total	C	N	O	S	0	0	0
			704	461	115	125	3			
1	D	87	Total	C	N	O	S	0	0	0
			695	455	113	124	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	GLY	-	EXPRESSION TAG	UNP O94817
A	51	SER	-	EXPRESSION TAG	UNP O94817
D	50	GLY	-	EXPRESSION TAG	UNP O94817
D	51	SER	-	EXPRESSION TAG	UNP O94817

- Molecule 2 is a protein called Autophagy protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	267	Total	C	N	O	S	0	0	0
			2227	1444	369	402	12			
2	E	265	Total	C	N	O	S	0	0	0
			2211	1436	367	396	12			

- Molecule 3 is a protein called Autophagy-related protein 16-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	34	Total	C	N	O	S	0	0	0
			311	196	65	49	1			
3	F	35	Total	C	N	O	S	0	0	0
			321	202	68	50	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	8	SER	-	EXPRESSION TAG	UNP Q676U5
C	9	HIS	-	EXPRESSION TAG	UNP Q676U5
C	10	MET	-	EXPRESSION TAG	UNP Q676U5
F	8	SER	-	EXPRESSION TAG	UNP Q676U5
F	9	HIS	-	EXPRESSION TAG	UNP Q676U5
F	10	MET	-	EXPRESSION TAG	UNP Q676U5

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Na 1 1	0	0
4	E	1	Total Na 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total O 3 3	0	0
5	B	44	Total O 44 44	0	0
5	C	7	Total O 7 7	0	0
5	D	1	Total O 1 1	0	0
5	E	13	Total O 13 13	0	0
5	F	8	Total O 8 8	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.76Å 113.15Å 210.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.06 – 2.70 44.06 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.06-2.70) 99.5 (44.06-2.70)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 2.69Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1069)	Depositor
R, $R_{free}$	0.176 , 0.236 0.180 , 0.240	Depositor DCC
$R_{free}$ test set	2000 reflections (6.76%)	DCC
Wilson B-factor (Å <sup>2</sup> )	61.1	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 36.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 29602 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6547	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/719	0.38	0/968
1	D	0.22	0/710	0.38	0/957
2	B	0.24	0/2293	0.41	0/3110
2	E	0.23	0/2277	0.39	0/3088
3	C	0.24	0/316	0.39	0/420
3	F	0.22	0/327	0.36	0/435
All	All	0.23	0/6642	0.39	0/8978

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	704	0	0	0	0
1	D	695	0	0	0	0
2	B	2227	0	0	7	0
2	E	2211	0	0	4	0
3	C	311	0	0	0	0
3	F	321	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	E	1	0	0	0	0
5	A	3	0	0	0	0
5	B	44	0	0	1	0
5	C	7	0	0	0	0
5	D	1	0	0	0	0
5	E	13	0	0	0	0
5	F	8	0	0	0	0
All	All	6547	0	0	11	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (11) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:275:ASP:OD2	3:F:26:ARG:NH1	2.35	0.59
2:E:50:ASP:OD2	2:E:118:LYS:NZ	2.42	0.53
2:E:109:GLU:OE1	2:E:114:HIS:NE2	2.42	0.51
2:B:61:ARG:NE	2:B:63:GLU:OE1	2.44	0.50
2:B:66:SER:OG	2:B:67:GLU:N	2.43	0.50
2:B:136:LYS:NZ	2:B:263:ASN:OD1	2.45	0.48
2:B:178:GLU:N	2:B:178:GLU:OE1	2.48	0.47
2:B:192:THR:N	5:B:431:HOH:O	2.47	0.46
2:B:204:ARG:O	2:B:213:HIS:ND1	2.53	0.42
2:E:133:ASP:OD1	2:E:137:HIS:ND1	2.53	0.41
2:E:237:GLN:N	2:E:270:ILE:O	2.54	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	86/91 (94%)	82 (95%)	4 (5%)	0	100 100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	85/91 (93%)	83 (98%)	2 (2%)	0	100	100
2	B	263/275 (96%)	257 (98%)	6 (2%)	0	100	100
2	E	261/275 (95%)	252 (97%)	9 (3%)	0	100	100
3	C	32/36 (89%)	31 (97%)	1 (3%)	0	100	100
3	F	33/36 (92%)	32 (97%)	1 (3%)	0	100	100
All	All	760/804 (94%)	737 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/79 (98%)	77 (100%)	0	100	100
1	D	76/79 (96%)	76 (100%)	0	100	100
2	B	247/254 (97%)	246 (100%)	1 (0%)	95	99
2	E	245/254 (96%)	245 (100%)	0	100	100
3	C	33/35 (94%)	33 (100%)	0	100	100
3	F	34/35 (97%)	34 (100%)	0	100	100
All	All	712/736 (97%)	711 (100%)	1 (0%)	96	99

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	71	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	88/91 (96%)	0.01	1 (1%) 77 82	55, 71, 92, 112	0
1	D	87/91 (95%)	0.22	1 (1%) 77 82	62, 77, 98, 112	0
2	B	267/275 (97%)	-0.08	1 (0%) 90 93	36, 56, 90, 115	0
2	E	265/275 (96%)	0.14	9 (3%) 43 48	41, 67, 106, 127	0
3	C	34/36 (94%)	-0.07	0 100 100	41, 57, 85, 94	0
3	F	35/36 (97%)	-0.13	0 100 100	41, 64, 98, 119	0
All	All	776/804 (96%)	0.04	12 (1%) 70 75	36, 65, 100, 127	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	27	ILE	3.5
2	E	30	ARG	3.5
2	E	61	ARG	3.3
2	B	227	ILE	3.1
2	E	209	ASP	2.8
2	E	210	GLY	2.5
2	E	212	LEU	2.3
2	E	54	LYS	2.3
1	D	64	ASP	2.3
2	E	65	ILE	2.1
2	E	208	ALA	2.1
1	A	58	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	E	301	1/1	0.11	-0.77	65,65,65,65	0
4	NA	B	301	1/1	0.10	-1.69	52,52,52,52	0

### 6.5 Other polymers

There are no such residues in this entry.