



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 01:35 PM GMT

PDB ID : 4GDO  
Title : Structure of a fragment of the rod domain of plectin  
Authors : De Pereda, J.M.; Buey, R.M.; Uson, I.; Sammito, M.D.; De Marino, I.  
Deposited on : 2012-08-01  
Resolution : 1.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	43	
1	B	43	
1	C	43	
1	D	43	
1	E	43	
1	F	43	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3410 atoms, of which 1594 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	42	Total 666	C 200	H 329	N 68	O 68	S 1	0	0	0
1	B	41	Total 655	C 197	H 324	N 67	O 66	S 1	0	0	0
1	C	38	Total 571	C 173	H 283	N 56	O 59		0	0	0
1	D	41	Total 619	C 186	H 304	N 64	O 64	S 1	0	0	0
1	E	31	Total 388	C 121	H 188	N 40	O 39		0	0	0
1	F	30	Total 348	C 110	H 166	N 36	O 36		0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1378	GLY	-	EXPRESSION TAG	UNP Q15149
A	1379	SER	-	EXPRESSION TAG	UNP Q15149
A	1380	HIS	-	EXPRESSION TAG	UNP Q15149
A	1381	MET	-	EXPRESSION TAG	UNP Q15149
B	1378	GLY	-	EXPRESSION TAG	UNP Q15149
B	1379	SER	-	EXPRESSION TAG	UNP Q15149
B	1380	HIS	-	EXPRESSION TAG	UNP Q15149
B	1381	MET	-	EXPRESSION TAG	UNP Q15149
C	1378	GLY	-	EXPRESSION TAG	UNP Q15149
C	1379	SER	-	EXPRESSION TAG	UNP Q15149
C	1380	HIS	-	EXPRESSION TAG	UNP Q15149
C	1381	MET	-	EXPRESSION TAG	UNP Q15149
D	1378	GLY	-	EXPRESSION TAG	UNP Q15149
D	1379	SER	-	EXPRESSION TAG	UNP Q15149
D	1380	HIS	-	EXPRESSION TAG	UNP Q15149
D	1381	MET	-	EXPRESSION TAG	UNP Q15149
E	1378	GLY	-	EXPRESSION TAG	UNP Q15149

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	1379	SER	-	EXPRESSION TAG	UNP Q15149
E	1380	HIS	-	EXPRESSION TAG	UNP Q15149
E	1381	MET	-	EXPRESSION TAG	UNP Q15149
F	1378	GLY	-	EXPRESSION TAG	UNP Q15149
F	1379	SER	-	EXPRESSION TAG	UNP Q15149
F	1380	HIS	-	EXPRESSION TAG	UNP Q15149
F	1381	MET	-	EXPRESSION TAG	UNP Q15149

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	48	Total O 48 48	0	0
2	B	41	Total O 41 41	0	0
2	C	26	Total O 26 26	0	0
2	D	39	Total O 39 39	0	0
2	E	2	Total O 2 2	0	0
2	F	7	Total O 7 7	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Plectin

Chain A:




- Molecule 1: Plectin

Chain B:




- Molecule 1: Plectin

Chain C:




- Molecule 1: Plectin

Chain D:



- Molecule 1: Plectin

Chain E:



- Molecule 1: Plectin

Chain F:



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.62Å 35.87Å 75.69Å 90.00° 94.11° 90.00°	Depositor
Resolution (Å)	45.19 – 1.70 45.19 – 1.70	Depositor EDS
% Data completeness (in resolution range)	85.3 (45.19-1.70) 85.4 (45.19-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1117)	Depositor
R, $R_{free}$	0.213 , 0.248 0.216 , 0.252	Depositor DCC
$R_{free}$ test set	1188 reflections (5.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtriage
Anisotropy	0.500	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 23087 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3410	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/338	0.54	0/447
1	B	0.46	0/332	0.54	0/439
1	C	0.40	0/288	0.47	0/383
1	D	0.43	0/315	0.57	0/418
1	E	0.28	0/200	0.34	0/270
1	F	0.26	0/182	0.28	0/247
All	All	0.42	0/1655	0.49	0/2204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	337	329	0	7	0
1	B	331	324	0	5	0
1	C	288	283	0	5	0
1	D	315	304	0	3	0
1	E	200	188	0	2	0
1	F	182	166	0	3	0
2	A	48	0	0	1	0
2	B	41	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	26	0	0	0	0
2	D	39	0	0	2	0
2	E	2	0	0	0	0
2	F	7	0	0	0	0
All	All	1816	1594	0	17	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:1380:HIS:N	2:B:1528:HOH:O	2.22	0.72
1:D:1382:ARG:NH2	2:D:1520:HOH:O	2.26	0.68
1:C:1419:GLU:OE2	1:D:1388:ARG:NH1	2.35	0.60
1:A:1398:LYS:NZ	2:A:1514:HOH:O	2.36	0.59
1:A:1380:HIS:NE2	1:C:1393:GLU:OE1	2.40	0.55

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	40/43 (93%)	40 (100%)	0	0	100	100
1	B	39/43 (91%)	39 (100%)	0	0	100	100
1	C	36/43 (84%)	36 (100%)	0	0	100	100
1	D	39/43 (91%)	39 (100%)	0	0	100	100
1	E	29/43 (67%)	29 (100%)	0	0	100	100
1	F	28/43 (65%)	28 (100%)	0	0	100	100
All	All	211/258 (82%)	211 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/31 (100%)	30 (97%)	1 (3%)	51	26
1	B	30/31 (97%)	30 (100%)	0	100	100
1	C	24/31 (77%)	24 (100%)	0	100	100
1	D	26/31 (84%)	26 (100%)	0	100	100
1	E	11/31 (36%)	10 (91%)	1 (9%)	14	3
1	F	8/31 (26%)	6 (75%)	2 (25%)	1	0
All	All	130/186 (70%)	126 (97%)	4 (3%)	52	27

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1382	ARG
1	E	1412	GLN
1	F	1397	GLU
1	F	1416	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1401	GLN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	42/43 (97%)	0.46	3 (7%) 16 19	12, 21, 49, 75	0
1	B	41/43 (95%)	0.41	2 (4%) 28 33	13, 22, 57, 70	0
1	C	38/43 (88%)	0.88	5 (13%) 4 5	15, 30, 101, 138	0
1	D	41/43 (95%)	0.87	8 (19%) 2 2	14, 27, 90, 104	0
1	E	31/43 (72%)	2.52	19 (61%) 0 2	59, 68, 100, 125	0
1	F	30/43 (69%)	4.16	27 (90%) 0 1	64, 94, 124, 150	0
All	All	223/258 (86%)	1.38	64 (28%) 1 2	12, 35, 108, 150	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	1402	LEU	8.2
1	F	1411	ALA	8.2
1	F	1404	GLU	8.1
1	F	1394	ALA	7.8
1	E	1405	ALA	7.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.