



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 01:12 AM GMT

PDB ID : 3GW6  
Title : Intramolecular Chaperone  
Authors : Schulz, E.C.; Dickmanns, A.; Ficner, R.  
Deposited on : 2009-03-31  
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	275	
1	B	275	
1	C	275	
1	D	275	
1	E	275	
1	F	275	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	PEG	A	17	-	X
2	PEG	C	29	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
2	PEG	C	3	-	X
2	PEG	D	15	-	X
2	PEG	E	16	-	X
2	PEG	E	18	-	X
2	PEG	E	22	-	X
2	PEG	E	27	-	X
2	PEG	E	33	-	X
2	PEG	E	6	-	X
2	PEG	F	24	-	X
2	PEG	F	30	-	X
2	PEG	F	5	-	X
3	TAM	A	32	-	X
3	TAM	A	35	-	X
3	TAM	C	28	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12617 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-N-acetylneuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	16	0	0
			2028	1270	351	400	7			
1	B	255	Total	C	N	O	S	24	0	0
			2020	1267	349	397	7			
1	E	257	Total	C	N	O	S	27	0	0
			2032	1272	351	402	7			
1	C	254	Total	C	N	O	S	11	0	0
			2011	1262	349	393	7			
1	D	252	Total	C	N	O	S	34	0	0
			1996	1253	346	390	7			
1	F	257	Total	C	N	O	S	24	0	0
			2035	1274	352	402	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	911	ALA	SER	CONFLICT	UNP Q858B1
B	911	ALA	SER	CONFLICT	UNP Q858B1
E	911	ALA	SER	CONFLICT	UNP Q858B1
C	911	ALA	SER	CONFLICT	UNP Q858B1
D	911	ALA	SER	CONFLICT	UNP Q858B1
F	911	ALA	SER	CONFLICT	UNP Q858B1

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		
2	D	1	Total	C	O	0	0
			7	4	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		
2	F	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: C<sub>7</sub>H<sub>17</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	1	0
			11	7	1	3		
3	A	1	Total	C	N	O	0	0
			11	7	1	3		
3	C	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	D	2	Total	Cl	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total 2	Cl 2	0	0
4	E	1	Total 1	Cl 1	0	0

- Molecule 5 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	3	Total 3	Br 3	0	0
5	C	1	Total 1	Br 1	0	0
5	F	1	Total 1	Br 1	0	0
5	E	1	Total 1	Br 1	0	0

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	1	Total 1	Ca 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	63	Total 63	O 63	0	0
7	B	47	Total 47	O 47	0	0
7	E	45	Total 45	O 45	0	0
7	C	61	Total 61	O 61	0	0
7	D	52	Total 52	O 52	0	0
7	F	55	Total 55	O 55	0	0









• Molecule 1: Endo-N-acetylneuraminidase

Chain C:




• Molecule 1: Endo-N-acetylneuraminidase

Chain D:



• Molecule 1: Endo-N-acetylneuraminidase

Chain F:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.65Å 79.95Å 109.58Å 81.71° 76.53° 87.11°	Depositor
Resolution (Å)	27.63 – 2.60 48.21 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.4 (27.63-2.60) 95.4 (48.21-2.60)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.209 , 0.255 0.204 , 0.249	Depositor DCC
$R_{free}$ test set	2997 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.7	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 42.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 59993 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12617	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TAM, CA, PEG, BR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/2071	0.43	0/2810
1	B	0.24	0/2062	0.44	0/2797
1	C	0.24	0/2053	0.43	0/2785
1	D	0.25	0/2039	0.47	0/2767
1	E	0.23	0/2074	0.45	0/2815
1	F	0.23	0/2078	0.43	0/2820
All	All	0.24	0/12377	0.44	0/16794

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2028	0	1425	54	0
1	B	2020	0	1416	59	0
1	C	2011	0	1416	55	0
1	D	1996	0	1418	59	0
1	E	2032	0	1429	58	0
1	F	2035	0	1425	51	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	14	0	20	5	0
2	B	7	0	10	0	0
2	C	21	0	30	5	0
2	D	7	0	10	0	0
2	E	49	0	70	14	0
2	F	28	0	40	14	0
3	A	22	0	34	3	0
3	C	11	0	17	1	0
4	A	1	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
4	E	1	0	0	0	0
5	B	3	0	0	0	0
5	C	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	F	1	0	0	0	0
7	A	63	0	0	3	0
7	B	47	0	0	0	0
7	C	61	0	0	3	0
7	D	52	0	0	2	0
7	E	45	0	0	1	0
7	F	55	0	0	3	0
All	All	12617	0	8760	281	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:845:GLY:H	1:B:850:THR:HG21	1.10	1.13
1:A:845:GLY:H	1:A:850:THR:HG21	1.18	1.06
1:F:845:GLY:H	1:F:850:THR:HG21	1.18	1.04
1:D:845:GLY:H	1:D:850:THR:HG21	1.23	1.04
3:A:35:TAM:H61	7:A:241:HOH:O	1.83	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/275 (91%)	237 (95%)	12 (5%)	1 (0%)	43	72
1	B	247/275 (90%)	237 (96%)	10 (4%)	0	100	100
1	C	246/275 (90%)	237 (96%)	9 (4%)	0	100	100
1	D	246/275 (90%)	238 (97%)	8 (3%)	0	100	100
1	E	251/275 (91%)	241 (96%)	10 (4%)	0	100	100
1	F	251/275 (91%)	244 (97%)	7 (3%)	0	100	100
All	All	1491/1650 (90%)	1434 (96%)	56 (4%)	1 (0%)	59	85

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	975	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/230 (94%)	194 (90%)	21 (10%)	12	21
1	B	214/230 (93%)	194 (91%)	20 (9%)	13	24
1	C	213/230 (93%)	199 (93%)	14 (7%)	24	45
1	D	211/230 (92%)	192 (91%)	19 (9%)	14	26
1	E	215/230 (94%)	196 (91%)	19 (9%)	14	27
1	F	216/230 (94%)	206 (95%)	10 (5%)	37	66
All	All	1284/1380 (93%)	1181 (92%)	103 (8%)	17	33

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	927	LEU
1	E	1051	LEU
1	F	927	LEU
1	E	936	ILE
1	E	995	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	912	ASN
1	C	999	HIS
1	F	969	HIS
1	E	969	HIS
1	D	800	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 34 ligands modelled in this entry, 13 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PEG	A	17	-	6,6,6	0.46	0	5,5,5	1.33	0
2	PEG	A	2	-	6,6,6	0.48	0	5,5,5	1.23	0
3	TAM	A	32	-	10,10,10	0.62	0	12,12,12	1.23	3 (25%)
3	TAM	A	35	-	10,10,10	0.62	0	12,12,12	1.77	4 (33%)
2	PEG	B	13	-	6,6,6	0.47	0	5,5,5	1.32	0
2	PEG	C	14	-	6,6,6	0.46	0	5,5,5	1.25	0
3	TAM	C	28	-	10,10,10	0.61	0	12,12,12	1.68	3 (25%)
2	PEG	C	29	-	6,6,6	0.48	0	5,5,5	1.26	0
2	PEG	C	3	-	6,6,6	0.49	0	5,5,5	1.30	0
2	PEG	D	15	-	6,6,6	0.47	0	5,5,5	1.27	0
2	PEG	E	16	-	6,6,6	0.48	0	5,5,5	1.25	0
2	PEG	E	18	-	6,6,6	0.47	0	5,5,5	1.22	0
2	PEG	E	21	-	6,6,6	0.49	0	5,5,5	1.28	0
2	PEG	E	22	-	6,6,6	0.48	0	5,5,5	1.23	0
2	PEG	E	27	-	6,6,6	0.47	0	5,5,5	1.23	0
2	PEG	E	33	-	6,6,6	0.47	0	5,5,5	1.27	0
2	PEG	E	6	-	6,6,6	0.49	0	5,5,5	1.33	0
2	PEG	F	24	-	6,6,6	0.48	0	5,5,5	1.38	0
2	PEG	F	30	-	6,6,6	0.47	0	5,5,5	1.31	0
2	PEG	F	34	-	6,6,6	0.47	0	5,5,5	1.36	0
2	PEG	F	5	-	6,6,6	0.49	0	5,5,5	1.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	17	-	-	0/4/4/4	0/0/0/0
2	PEG	A	2	-	-	0/4/4/4	0/0/0/0
3	TAM	A	32	-	-	0/12/12/12	0/0/0/0
3	TAM	A	35	-	-	0/12/12/12	0/0/0/0
2	PEG	B	13	-	-	0/4/4/4	0/0/0/0
2	PEG	C	14	-	-	0/4/4/4	0/0/0/0
3	TAM	C	28	-	-	0/12/12/12	0/0/0/0
2	PEG	C	29	-	-	0/4/4/4	0/0/0/0
2	PEG	C	3	-	-	0/4/4/4	0/0/0/0
2	PEG	D	15	-	-	0/4/4/4	0/0/0/0
2	PEG	E	16	-	-	0/4/4/4	0/0/0/0
2	PEG	E	18	-	-	0/4/4/4	0/0/0/0
2	PEG	E	21	-	-	0/4/4/4	0/0/0/0
2	PEG	E	22	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	E	27	-	-	0/4/4/4	0/0/0/0
2	PEG	E	33	-	-	0/4/4/4	0/0/0/0
2	PEG	E	6	-	-	0/4/4/4	0/0/0/0
2	PEG	F	24	-	-	0/4/4/4	0/0/0/0
2	PEG	F	30	-	-	0/4/4/4	0/0/0/0
2	PEG	F	34	-	-	0/4/4/4	0/0/0/0
2	PEG	F	5	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	28	TAM	C5-C2-C	3.17	120.98	115.81
3	A	35	TAM	C6-C3-C	3.11	120.88	115.81
3	C	28	TAM	O5-C5-C2	2.82	119.86	111.33
3	A	35	TAM	O6-C6-C3	2.51	118.91	111.33
3	C	28	TAM	O6-C6-C3	2.42	118.63	111.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/275 (93%)	0.10	14 (5%)	24 20	24, 49, 81, 147	26 (10%)
1	B	255/275 (92%)	0.21	12 (4%)	30 27	25, 50, 95, 143	41 (16%)
1	C	254/275 (92%)	-0.00	7 (2%)	50 48	26, 47, 79, 123	14 (5%)
1	D	252/275 (91%)	0.25	15 (5%)	21 18	23, 55, 90, 116	44 (17%)
1	E	257/275 (93%)	0.07	13 (5%)	27 23	24, 53, 93, 131	44 (17%)
1	F	257/275 (93%)	0.11	6 (2%)	57 54	26, 52, 82, 127	34 (13%)
All	All	1531/1650 (92%)	0.12	67 (4%)	33 29	23, 51, 88, 147	203 (13%)

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1056	ALA	8.9
1	D	1061	LEU	7.4
1	A	1058	LEU	5.8
1	B	1055	GLU	5.4
1	A	1056	ALA	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	PEG	E	27	7/7	0.50	81.89	78,78,79,79	5
2	PEG	E	33	7/7	0.32	50.90	80,81,83,83	3
2	PEG	C	3	7/7	0.33	11.83	71,75,79,80	0
2	PEG	F	5	7/7	0.51	9.02	50,57,58,58	0
3	TAM	A	35	11/11	0.28	7.37	56,61,64,66	0
2	PEG	F	30	7/7	0.32	5.98	29,47,55,57	0
2	PEG	E	22	7/7	0.30	5.97	59,60,62,63	0
2	PEG	E	6	7/7	0.36	5.82	73,73,75,75	5
3	TAM	C	28	11/11	0.25	5.58	59,64,66,66	0
3	TAM	A	32	11/11	0.22	5.38	71,74,75,76	2
2	PEG	E	16	7/7	0.20	5.16	94,97,98,99	0
2	PEG	A	17	7/7	0.37	3.72	52,60,71,72	0
2	PEG	E	18	7/7	0.38	3.60	45,48,52,52	0
2	PEG	D	15	7/7	0.29	3.49	63,67,68,68	0
2	PEG	F	24	7/7	0.22	3.16	86,87,88,88	0
2	PEG	C	29	7/7	0.42	3.05	66,73,78,79	0
2	PEG	A	2	7/7	0.18	1.56	42,48,57,60	0
2	PEG	E	21	7/7	0.19	1.48	43,48,56,59	0
2	PEG	F	34	7/7	0.20	1.33	51,52,60,62	0
2	PEG	C	14	7/7	0.17	0.96	66,69,72,73	0
2	PEG	B	13	7/7	0.23	0.95	75,78,80,80	2
6	CA	F	1	1/1	0.19	0.79	44,44,44,44	0
4	CL	D	10	1/1	0.15	-0.60	42,42,42,42	0
4	CL	C	12	1/1	0.16	-1.00	66,66,66,66	0
4	CL	A	40	1/1	0.11	-1.22	44,44,44,44	0
4	CL	C	9	1/1	0.11	-1.91	41,41,41,41	0
5	BR	E	36	1/1	0.10	-2.37	61,61,61,61	0
5	BR	B	39	1/1	0.06	-2.64	77,77,77,77	1
5	BR	B	11	1/1	0.07	-2.86	78,78,78,78	0
4	CL	E	42	1/1	0.06	-3.17	61,61,61,61	0
5	BR	C	8	1/1	0.07	-3.33	64,64,64,64	0
4	CL	D	41	1/1	0.09	-3.79	45,45,45,45	0
5	BR	F	37	1/1	0.09	-4.63	68,68,68,68	0
5	BR	B	38	1/1	0.05	-8.34	66,66,66,66	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.