



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 02:44 PM GMT

PDB ID : 1H0A  
Title : EPSIN ENTH BOUND TO INS(1,4,5)P3  
Authors : Ford, M.G.J.; McMahon, H.T.; Evans, P.R.  
Deposited on : 2002-06-12  
Resolution : 1.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	158	

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 1611 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPSIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	158	Total	C	N	O	S	0	15	0
			1353	844	241	256	12			

- Molecule 2 is 1,4-DIETHYLENE DIOXIDE (three-letter code: DIO) (formula: C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	4	2		
2	A	1	Total	C	O	0	0
			6	4	2		
2	A	1	Total	C	O	0	0
			6	4	2		
2	A	1	Total	C	O	0	0
			6	4	2		
2	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 3 is D-MYO-INOSITOL-1,4,5-TRIPHOSPHATE (three-letter code: I3P) (formula:  $C_6H_{15}O_{15}P_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
3	A	1	24	6	15	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	204	Total	O	0	1
			204	204		

i

- Molecule 1: EPSIN

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.21Å 95.82Å 118.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.54 – 1.70 31.87 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (74.54-1.70) 97.4 (31.87-1.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.80 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, $R_{free}$	0.185 , 0.228 0.195 , 0.234	Depositor DCC
$R_{free}$ test set	2559 reflections (11.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtriage
Anisotropy	0.611	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 66.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 25488 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1611	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: DIO, I3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.38	11/1442 (0.8%)	1.19	6/1935 (0.3%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	62	LYS	CE-NZ	6.15	1.64	1.49
1	A	57	MET	SD-CE	-5.99	1.44	1.77
1	A	126	LYS	CD-CE	-5.81	1.36	1.51
1	A	139	ARG	CG-CD	-5.73	1.37	1.51
1	A	40	MET	CG-SD	5.72	1.96	1.81
1	A	14	VAL	CB-CG2	-5.66	1.41	1.52
1	A	10[A]	MET	SD-CE	-5.50	1.47	1.77
1	A	10[B]	MET	SD-CE	-5.50	1.47	1.77
1	A	8	ARG	CD-NE	-5.34	1.37	1.46
1	A	129	GLN	CG-CD	5.25	1.63	1.51
1	A	63	ARG	NE-CZ	-5.08	1.26	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	8	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	A	115	ASP	CB-CG-OD2	6.90	124.51	118.30
1	A	45	ASP	CB-CG-OD2	6.27	123.95	118.30
1	A	57	MET	CG-SD-CE	-5.99	90.61	100.20
1	A	136	ASP	CB-CG-OD1	5.92	123.63	118.30
1	A	7	ARG	NE-CZ-NH2	5.73	123.17	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1353	0	1332	22	0
2	A	30	0	40	2	0
3	A	24	0	9	0	0
4	A	204	0	0	12	2
All	All	1611	0	1381	23	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (23) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:94[A]:GLN:HG3	4:A:2126:HOH:O	1.63	0.97
1:A:136:ASP:OD2	4:A:2172:HOH:O	1.95	0.83
1:A:90:GLU:O	1:A:94[A]:GLN:HG2	1.90	0.72
1:A:147:ALA:HB3	2:A:1161:DIO:H1'2	1.80	0.64
1:A:86:LYS:NZ	4:A:2119:HOH:O	2.33	0.61
1:A:100:MET:HE1	1:A:135:ARG:HD3	1.82	0.61
1:A:35:PRO:CG	4:A:2113:HOH:O	2.49	0.60
1:A:35:PRO:HG3	4:A:2113:HOH:O	2.05	0.57
1:A:104[B]:GLN:OE1	1:A:131:VAL:HG11	2.07	0.54
1:A:62:LYS:NZ	4:A:2099:HOH:O	2.48	0.46
1:A:101[B]:TYR:HD2	4:A:2134:HOH:O	1.97	0.46
1:A:147:ALA:CB	2:A:1161:DIO:H1'2	2.48	0.42
1:A:151:LYS:HB3	1:A:151:LYS:HE3	1.79	0.42
1:A:101[B]:TYR:CD1	1:A:102:ALA:N	2.87	0.42
1:A:72:ARG:NH1	4:A:2111:HOH:O	2.50	0.42
1:A:1[B]:MET:HB2	4:A:2003:HOH:O	2.19	0.41
1:A:1[A]:MET:HG2	1:A:2:SER:N	2.35	0.41
1:A:65:ASN:O	1:A:67[A]:HIS:CD2	2.73	0.41
1:A:94[A]:GLN:NE2	4:A:2129:HOH:O	2.54	0.41
1:A:100:MET:HE3	1:A:100:MET:HB3	1.83	0.41
1:A:62:LYS:CE	4:A:2099:HOH:O	2.70	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:2018:HOH:O	4:A:2020:HOH:O[3_655]	1.82	0.38
4:A:2017:HOH:O	4:A:2020:HOH:O[3_655]	1.94	0.26

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/158 (108%)	168 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/141 (111%)	150 (96%)	6 (4%)	44	19

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	MET
1	A	1[B]	MET
1	A	37	SER
1	A	69	LYS
1	A	94[A]	GLN
1	A	94[B]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	65	ASN
1	A	95	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DIO	A	1159	-	6,6,6	0.66	0	6,6,6	0.65	0
2	DIO	A	1160	-	6,6,6	0.72	0	6,6,6	0.74	0
2	DIO	A	1161	-	6,6,6	0.78	0	6,6,6	0.67	0
2	DIO	A	1162	-	6,6,6	0.69	0	6,6,6	0.76	0
2	DIO	A	1163	-	6,6,6	0.72	0	6,6,6	1.20	1 (16%)
3	I3P	A	1164	-	24,24,24	1.46	3 (12%)	39,39,39	2.08	16 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DIO	A	1159	-	-	0/0/6/6	0/1/1/1
2	DIO	A	1160	-	-	0/0/6/6	0/1/1/1
2	DIO	A	1161	-	-	0/0/6/6	0/1/1/1
2	DIO	A	1162	-	-	0/0/6/6	0/1/1/1
2	DIO	A	1163	-	-	0/0/6/6	0/1/1/1
3	I3P	A	1164	-	-	0/15/39/39	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1164	I3P	P1-O13	-3.48	1.42	1.54
3	A	1164	I3P	P4-O43	-2.85	1.44	1.54
3	A	1164	I3P	P4-O41	2.11	1.58	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1164	I3P	P1-O1-C1	-4.40	112.70	121.96
3	A	1164	I3P	O52-P5-O51	-4.24	96.58	110.44
3	A	1164	I3P	C6-C1-C2	3.86	116.17	110.68
3	A	1164	I3P	O12-P1-O1	-3.51	96.98	107.09
3	A	1164	I3P	P4-O4-C4	3.43	129.19	121.96
3	A	1164	I3P	O43-P4-O42	3.05	119.50	107.61
3	A	1164	I3P	O6-C6-C1	2.72	116.19	109.85
3	A	1164	I3P	O6-C6-C5	2.65	116.02	109.85
3	A	1164	I3P	O53-P5-O52	2.49	117.31	107.61
3	A	1164	I3P	C6-C5-C4	2.47	116.77	111.39
3	A	1164	I3P	O3-C3-C4	2.41	115.47	109.85
3	A	1164	I3P	P5-O5-C5	2.30	126.81	121.96
3	A	1164	I3P	O43-P4-O4	-2.28	100.54	107.09
3	A	1164	I3P	O1-C1-C6	-2.26	103.83	108.44
3	A	1164	I3P	O2-C2-C1	2.24	115.08	109.85
3	A	1164	I3P	O1-C1-C2	2.12	112.75	108.44
2	A	1163	DIO	C2-O1-C1	2.07	116.98	109.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	158/158 (100%)	0.39	13 (8%)	12 14	19, 29, 54, 90	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	ALA	8.2
1	A	157	THR	6.6
1	A	156	GLN	5.1
1	A	154	LEU	4.9
1	A	101[A]	TYR	4.6
1	A	155	ALA	3.9
1	A	61[A]	TRP	3.3
1	A	114	ARG	2.9
1	A	1[A]	MET	2.9
1	A	2	SER	2.8
1	A	153	LYS	2.7
1	A	37	SER	2.5
1	A	34	GLY	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	DIO	A	1163	6/6	0.20	1.60	62,62,64,64	0
2	DIO	A	1161	6/6	0.13	1.51	52,53,56,56	0
2	DIO	A	1162	6/6	0.14	0.25	68,70,70,71	0
3	I3P	A	1164	24/24	0.09	-0.03	21,26,35,41	0
2	DIO	A	1159	6/6	0.11	-0.21	55,56,58,60	0
2	DIO	A	1160	6/6	0.10	-0.29	39,40,42,46	0

## 6.5 Other polymers

There are no such residues in this entry.