



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 06:11 AM GMT

PDB ID : 4H0M
Title : X-Ray Crystal Structure of Phycocyanin from *Synechococcus elongatus* sp.
PCC 7942
Authors : Marx, A.; Adir, N.
Deposited on : 2012-09-09
Resolution : 2.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2938 (2.20-2.20)
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	163	
1	C	163	
1	E	163	
1	G	163	
1	I	163	
1	K	163	
1	M	163	
1	O	163	
1	Q	163	
1	S	163	
1	U	163	
1	W	163	
2	B	173	
2	D	173	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	173	
2	H	173	
2	J	173	
2	L	173	
2	N	173	
2	P	173	
2	R	173	
2	T	173	
2	V	173	
2	X	173	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 32152 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-phycocyanin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	C	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	E	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	G	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	I	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	K	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	M	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	O	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	Q	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	S	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	U	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			
1	W	162	Total	C	N	O	S	0	0	0
			1213	766	204	242	1			

- Molecule 2 is a protein called C-phycocyanin beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	D	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	H	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	J	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	L	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	N	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	P	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	R	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	T	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	V	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			
2	X	172	Total	C	N	O	S	0	0	0
			1273	789	228	250	6			

- Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: $C_{33}H_{40}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			43	33	4	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total 43	C 33	N 4	O 6	0	0
3	B	1	Total 43	C 33	N 4	O 6	0	0
3	C	1	Total 43	C 33	N 4	O 6	0	0
3	D	1	Total 43	C 33	N 4	O 6	0	0
3	D	1	Total 43	C 33	N 4	O 6	0	0
3	E	1	Total 43	C 33	N 4	O 6	0	0
3	F	1	Total 43	C 33	N 4	O 6	0	0
3	F	1	Total 43	C 33	N 4	O 6	0	0
3	G	1	Total 43	C 33	N 4	O 6	0	0
3	H	1	Total 43	C 33	N 4	O 6	0	0
3	H	1	Total 43	C 33	N 4	O 6	0	0
3	I	1	Total 43	C 33	N 4	O 6	0	0
3	J	1	Total 43	C 33	N 4	O 6	0	0
3	J	1	Total 43	C 33	N 4	O 6	0	0
3	K	1	Total 43	C 33	N 4	O 6	0	0
3	L	1	Total 43	C 33	N 4	O 6	0	0
3	L	1	Total 43	C 33	N 4	O 6	0	0
3	M	1	Total 43	C 33	N 4	O 6	0	0
3	N	1	Total 43	C 33	N 4	O 6	0	0
3	N	1	Total 43	C 33	N 4	O 6	0	0
3	O	1	Total 43	C 33	N 4	O 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	P	1	Total	C	N	O	0	0
			43	33	4	6		
3	P	1	Total	C	N	O	0	0
			43	33	4	6		
3	Q	1	Total	C	N	O	0	0
			43	33	4	6		
3	R	1	Total	C	N	O	0	0
			43	33	4	6		
3	R	1	Total	C	N	O	0	0
			43	33	4	6		
3	S	1	Total	C	N	O	0	0
			43	33	4	6		
3	T	1	Total	C	N	O	0	0
			43	33	4	6		
3	T	1	Total	C	N	O	0	0
			43	33	4	6		
3	U	1	Total	C	N	O	0	0
			43	33	4	6		
3	V	1	Total	C	N	O	0	0
			43	33	4	6		
3	V	1	Total	C	N	O	0	0
			43	33	4	6		
3	W	1	Total	C	N	O	0	0
			43	33	4	6		
3	X	1	Total	C	N	O	0	0
			43	33	4	6		
3	X	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	41	Total	O	0	0
			41	41		
4	B	41	Total	O	0	0
			41	41		
4	C	31	Total	O	0	0
			31	31		
4	D	38	Total	O	0	0
			38	38		
4	E	46	Total	O	0	0
			46	46		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	19	Total 19	O 19	0	0
4	G	30	Total 30	O 30	0	0
4	H	17	Total 17	O 17	0	0
4	I	34	Total 34	O 34	0	0
4	J	25	Total 25	O 25	0	0
4	K	33	Total 33	O 33	0	0
4	L	20	Total 20	O 20	0	0
4	M	37	Total 37	O 37	0	0
4	N	23	Total 23	O 23	0	0
4	O	35	Total 35	O 35	0	0
4	P	27	Total 27	O 27	0	0
4	Q	38	Total 38	O 38	0	0
4	R	25	Total 25	O 25	0	0
4	S	47	Total 47	O 47	0	0
4	T	21	Total 21	O 21	0	0
4	U	41	Total 41	O 41	0	0
4	V	31	Total 31	O 31	0	0
4	W	40	Total 40	O 40	0	0
4	X	32	Total 32	O 32	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C-phycocyanin alpha chain

Chain A:



- Molecule 1: C-phycocyanin alpha chain

Chain C:




- Molecule 1: C-phycocyanin alpha chain

Chain E:



- Molecule 1: C-phycocyanin alpha chain

Chain G:




- Molecule 1: C-phycocyanin alpha chain

Chain I:




- Molecule 1: C-phycoerythrin alpha chain

Chain K:




- Molecule 1: C-phycoerythrin alpha chain

Chain M:



- Molecule 1: C-phycoerythrin alpha chain

Chain O:



- Molecule 1: C-phycoerythrin alpha chain

Chain Q:



- Molecule 1: C-phycoerythrin alpha chain


Chain S:






- Molecule 1: C-phycocyanin alpha chain

Chain U:




- Molecule 1: C-phycocyanin alpha chain

Chain W:



- Molecule 2: C-phycocyanin beta chain

Chain B:



- Molecule 2: C-phycocyanin beta chain

Chain D:




- Molecule 2: C-phycocyanin beta chain

Chain F:




- Molecule 2: C-phycocyanin beta chain

Chain H:




• Molecule 2: C-phycoerythrin beta chain

Chain J:




• Molecule 2: C-phycoerythrin beta chain

Chain L:




• Molecule 2: C-phycoerythrin beta chain

Chain N:



• Molecule 2: C-phycoerythrin beta chain


Chain P:



• Molecule 2: C-phycoerythrin beta chain

Chain R:





- Molecule 2: C-phycocyanin beta chain

Chain T:



- Molecule 2: C-phycocyanin beta chain

Chain V:



- Molecule 2: C-phycocyanin beta chain

Chain X:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.84Å 113.52Å 184.44Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	40.00 – 2.20 36.89 – 2.20	Depositor EDS
% Data completeness (in resolution range)	87.4 (40.00-2.20) 87.3 (36.89-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.242 , 0.295 0.244 , 0.298	Depositor DCC
R_{free} test set	9751 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	13.6	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 8.0	EDS
Estimated twinning fraction	0.447 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 195065 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	32152	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.07 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.2934e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	1/1237 (0.1%)	0.61	0/1681
1	C	0.79	0/1237	0.70	1/1681 (0.1%)
1	E	0.78	3/1237 (0.2%)	0.59	0/1681
1	G	0.68	0/1237	0.62	0/1681
1	I	0.81	3/1237 (0.2%)	0.61	0/1681
1	K	0.66	0/1237	0.62	1/1681 (0.1%)
1	M	0.64	1/1237 (0.1%)	0.63	1/1681 (0.1%)
1	O	0.68	1/1237 (0.1%)	0.62	1/1681 (0.1%)
1	Q	0.66	0/1237	0.66	1/1681 (0.1%)
1	S	0.84	0/1237	0.64	1/1681 (0.1%)
1	U	0.61	0/1237	0.57	0/1681
1	W	0.67	2/1237 (0.2%)	0.63	1/1681 (0.1%)
2	B	0.88	4/1276 (0.3%)	0.76	2/1724 (0.1%)
2	D	0.75	4/1276 (0.3%)	0.67	2/1724 (0.1%)
2	F	0.78	1/1276 (0.1%)	0.66	1/1724 (0.1%)
2	H	0.78	3/1276 (0.2%)	0.65	1/1724 (0.1%)
2	J	0.94	3/1276 (0.2%)	0.67	0/1724
2	L	0.86	2/1276 (0.2%)	0.71	0/1724
2	N	0.88	2/1276 (0.2%)	0.66	0/1724
2	P	0.87	4/1276 (0.3%)	0.75	2/1724 (0.1%)
2	R	0.91	2/1276 (0.2%)	0.74	3/1724 (0.2%)
2	T	0.82	4/1276 (0.3%)	0.74	3/1724 (0.2%)
2	V	0.72	2/1276 (0.2%)	0.66	0/1724
2	X	0.80	0/1276	0.73	1/1724 (0.1%)
All	All	0.78	42/30156 (0.1%)	0.67	22/40860 (0.1%)

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	153	CYS	CB-SG	-17.54	1.52	1.82
2	N	153	CYS	CB-SG	-16.53	1.54	1.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	153	CYS	CB-SG	-13.73	1.58	1.82
2	H	153	CYS	CB-SG	-11.14	1.63	1.82
2	L	153	CYS	CB-SG	-7.89	1.68	1.82

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	140	HIS	CB-CA-C	-11.58	87.23	110.40
2	P	74	TYR	C-N-CA	-7.38	103.24	121.70
2	T	153	CYS	CA-CB-SG	6.94	126.50	114.00
1	K	111	LEU	CA-CB-CG	6.75	130.82	115.30
2	P	74	TYR	O-C-N	-6.42	112.42	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1213	0	1182	39	0
1	C	1213	0	1183	48	0
1	E	1213	0	1183	31	0
1	G	1213	0	1183	41	0
1	I	1213	0	1183	33	0
1	K	1213	0	1183	35	0
1	M	1213	0	1183	46	0
1	O	1213	0	1183	36	0
1	Q	1213	0	1183	44	0
1	S	1213	0	1183	48	0
1	U	1213	0	1183	37	0
1	W	1213	0	1183	40	0
2	B	1273	0	1275	74	0
2	D	1273	0	1275	62	0
2	F	1273	0	1276	71	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1273	0	1275	62	0
2	J	1273	0	1276	65	0
2	L	1273	0	1275	56	0
2	N	1273	0	1275	100	0
2	P	1273	0	1275	65	0
2	R	1273	0	1274	60	0
2	T	1273	0	1274	56	0
2	V	1273	0	1276	59	0
2	X	1273	0	1275	47	0
3	A	43	0	37	5	0
3	B	86	0	71	19	0
3	C	43	0	37	6	0
3	D	86	0	74	18	0
3	E	43	0	37	4	0
3	F	86	0	74	13	0
3	G	43	0	37	5	0
3	H	86	0	73	8	0
3	I	43	0	37	7	0
3	J	86	0	76	16	0
3	K	43	0	37	5	0
3	L	86	0	74	13	0
3	M	43	0	37	2	0
3	N	86	0	74	11	0
3	O	43	0	37	4	0
3	P	86	0	73	16	0
3	Q	43	0	37	3	0
3	R	86	0	74	9	0
3	S	43	0	37	4	0
3	T	86	0	71	17	0
3	U	43	0	37	4	0
3	V	86	0	75	18	0
3	W	43	0	37	5	0
3	X	86	0	72	10	0
4	A	41	0	0	2	0
4	B	41	0	0	2	0
4	C	31	0	0	1	0
4	D	38	0	0	3	0
4	E	46	0	0	3	0
4	F	19	0	0	1	0
4	G	30	0	0	3	0
4	H	17	0	0	3	0
4	I	34	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	J	25	0	0	2	0
4	K	33	0	0	4	0
4	L	20	0	0	0	0
4	M	37	0	0	0	0
4	N	23	0	0	3	0
4	O	35	0	0	5	0
4	P	27	0	0	1	0
4	Q	38	0	0	4	0
4	R	25	0	0	2	0
4	S	47	0	0	2	0
4	T	21	0	0	2	0
4	U	41	0	0	1	0
4	V	31	0	0	0	0
4	W	40	0	0	4	0
4	X	32	0	0	1	0
All	All	32152	0	30821	1235	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 20.

The worst 5 of 1235 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:F:153:CYS:SG	3:F:202:CYC:HAC2	1.28	1.66
2:D:153:CYS:SG	3:D:202:CYC:HAC2	1.12	1.62
2:J:153:CYS:SG	3:J:202:CYC:HAC2	1.38	1.59
2:V:153:CYS:SG	3:V:202:CYC:HAC2	1.60	1.40
2:N:72:MEN:CE2	2:N:123:PRO:HD2	1.52	1.39

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	160/163 (98%)	154 (96%)	6 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	160/163 (98%)	154 (96%)	6 (4%)	0	100	100
1	E	160/163 (98%)	156 (98%)	4 (2%)	0	100	100
1	G	160/163 (98%)	154 (96%)	6 (4%)	0	100	100
1	I	160/163 (98%)	156 (98%)	4 (2%)	0	100	100
1	K	160/163 (98%)	153 (96%)	7 (4%)	0	100	100
1	M	160/163 (98%)	156 (98%)	4 (2%)	0	100	100
1	O	160/163 (98%)	155 (97%)	5 (3%)	0	100	100
1	Q	160/163 (98%)	153 (96%)	7 (4%)	0	100	100
1	S	160/163 (98%)	158 (99%)	2 (1%)	0	100	100
1	U	160/163 (98%)	153 (96%)	7 (4%)	0	100	100
1	W	160/163 (98%)	156 (98%)	4 (2%)	0	100	100
2	B	169/173 (98%)	166 (98%)	2 (1%)	1 (1%)	33	32
2	D	169/173 (98%)	161 (95%)	8 (5%)	0	100	100
2	F	169/173 (98%)	165 (98%)	2 (1%)	2 (1%)	19	14
2	H	169/173 (98%)	165 (98%)	4 (2%)	0	100	100
2	J	169/173 (98%)	166 (98%)	2 (1%)	1 (1%)	33	32
2	L	169/173 (98%)	166 (98%)	3 (2%)	0	100	100
2	N	169/173 (98%)	164 (97%)	5 (3%)	0	100	100
2	P	169/173 (98%)	160 (95%)	8 (5%)	1 (1%)	33	32
2	R	169/173 (98%)	165 (98%)	4 (2%)	0	100	100
2	T	169/173 (98%)	166 (98%)	2 (1%)	1 (1%)	33	32
2	V	169/173 (98%)	164 (97%)	2 (1%)	3 (2%)	13	7
2	X	169/173 (98%)	166 (98%)	3 (2%)	0	100	100
All	All	3948/4032 (98%)	3832 (97%)	107 (3%)	9 (0%)	56	62

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	74	TYR
2	V	103	SER
2	V	104	ILE
2	P	71	GLY
2	V	75	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/123 (99%)	121 (99%)	1 (1%)	89	95
1	C	122/123 (99%)	120 (98%)	2 (2%)	75	85
1	E	122/123 (99%)	120 (98%)	2 (2%)	75	85
1	G	122/123 (99%)	120 (98%)	2 (2%)	75	85
1	I	122/123 (99%)	121 (99%)	1 (1%)	89	95
1	K	122/123 (99%)	120 (98%)	2 (2%)	75	85
1	M	122/123 (99%)	119 (98%)	3 (2%)	60	71
1	O	122/123 (99%)	119 (98%)	3 (2%)	60	71
1	Q	122/123 (99%)	119 (98%)	3 (2%)	60	71
1	S	122/123 (99%)	119 (98%)	3 (2%)	60	71
1	U	122/123 (99%)	121 (99%)	1 (1%)	89	95
1	W	122/123 (99%)	122 (100%)	0	100	100
2	B	127/128 (99%)	123 (97%)	4 (3%)	52	63
2	D	127/128 (99%)	125 (98%)	2 (2%)	75	85
2	F	127/128 (99%)	125 (98%)	2 (2%)	75	85
2	H	127/128 (99%)	125 (98%)	2 (2%)	75	85
2	J	127/128 (99%)	124 (98%)	3 (2%)	61	73
2	L	127/128 (99%)	127 (100%)	0	100	100
2	N	127/128 (99%)	125 (98%)	2 (2%)	75	85
2	P	127/128 (99%)	124 (98%)	3 (2%)	61	73
2	R	127/128 (99%)	119 (94%)	8 (6%)	25	27
2	T	127/128 (99%)	125 (98%)	2 (2%)	75	85
2	V	127/128 (99%)	123 (97%)	4 (3%)	52	63
2	X	127/128 (99%)	124 (98%)	3 (2%)	61	73
All	All	2988/3012 (99%)	2930 (98%)	58 (2%)	69	81

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	74	TYR
2	P	67	ILE
2	V	111	ASN
2	N	75	THR
1	O	112	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	53	ASN
1	M	33	GLN
2	V	35	ASN
1	K	140	HIS
2	L	42	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MEN	B	72	2	8,8,9	3.81	3 (37%)	7,9,11	0.81	1 (14%)
2	MEN	D	72	2	8,8,9	3.81	2 (25%)	7,9,11	2.38	1 (14%)
2	MEN	F	72	2	8,8,9	2.05	2 (25%)	7,9,11	1.39	1 (14%)
2	MEN	H	72	2	8,8,9	3.58	2 (25%)	7,9,11	2.19	1 (14%)
2	MEN	J	72	2	8,8,9	1.64	2 (25%)	7,9,11	1.45	2 (28%)
2	MEN	L	72	2	8,8,9	6.28	2 (25%)	7,9,11	0.50	0
2	MEN	N	72	2	8,8,9	6.30	2 (25%)	7,9,11	0.69	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MEN	P	72	2	8,8,9	3.25	2 (25%)	7,9,11	0.93	1 (14%)
2	MEN	R	72	2	8,8,9	4.97	3 (37%)	7,9,11	1.95	2 (28%)
2	MEN	T	72	2	8,8,9	3.49	2 (25%)	7,9,11	1.09	0
2	MEN	V	72	2	8,8,9	3.76	5 (62%)	7,9,11	2.07	2 (28%)
2	MEN	X	72	2	8,8,9	5.14	2 (25%)	7,9,11	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	72	2	-	1/6/8/10	0/0/0/0
2	MEN	D	72	2	-	1/6/8/10	0/0/0/0
2	MEN	F	72	2	-	1/6/8/10	0/0/0/0
2	MEN	H	72	2	-	0/6/8/10	0/0/0/0
2	MEN	J	72	2	-	0/6/8/10	0/0/0/0
2	MEN	L	72	2	-	0/6/8/10	0/0/0/0
2	MEN	N	72	2	-	1/6/8/10	0/0/0/0
2	MEN	P	72	2	-	0/6/8/10	0/0/0/0
2	MEN	R	72	2	-	1/6/8/10	0/0/0/0
2	MEN	T	72	2	-	0/6/8/10	0/0/0/0
2	MEN	V	72	2	-	1/6/8/10	0/0/0/0
2	MEN	X	72	2	-	1/6/8/10	0/0/0/0

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	72	MEN	O-C	17.62	1.23	1.11
2	L	72	MEN	O-C	17.59	1.23	1.11
2	X	72	MEN	O-C	14.30	1.21	1.11
2	R	72	MEN	CE2-ND2	-10.51	1.27	1.45
2	D	72	MEN	O-C	10.32	1.18	1.11

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	72	MEN	C-CA-N	-6.18	107.66	113.83
2	H	72	MEN	C-CA-N	5.73	119.55	113.83
2	V	72	MEN	CB-CG-ND2	-4.28	109.32	115.87
2	R	72	MEN	OD1-CG-CB	-3.95	115.59	121.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	72	MEN	C-CA-N	3.47	117.29	113.83

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	72	MEN	CB-CG-ND2-CE2
2	X	72	MEN	CB-CG-ND2-CE2
2	D	72	MEN	CB-CG-ND2-CE2
2	B	72	MEN	CB-CG-ND2-CE2
2	N	72	MEN	CB-CG-ND2-CE2

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CYC	A	201	1	46,46,46	2.09	15 (32%)	65,67,67	3.83	28 (43%)
3	CYC	B	201	2	46,46,46	2.34	14 (30%)	65,67,67	3.71	25 (38%)
3	CYC	B	202	2	46,46,46	3.00	17 (36%)	65,67,67	4.12	29 (44%)
3	CYC	C	201	1	46,46,46	2.15	15 (32%)	65,67,67	3.73	25 (38%)
3	CYC	D	201	2	46,46,46	2.27	15 (32%)	65,67,67	3.82	25 (38%)
3	CYC	D	202	2	46,46,46	3.73	17 (36%)	65,67,67	4.46	30 (46%)
3	CYC	E	201	1	46,46,46	2.13	15 (32%)	65,67,67	3.75	24 (36%)
3	CYC	F	201	2	46,46,46	2.19	13 (28%)	65,67,67	3.72	21 (32%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CYC	F	202	2	46,46,46	2.55	17 (36%)	65,67,67	4.09	31 (47%)
3	CYC	G	201	1	46,46,46	2.25	17 (36%)	65,67,67	3.76	29 (44%)
3	CYC	H	201	2	46,46,46	2.30	17 (36%)	65,67,67	3.75	25 (38%)
3	CYC	H	202	2	46,46,46	2.16	15 (32%)	65,67,67	4.06	29 (44%)
3	CYC	I	201	1	46,46,46	2.23	16 (34%)	65,67,67	3.79	25 (38%)
3	CYC	J	201	2	46,46,46	2.25	16 (34%)	65,67,67	3.79	26 (40%)
3	CYC	J	202	2	46,46,46	2.51	15 (32%)	65,67,67	4.18	28 (43%)
3	CYC	K	201	1	46,46,46	2.12	15 (32%)	65,67,67	3.73	24 (36%)
3	CYC	L	201	-	46,46,46	2.42	15 (32%)	65,67,67	3.75	25 (38%)
3	CYC	L	202	2	46,46,46	2.46	15 (32%)	65,67,67	4.02	31 (47%)
3	CYC	M	201	1	46,46,46	2.25	13 (28%)	65,67,67	3.77	26 (40%)
3	CYC	N	201	2	46,46,46	2.27	14 (30%)	65,67,67	3.74	27 (41%)
3	CYC	N	202	2	46,46,46	2.92	18 (39%)	65,67,67	3.85	27 (41%)
3	CYC	O	201	1	46,46,46	2.14	16 (34%)	65,67,67	3.68	24 (36%)
3	CYC	P	201	2	46,46,46	2.20	16 (34%)	65,67,67	3.79	25 (38%)
3	CYC	P	202	2	46,46,46	2.69	15 (32%)	65,67,67	3.93	26 (40%)
3	CYC	Q	201	1	46,46,46	2.25	17 (36%)	65,67,67	3.67	28 (43%)
3	CYC	R	201	2	46,46,46	2.25	14 (30%)	65,67,67	3.68	25 (38%)
3	CYC	R	202	2	46,46,46	2.16	16 (34%)	65,67,67	3.81	29 (44%)
3	CYC	S	201	1	46,46,46	2.16	16 (34%)	65,67,67	3.75	26 (40%)
3	CYC	T	201	2	46,46,46	2.13	13 (28%)	65,67,67	3.70	24 (36%)
3	CYC	T	202	2	46,46,46	3.40	17 (36%)	65,67,67	4.21	33 (50%)
3	CYC	U	201	1	46,46,46	2.08	15 (32%)	65,67,67	3.83	24 (36%)
3	CYC	V	201	-	46,46,46	2.32	15 (32%)	65,67,67	3.74	25 (38%)
3	CYC	V	202	2	46,46,46	2.64	16 (34%)	65,67,67	3.98	28 (43%)
3	CYC	W	201	1	46,46,46	2.17	15 (32%)	65,67,67	3.81	25 (38%)
3	CYC	X	201	2	46,46,46	2.31	15 (32%)	65,67,67	3.79	23 (35%)
3	CYC	X	202	2	46,46,46	3.09	18 (39%)	65,67,67	4.40	29 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	A	201	1	-	3/25/74/74	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	B	201	2	-	3/25/74/74	0/4/4/4
3	CYC	B	202	2	-	3/25/74/74	0/4/4/4
3	CYC	C	201	1	-	3/25/74/74	0/4/4/4
3	CYC	D	201	2	-	3/25/74/74	0/4/4/4
3	CYC	D	202	2	-	3/25/74/74	0/4/4/4
3	CYC	E	201	1	-	3/25/74/74	0/4/4/4
3	CYC	F	201	2	-	3/25/74/74	0/4/4/4
3	CYC	F	202	2	-	3/25/74/74	0/4/4/4
3	CYC	G	201	1	-	3/25/74/74	0/4/4/4
3	CYC	H	201	2	-	3/25/74/74	0/4/4/4
3	CYC	H	202	2	-	3/25/74/74	0/4/4/4
3	CYC	I	201	1	-	3/25/74/74	0/4/4/4
3	CYC	J	201	2	-	3/25/74/74	0/4/4/4
3	CYC	J	202	2	-	3/25/74/74	0/4/4/4
3	CYC	K	201	1	-	3/25/74/74	0/4/4/4
3	CYC	L	201	-	-	3/25/74/74	0/4/4/4
3	CYC	L	202	2	-	3/25/74/74	0/4/4/4
3	CYC	M	201	1	-	3/25/74/74	0/4/4/4
3	CYC	N	201	2	-	3/25/74/74	0/4/4/4
3	CYC	N	202	2	-	3/25/74/74	0/4/4/4
3	CYC	O	201	1	-	3/25/74/74	0/4/4/4
3	CYC	P	201	2	-	3/25/74/74	0/4/4/4
3	CYC	P	202	2	-	3/25/74/74	0/4/4/4
3	CYC	Q	201	1	-	3/25/74/74	0/4/4/4
3	CYC	R	201	2	-	3/25/74/74	0/4/4/4
3	CYC	R	202	2	-	3/25/74/74	0/4/4/4
3	CYC	S	201	1	-	3/25/74/74	0/4/4/4
3	CYC	T	201	2	-	3/25/74/74	0/4/4/4
3	CYC	T	202	2	-	3/25/74/74	0/4/4/4
3	CYC	U	201	1	-	3/25/74/74	0/4/4/4
3	CYC	V	201	-	-	3/25/74/74	0/4/4/4
3	CYC	V	202	2	-	3/25/74/74	0/4/4/4
3	CYC	W	201	1	-	3/25/74/74	0/4/4/4
3	CYC	X	201	2	-	3/25/74/74	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	X	202	2	-	3/25/74/74	0/4/4/4

The worst 5 of 558 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	202	CYC	CBD-CGD	-18.72	1.02	1.50
3	T	202	CYC	CBD-CGD	-16.99	1.06	1.50
3	B	202	CYC	CBD-CGD	-12.85	1.17	1.50
3	X	202	CYC	CBD-CGD	-12.64	1.18	1.50
3	P	202	CYC	CBD-CGD	-11.28	1.21	1.50

The worst 5 of 954 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	U	201	CYC	C4B-C3B-C2B	-14.15	99.48	108.04
3	W	201	CYC	C4B-C3B-C2B	-13.94	99.61	108.04
3	F	201	CYC	C4B-C3B-C2B	-13.82	99.68	108.04
3	A	201	CYC	C4B-C3B-C2B	-13.71	99.75	108.04
3	E	201	CYC	C4B-C3B-C2B	-13.67	99.77	108.04

There are no chirality outliers.

5 of 108 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	T	202	CYC	C1B-CHB-C4A-NA
3	T	202	CYC	C1B-CHB-C4A-C3A
3	F	202	CYC	C1B-CHB-C4A-NA
3	X	202	CYC	C1B-CHB-C4A-NA
3	N	201	CYC	C4C-CHD-C1D-ND

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/163 (99%)	-0.30	0 100 100	3, 11, 25, 35	0
1	C	162/163 (99%)	-0.24	1 (0%) 86 88	5, 14, 27, 56	0
1	E	162/163 (99%)	-0.25	0 100 100	5, 12, 26, 57	0
1	G	162/163 (99%)	-0.22	0 100 100	5, 14, 29, 40	0
1	I	162/163 (99%)	-0.29	0 100 100	3, 12, 28, 55	0
1	K	162/163 (99%)	-0.23	0 100 100	3, 13, 27, 65	0
1	M	162/163 (99%)	-0.21	0 100 100	6, 13, 30, 51	0
1	O	162/163 (99%)	-0.27	0 100 100	4, 13, 25, 37	0
1	Q	162/163 (99%)	-0.19	1 (0%) 86 88	4, 13, 30, 65	0
1	S	162/163 (99%)	-0.26	0 100 100	3, 12, 23, 38	0
1	U	162/163 (99%)	-0.32	0 100 100	3, 12, 21, 30	0
1	W	162/163 (99%)	-0.19	0 100 100	4, 14, 28, 41	0
2	B	172/173 (99%)	-0.14	1 (0%) 86 88	3, 14, 31, 53	0
2	D	172/173 (99%)	-0.18	0 100 100	4, 15, 30, 55	0
2	F	172/173 (99%)	-0.07	2 (1%) 75 76	5, 16, 34, 48	0
2	H	172/173 (99%)	-0.14	0 100 100	7, 18, 36, 53	0
2	J	172/173 (99%)	-0.01	0 100 100	6, 16, 36, 50	0
2	L	172/173 (99%)	-0.08	2 (1%) 75 76	5, 16, 35, 51	0
2	N	172/173 (99%)	-0.12	3 (1%) 67 68	5, 15, 35, 54	0
2	P	172/173 (99%)	-0.11	2 (1%) 75 76	6, 15, 34, 61	0
2	R	172/173 (99%)	-0.08	1 (0%) 86 88	6, 17, 40, 53	0
2	T	172/173 (99%)	-0.14	1 (0%) 86 88	5, 16, 34, 51	0
2	V	172/173 (99%)	-0.19	0 100 100	5, 13, 29, 45	0
2	X	172/173 (99%)	-0.18	1 (0%) 86 88	5, 14, 33, 43	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	4008/4032 (99%)	-0.18	15 (0%) 90 92	3, 14, 33, 65	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	21	ASP	3.1
2	F	172	ALA	3.1
2	B	153	CYS	3.0
2	T	145	ARG	3.0
2	N	72	MEN	2.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MEN	L	72	9/10	0.22	4.01	20,20,23,30	0
2	MEN	N	72	9/10	0.26	2.45	15,19,24,24	0
2	MEN	X	72	9/10	0.17	2.22	19,21,23,23	0
2	MEN	J	72	9/10	0.21	2.08	15,19,27,29	0
2	MEN	V	72	9/10	0.17	1.66	20,21,27,27	0
2	MEN	T	72	9/10	0.17	1.41	20,20,30,30	0
2	MEN	P	72	9/10	0.17	1.22	20,20,31,34	0
2	MEN	H	72	9/10	0.18	1.05	20,20,32,32	0
2	MEN	R	72	9/10	0.15	0.94	15,19,28,28	0
2	MEN	F	72	9/10	0.20	0.92	26,28,35,37	0
2	MEN	B	72	9/10	0.14	0.54	15,19,20,20	0
2	MEN	D	72	9/10	0.12	-0.23	18,19,25,28	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CYC	R	202	43/43	0.19	1.80	9,20,32,39	0
3	CYC	A	201	43/43	0.15	1.43	5,11,18,23	0
3	CYC	X	202	43/43	0.17	1.31	6,18,29,40	0
3	CYC	P	202	43/43	0.15	1.25	5,16,33,41	0
3	CYC	T	201	43/43	0.16	0.99	15,19,26,29	0
3	CYC	F	202	43/43	0.16	0.96	4,21,35,43	0
3	CYC	V	202	43/43	0.16	0.88	7,17,32,41	0
3	CYC	L	201	43/43	0.15	0.81	6,16,25,31	0
3	CYC	M	201	43/43	0.15	0.77	7,12,17,23	0
3	CYC	B	202	43/43	0.16	0.68	5,15,31,40	0
3	CYC	H	202	43/43	0.15	0.65	4,15,32,41	0
3	CYC	D	202	43/43	0.15	0.62	4,17,31,39	0
3	CYC	L	202	43/43	0.15	0.60	2,18,35,46	0
3	CYC	U	201	43/43	0.13	0.50	6,9,14,17	0
3	CYC	S	201	43/43	0.13	0.40	3,7,12,16	0
3	CYC	P	201	43/43	0.14	0.40	8,17,25,26	0
3	CYC	T	202	43/43	0.16	0.39	7,18,32,39	0
3	CYC	H	201	43/43	0.14	0.34	14,18,22,24	0
3	CYC	V	201	43/43	0.13	0.33	3,11,20,23	0
3	CYC	N	202	43/43	0.15	0.23	4,17,31,38	0
3	CYC	F	201	43/43	0.13	0.23	13,18,22,25	0
3	CYC	J	201	43/43	0.14	0.22	6,19,24,27	0
3	CYC	R	201	43/43	0.14	0.18	16,18,24,25	0
3	CYC	O	201	43/43	0.13	0.12	5,11,14,20	0
3	CYC	E	201	43/43	0.13	0.11	3,9,13,15	0
3	CYC	I	201	43/43	0.12	0.06	5,10,13,15	0
3	CYC	W	201	43/43	0.12	-0.03	7,12,16,18	0
3	CYC	D	201	43/43	0.13	-0.05	11,14,19,22	0
3	CYC	B	201	43/43	0.13	-0.05	8,14,25,27	0
3	CYC	K	201	43/43	0.12	-0.06	2,10,16,20	0
3	CYC	N	201	43/43	0.13	-0.08	8,17,28,32	0
3	CYC	X	201	43/43	0.12	-0.16	8,15,18,22	0
3	CYC	J	202	43/43	0.14	-0.17	4,13,34,45	0
3	CYC	C	201	43/43	0.12	-0.31	5,10,15,21	0
3	CYC	G	201	43/43	0.11	-0.40	5,15,18,18	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CYC	Q	201	43/43	0.11	-0.58	5,10,13,14	0

6.5 Other polymers ⓘ

There are no such residues in this entry.