



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 01:37 PM GMT

PDB ID : 2H2S
Title : Crystal Structure of E148A mutant of CLC-ec1 in SeCN-
Authors : Nguitragool, W.; Miller, C.
Deposited on : 2006-05-19
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	FAILED
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1007 (3.18-3.02)
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)
RSRZ outliers	66119	1008 (3.18-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	465	
1	B	465	
2	C	221	
2	E	221	
3	D	211	
3	F	211	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13221 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLC Cl transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3329	2188	560	561	20			
1	B	441	Total	C	N	O	S	0	0	0
			3300	2172	553	555	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	148	ALA	GLU	ENGINEERED	UNP P37019
B	148	ALA	GLU	ENGINEERED	UNP P37019

- Molecule 2 is a protein called FAB fragment, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	221	Total	C	N	O	S	0	0	0
			1672	1077	274	315	6			
2	E	221	Total	C	N	O	S	0	0	0
			1672	1077	274	315	6			

- Molecule 3 is a protein called FAB fragment, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	211	Total	C	N	O	S	0	0	0
			1621	1008	271	334	8			
3	F	211	Total	C	N	O	S	0	0	0
			1621	1008	271	334	8			

- Molecule 4 is SELENOCYANATE ION (three-letter code: SEK) (formula: CNSe).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Se	0	0
			1	1		
4	B	1	Total	Se	0	0
			1	1		
4	A	1	Total	Se	1	0
			1	1		
4	B	1	Total	Se	1	0
			1	1		
4	A	1	Total	Se	1	0
			1	1		
4	B	1	Total	Se	1	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CLC Cl transporter


Chain A:



• Molecule 1: CLC Cl transporter


Chain B:






• Molecule 2: FAB fragment, heavy chain

Chain C:




• Molecule 2: FAB fragment, heavy chain


Chain E:



• Molecule 3: FAB fragment, light chain

Chain D:





- Molecule 3: FAB fragment, light chain

Chain F:



4 Data and refinement statistics

Xtriage (Phenix) failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	216.93Å 118.69Å 148.91Å 90.00° 127.44° 90.00°	Depositor
Resolution (Å)	50.00 – 3.10 48.87 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-3.10) 99.3 (48.87-3.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.281 , 0.282 0.278 , 0.280	Depositor DCC
R_{free} test set	2714 reflections (5.28%)	DCC
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 66.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	13221	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SEK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3401	0.67	0/4616
1	B	0.47	0/3372	0.67	0/4578
2	C	0.53	0/1721	0.79	0/2355
2	E	0.50	0/1721	0.74	0/2355
3	D	0.56	1/1660 (0.1%)	0.80	1/2257 (0.0%)
3	F	0.53	0/1660	0.76	1/2257 (0.0%)
All	All	0.51	1/13535 (0.0%)	0.72	2/18418 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	167	SER	N-CA	-5.62	1.35	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	167	SER	N-CA-CB	6.00	119.50	110.50
3	F	169	ASP	N-CA-C	5.92	126.98	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3329	0	3483	400	0
1	B	3300	0	3456	386	0
2	C	1672	0	1653	190	0
2	E	1672	0	1654	163	0
3	D	1621	0	1546	263	0
3	F	1621	0	1546	203	0
4	A	3	0	0	5	0
4	B	3	0	0	5	0
All	All	13221	0	13338	1503	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 57.

The worst 5 of 1503 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:131:PRO:CG	2:E:216:LYS:HE3	1.29	1.56
2:E:131:PRO:CD	2:E:216:LYS:HE2	1.27	1.55
2:C:53:PRO:CA	2:C:72:ARG:NH2	1.68	1.53
2:E:131:PRO:CG	2:E:216:LYS:CE	1.82	1.50
3:D:88:GLN:HG3	3:D:97:PHE:CE1	1.60	1.33

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/465 (95%)	318 (72%)	101 (23%)	23 (5%)	3	21
1	B	439/465 (94%)	313 (71%)	101 (23%)	25 (6%)	3	18
2	C	219/221 (99%)	170 (78%)	29 (13%)	20 (9%)	1	8
2	E	219/221 (99%)	173 (79%)	35 (16%)	11 (5%)	3	22
3	D	209/211 (99%)	151 (72%)	33 (16%)	25 (12%)	1	4
3	F	209/211 (99%)	166 (79%)	30 (14%)	13 (6%)	2	16
All	All	1737/1794 (97%)	1291 (74%)	329 (19%)	117 (7%)	2	14

5 of 117 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	PHE
1	A	107	SER
1	A	132	PHE
1	A	148	ALA
1	A	167	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/352 (95%)	315 (94%)	19 (6%)	29	70
1	B	331/352 (94%)	312 (94%)	19 (6%)	29	70
2	C	181/181 (100%)	163 (90%)	18 (10%)	11	39
2	E	181/181 (100%)	165 (91%)	16 (9%)	14	49
3	D	185/185 (100%)	166 (90%)	19 (10%)	10	36
3	F	185/185 (100%)	167 (90%)	18 (10%)	12	41
All	All	1397/1436 (97%)	1288 (92%)	109 (8%)	18	57

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	172	HIS
3	D	70	TYR
3	F	119	PRO
2	C	182	LEU
2	C	204	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	207	GLN
1	B	327	ASN
3	F	36	GLN
1	B	270	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	277	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/465 (95%)	0.11	2 (0%) 88 39	78, 78, 78, 78	0
1	B	441/465 (94%)	0.16	2 (0%) 88 39	78, 78, 78, 78	0
2	C	221/221 (100%)	0.07	2 (0%) 81 25	78, 78, 78, 78	0
2	E	221/221 (100%)	-0.01	0 100 100	78, 78, 78, 78	0
3	D	211/211 (100%)	0.11	2 (0%) 81 25	78, 78, 78, 78	0
3	F	211/211 (100%)	0.07	1 (0%) 88 39	78, 78, 78, 78	0
All	All	1749/1794 (97%)	0.10	9 (0%) 88 39	78, 78, 78, 78	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	168	LEU	2.4
1	B	75	TYR	2.3
3	F	182	LYS	2.3
1	A	282	ARG	2.2
1	B	64	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SEK	A	468	1/3	-	-	78,78,78,78	1
4	SEK	A	466	1/3	0.29	-	78,78,78,78	0
4	SEK	B	466	1/3	0.46	-	78,78,78,78	0
4	SEK	A	467	1/3	-	-	78,78,78,78	1
4	SEK	B	467	1/3	-	-	78,78,78,78	1
4	SEK	B	468	1/3	-	-	78,78,78,78	1

6.5 Other polymers

There are no such residues in this entry.