



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 02:38 PM GMT

PDB ID : 1H3Y  
Title : CRYSTAL STRUCTURE OF A HUMAN IGG1 FC-FRAGMENT,HIGH  
SALT CONDITION  
Authors : Krapp, S.; Mimura, Y.; Jefferis, R.; Huber, R.; Sondermann, P.  
Deposited on : 2002-09-19  
Resolution : 4.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1002 (4.70-3.48)
Clashscore	79885	1248 (4.70-3.50)
Ramachandran outliers	78287	1183 (4.70-3.50)
Sidechain outliers	78261	1168 (4.70-3.50)
RSRZ outliers	66119	1002 (4.70-3.48)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	223	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
1	B	223	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3015 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IG GAMMA-1 CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	1
			1373	848	245	274	6			
1	B	206	Total	C	N	O	S	0	0	0
			1422	886	246	283	7			

- Molecule 2 is a polymer of unknown type called SUGAR (9-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	9	Total	C	N	O	0	0
			110	62	4	44		

- Molecule 3 is a polymer of unknown type called SUGAR (9-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	9	Total	C	N	O	0	0
			110	62	4	44		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	152.94Å 152.94Å 116.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 4.10 38.24 – 4.13	Depositor EDS
% Data completeness (in resolution range)	80.0 (50.00-4.10) 79.2 (38.24-4.13)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 4.13Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.285 , 0.349 0.278 , 0.340	Depositor DCC
$R_{free}$ test set	285 reflections (5.78%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.4	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 112.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 5218 reflections	Xtriage
$F_o, F_c$ correlation	0.81	EDS
Total number of atoms	3015	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, FUL, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/1415	0.83	1/1927 (0.1%)
1	B	0.53	0/1466	0.87	4/1988 (0.2%)
All	All	0.52	0/2881	0.85	5/3915 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	1	0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	402	GLY	N-CA-C	-6.03	98.02	113.10
1	A	402	GLY	N-CA-C	-5.78	98.65	113.10
1	B	443	LEU	CA-CB-CG	-5.73	102.13	115.30
1	B	297	ASN	N-CA-CB	-5.18	101.27	110.60
1	B	395	PRO	N-CA-C	-5.11	98.81	112.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1449	NAG	C1

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1373	0	997	88	0
1	B	1422	0	1100	114	1
2	A	110	0	94	22	0
3	B	110	0	94	13	0
All	All	3015	0	2285	221	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 42.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:1452:NAG:H83	3:B:1445:FUL:H4	1.25	1.18
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.24	1.10
1:B:325:ASN:ND2	1:B:326:LYS:H	1.52	1.06
1:A:283:GLN:HG2	1:A:285:HIS:H	1.30	0.96
3:B:1451:MAN:H2	3:B:1452:NAG:N2	1.78	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:309:LEU:CD1	1:B:309:LEU:CD1[9_765]	1.59	0.61

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/223 (92%)	165 (80%)	22 (11%)	18 (9%)	1	25
1	B	204/223 (92%)	159 (78%)	29 (14%)	16 (8%)	1	28
All	All	409/446 (92%)	324 (79%)	51 (12%)	34 (8%)	1	27

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	ALA
1	A	377	ILE
1	A	443	LEU
1	B	297	ASN
1	B	377	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/206 (52%)	101 (94%)	7 (6%)	24	72
1	B	127/206 (62%)	116 (91%)	11 (9%)	15	59
All	All	235/412 (57%)	217 (92%)	18 (8%)	18	65

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	271	PRO
1	B	325	ASN
1	B	411	THR
1	A	443	LEU
1	B	260	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
1	A	434	ASN
1	B	325	ASN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	325	ASN
1	B	312	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	1444	1,2	12,14,15	1.01	1 (8%)	15,19,21	1.53	4 (26%)
2	FUL	A	1445	2	9,10,11	0.61	0	10,14,16	0.65	0
2	NAG	A	1446	2	12,14,15	0.96	1 (8%)	15,19,21	1.42	3 (20%)
2	MAN	A	1447	2	10,11,12	0.67	0	11,15,17	1.41	1 (9%)
2	MAN	A	1448	2	10,11,12	0.81	0	11,15,17	0.75	0
2	NAG	A	1449	2	12,14,15	0.66	0	15,19,21	1.56	3 (20%)
2	GLA	A	1450	2	10,11,12	0.74	0	11,15,17	0.88	1 (9%)
2	MAN	A	1451	2	10,11,12	0.64	0	11,15,17	0.70	0
2	NAG	A	1452	2	12,14,15	0.67	0	15,19,21	1.03	1 (6%)
3	NAG	B	1444	1,3	12,14,15	0.74	0	15,19,21	1.21	2 (13%)
3	FUL	B	1445	3	9,10,11	0.72	0	10,14,16	0.64	0
3	NAG	B	1446	3	12,14,15	0.77	0	15,19,21	1.10	1 (6%)
3	BMA	B	1447	3	10,11,12	0.80	1 (10%)	11,15,17	0.86	0
3	MAN	B	1448	3	10,11,12	0.64	0	11,15,17	1.20	1 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	1449	3	12,14,15	0.83	0	15,19,21	1.24	2 (13%)
3	GLA	B	1450	3	10,11,12	0.46	0	11,15,17	0.74	0
3	MAN	B	1451	3	10,11,12	0.87	0	11,15,17	1.04	1 (9%)
3	NAG	B	1452	3	12,14,15	0.66	0	15,19,21	1.14	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1444	1,2	-	1/6/23/26	0/1/1/1
2	FUL	A	1445	2	-	0/0/17/20	1/1/1/1
2	NAG	A	1446	2	-	0/6/23/26	1/1/1/1
2	MAN	A	1447	2	-	0/2/19/22	0/1/1/1
2	MAN	A	1448	2	-	0/2/19/22	0/1/1/1
2	NAG	A	1449	2	1/1/5/7	0/6/23/26	0/1/1/1
2	GLA	A	1450	2	-	0/2/19/22	0/1/1/1
2	MAN	A	1451	2	-	0/2/19/22	0/1/1/1
2	NAG	A	1452	2	-	0/6/23/26	0/1/1/1
3	NAG	B	1444	1,3	-	0/6/23/26	0/1/1/1
3	FUL	B	1445	3	-	0/0/17/20	1/1/1/1
3	NAG	B	1446	3	-	0/6/23/26	0/1/1/1
3	BMA	B	1447	3	-	0/2/19/22	0/1/1/1
3	MAN	B	1448	3	-	0/2/19/22	0/1/1/1
3	NAG	B	1449	3	-	0/6/23/26	0/1/1/1
3	GLA	B	1450	3	-	0/2/19/22	1/1/1/1
3	MAN	B	1451	3	-	0/2/19/22	0/1/1/1
3	NAG	B	1452	3	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1444	NAG	C4-C5	2.43	1.58	1.53
2	A	1446	NAG	O4-C4	2.15	1.48	1.43
3	B	1447	BMA	C3-C2	2.03	1.57	1.52

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1447	MAN	C4-C3-C2	3.59	115.32	110.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1444	NAG	C2-N2-C7	-3.47	117.27	123.09
2	A	1449	NAG	C4-C3-C2	-3.45	102.86	111.32
2	A	1444	NAG	C3-C4-C5	2.93	115.44	110.20
2	A	1449	NAG	C3-C4-C5	-2.87	105.08	110.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1449	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1444	NAG	O7-C7-N2-C2

All (4) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1445	FUL	C1-C2-C3-C4-C5-O5
2	A	1445	FUL	C1-C2-C3-C4-C5-O5
2	A	1446	NAG	C1-C2-C3-C4-C5-O5
3	B	1450	GLA	C1-C2-C3-C4-C5-O5

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/223 (92%)	-0.05	0 100 100	28, 32, 47, 47	0
1	B	206/223 (92%)	-0.27	0 100 100	14, 17, 22, 23	0
All	All	413/446 (92%)	-0.16	0 100 100	14, 23, 46, 47	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GLA	B	1450	11/12	0.24	1.20	33,34,38,39	0
2	GLA	A	1450	11/12	0.27	0.54	71,72,76,76	0
2	NAG	A	1444	14/15	0.26	-0.36	71,73,76,76	0
2	NAG	A	1449	14/15	0.26	-0.59	71,74,76,76	0
3	NAG	B	1444	14/15	0.18	-0.68	33,35,39,39	0
3	NAG	B	1449	14/15	0.18	-0.77	33,35,38,38	0
2	FUL	A	1445	10/11	0.19	-1.64	70,73,75,76	0
3	NAG	B	1446	14/15	0.16	-1.67	33,35,38,39	0
3	MAN	B	1448	11/12	0.23	-1.95	33,35,38,38	0
2	MAN	A	1448	11/12	0.21	-2.28	71,73,76,76	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	1446	14/15	0.24	-3.20	71,74,75,76	0
3	BMA	B	1447	11/12	0.12	-3.39	33,36,38,39	0
2	NAG	A	1452	14/15	0.25	-	71,73,75,76	0
2	MAN	A	1447	11/12	0.19	-	71,73,76,76	0
3	FUL	B	1445	10/11	0.18	-	33,36,38,39	0
2	MAN	A	1451	11/12	0.23	-	73,75,76,76	0
3	MAN	B	1451	11/12	0.28	-	33,35,38,38	0
3	NAG	B	1452	14/15	0.25	-	33,36,39,39	0

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.