



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 03:41 AM GMT

PDB ID : 2H4Q
Title : Crystal structure of a M-loop deletion variant of MENT in the cleaved conformation
Authors : Whisstock, J.C.; Buckle, A.M.; McGowan, S.; Irving, J.A.
Deposited on : 2006-05-25
Resolution : 2.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	382	
2	B	34	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3443 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heterochromatin-associated protein MENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	352	2814	1800	463	537	14	0	0	0

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-40	MET	-	CLONING ARTIFACT	UNP O73790
A	-39	GLY	-	CLONING ARTIFACT	UNP O73790
A	-38	GLY	-	CLONING ARTIFACT	UNP O73790
A	-37	SER	-	CLONING ARTIFACT	UNP O73790
A	-36	HIS	-	CLONING ARTIFACT	UNP O73790
A	-35	HIS	-	CLONING ARTIFACT	UNP O73790
A	-34	HIS	-	CLONING ARTIFACT	UNP O73790
A	-33	HIS	-	CLONING ARTIFACT	UNP O73790
A	-32	HIS	-	CLONING ARTIFACT	UNP O73790
A	-31	HIS	-	CLONING ARTIFACT	UNP O73790
A	-30	GLY	-	CLONING ARTIFACT	UNP O73790
A	-29	MET	-	CLONING ARTIFACT	UNP O73790
A	-28	ALA	-	CLONING ARTIFACT	UNP O73790
A	-27	SER	-	CLONING ARTIFACT	UNP O73790
A	-26	MET	-	CLONING ARTIFACT	UNP O73790
A	-25	THR	-	CLONING ARTIFACT	UNP O73790
A	-24	GLY	-	CLONING ARTIFACT	UNP O73790
A	-23	GLY	-	CLONING ARTIFACT	UNP O73790
A	-22	GLN	-	CLONING ARTIFACT	UNP O73790
A	-21	GLN	-	CLONING ARTIFACT	UNP O73790
A	-20	MET	-	CLONING ARTIFACT	UNP O73790
A	-19	GLY	-	CLONING ARTIFACT	UNP O73790
A	-18	ARG	-	CLONING ARTIFACT	UNP O73790
A	-17	ASP	-	CLONING ARTIFACT	UNP O73790
A	-16	LEU	-	CLONING ARTIFACT	UNP O73790
A	-15	TYR	-	CLONING ARTIFACT	UNP O73790
A	-14	ASP	-	CLONING ARTIFACT	UNP O73790

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	ASP	-	CLONING ARTIFACT	UNP O73790
A	-12	ASP	-	CLONING ARTIFACT	UNP O73790
A	-11	ASP	-	CLONING ARTIFACT	UNP O73790
A	-10	LYS	-	CLONING ARTIFACT	UNP O73790
A	-9	ASP	-	CLONING ARTIFACT	UNP O73790
A	-8	ARG	-	CLONING ARTIFACT	UNP O73790
A	-7	TRP	-	CLONING ARTIFACT	UNP O73790
A	-6	GLY	-	CLONING ARTIFACT	UNP O73790
A	-5	SER	-	CLONING ARTIFACT	UNP O73790
A	-4	GLU	-	CLONING ARTIFACT	UNP O73790
A	-3	LEU	-	CLONING ARTIFACT	UNP O73790
A	-2	GLU	-	CLONING ARTIFACT	UNP O73790
A	-1	ILE	-	CLONING ARTIFACT	UNP O73790
A	0	SER	-	CLONING ARTIFACT	UNP O73790
A	?	-	THR	DELETION	UNP O73790
A	?	-	GLU	DELETION	UNP O73790
A	?	-	ALA	DELETION	UNP O73790
A	?	-	VAL	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	ALA	DELETION	UNP O73790
A	?	-	GLU	DELETION	UNP O73790
A	?	-	SER	DELETION	UNP O73790
A	?	-	SER	DELETION	UNP O73790
A	?	-	SER	DELETION	UNP O73790
A	?	-	VAL	DELETION	UNP O73790
A	?	-	ALA	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	PRO	DELETION	UNP O73790
A	?	-	SER	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	GLY	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	PRO	DELETION	UNP O73790
A	?	-	LYS	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	ARG	DELETION	UNP O73790
A	?	-	MET	DELETION	UNP O73790
A	?	-	ASP	DELETION	UNP O73790
A	?	-	PRO	DELETION	UNP O73790
A	?	-	GLU	DELETION	UNP O73790
A	?	-	HIS	DELETION	UNP O73790

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	361	VAL	ALA	ENGINEERED	UNP O73790

- Molecule 2 is a protein called Heterochromatin-associated protein MENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	33	Total	C	N	O	S	0	0	0
			279	190	49	38	2			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	323	Total	O	0	0
			323	323		
3	B	27	Total	O	0	0
			27	27		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heterochromatin-associatedprotein MENT

Chain A: 



- Molecule 2: Heterochromatin-associatedprotein MENT

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	150.19Å 150.19Å 175.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.35 – 2.10 24.34 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (24.35-2.10) 99.4 (24.34-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.188 , 0.213 0.208 , 0.236	Depositor DCC
R_{free} test set	1782 reflections (4.21%)	DCC
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.2	EDS
Estimated twinning fraction	0.021 for -2/3*h-1/3*k+2/3*l,-1/3*h-2/3*k-2/3*l,2/3*h-2/3*k+1/3*l 0.012 for -h,1/3*h-1/3*k+2/3*l,2/3*h+4/3*k+1/3*l 0.011 for -1/3*h+1/3*k-2/3*l,-k,-4/3*h-2/3*k+1/3*l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 44120 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3443	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	1/2866 (0.0%)	0.46	0/3864
2	B	0.42	0/290	0.45	0/389
All	All	0.36	1/3156 (0.0%)	0.46	0/4253

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	GLU	CB-CG	-5.15	1.42	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2814	0	2780	10	0
2	B	279	0	274	1	0
3	A	323	0	0	3	2
3	B	27	0	0	0	0
All	All	3443	0	3054	10	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

The worst 5 of 10 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:0:SER:OG	3:A:643:HOH:O	2.07	0.73
1:A:244:MET:HE2	1:A:286:TRP:HB3	1.82	0.61
1:A:244:MET:CE	1:A:286:TRP:HB3	2.36	0.54
1:A:244:MET:HE1	1:A:286:TRP:C	2.35	0.47
1:A:56:GLU:CG	3:A:469:HOH:O	2.62	0.46

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:642:HOH:O	3:A:657:HOH:O[4_555]	2.06	0.14
3:A:644:HOH:O	3:A:657:HOH:O[4_555]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/382 (92%)	342 (98%)	8 (2%)	0	100	100
2	B	31/34 (91%)	30 (97%)	1 (3%)	0	100	100
All	All	381/416 (92%)	372 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/340 (89%)	302 (99%)	2 (1%)	91	94

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	30/33 (91%)	30 (100%)	0	100	100
All	All	334/373 (90%)	332 (99%)	2 (1%)	92	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-10	LYS
1	A	-2	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	27	ASN
1	A	241	ASN
1	A	318	ASN
1	A	347	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/382 (92%)	-0.00	9 (2%) 53 58	22, 35, 53, 59	0
2	B	33/34 (97%)	-0.01	1 (3%) 48 53	23, 28, 43, 50	0
All	All	385/416 (92%)	-0.00	10 (2%) 53 58	22, 35, 53, 59	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	377	HIS	3.9
1	A	328	ASN	3.5
1	A	173	LYS	2.8
1	A	327	THR	2.7
1	A	188	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.