



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 06:23 AM GMT

PDB ID : 1HF8
Title : CALM-N N-TERMINAL DOMAIN OF CLATHRIN ASSEMBLY LYMPHOID MYELOID LEUKAEMIA PROTEIN
Authors : Ford, M.G.J.; Evans, P.R.; McMahon, H.T.
Deposited on : 2000-11-30
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	289	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2244 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLATHRIN ASSEMBLY PROTEIN SHORT FORM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2114	1345	363	393	13			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	130	Total	O	0	0
			130	130		

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CLATHRIN ASSEMBLY PROTEIN SHORT FORM

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	77.93Å 77.93Å 121.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.94 – 2.00 36.02 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (65.94-2.00) 100.0 (36.02-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.190 , 0.223 0.201 , 0.237	Depositor DCC
R_{free} test set	1324 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	35.2	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 26057 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2244	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.46	15/2150 (0.7%)	1.31	21/2896 (0.7%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	MET	SD-CE	-13.32	1.03	1.77
1	A	211	TYR	CD1-CE1	9.01	1.52	1.39
1	A	206	ARG	CG-CD	-7.99	1.31	1.51
1	A	257	VAL	CB-CG2	-7.46	1.37	1.52
1	A	96	TYR	CD2-CE2	-6.80	1.29	1.39
1	A	96	TYR	CD1-CE1	-6.27	1.29	1.39
1	A	255	GLU	CD-OE2	-5.79	1.19	1.25
1	A	92	ARG	CG-CD	5.78	1.66	1.51
1	A	121	SER	CA-CB	5.77	1.61	1.52
1	A	264	ILE	CA-CB	-5.48	1.42	1.54
1	A	35	MET	SD-CE	-5.37	1.47	1.77
1	A	52	MET	SD-CE	5.29	2.07	1.77
1	A	136	VAL	CB-CG2	-5.29	1.41	1.52
1	A	257	VAL	C-O	-5.10	1.13	1.23
1	A	88	TYR	CB-CG	-5.04	1.44	1.51

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	ARG	CA-CB-CG	-7.25	97.44	113.40
1	A	261	ARG	NE-CZ-NH1	7.25	123.92	120.30
1	A	211	TYR	CB-CG-CD2	-6.81	116.92	121.00
1	A	268	SER	N-CA-C	-6.59	93.21	111.00
1	A	181	PHE	CB-CG-CD2	-6.42	116.30	120.80
1	A	260	ASP	CB-CG-OD1	6.38	124.04	118.30
1	A	129	ARG	NE-CZ-NH2	-6.37	117.11	120.30
1	A	144	ASP	CB-CG-OD2	6.33	124.00	118.30
1	A	76	PHE	CB-CG-CD1	6.30	125.21	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	211	TYR	CB-CG-CD1	6.21	124.73	121.00
1	A	76	PHE	CB-CG-CD2	-6.05	116.56	120.80
1	A	224	ASP	CB-CG-OD2	5.71	123.44	118.30
1	A	268	SER	CB-CA-C	5.54	120.62	110.10
1	A	34	ILE	CG1-CB-CG2	-5.47	99.37	111.40
1	A	129	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	120	MET	CG-SD-CE	5.40	108.84	100.20
1	A	103	LEU	CB-CG-CD2	-5.39	101.84	111.00
1	A	35	MET	CG-SD-CE	5.39	108.82	100.20
1	A	86	MET	CG-SD-CE	-5.30	91.72	100.20
1	A	247	ILE	CG1-CB-CG2	-5.17	100.03	111.40
1	A	131	LEU	CB-CG-CD2	5.14	119.74	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2137	75	0
2	A	130	0	0	6	1
All	All	2244	0	2137	75	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

All (75) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:52:MET:SD	1:A:52:MET:CE	2.07	1.42
1:A:86:MET:CG	1:A:86:MET:CE	2.08	1.30
1:A:86:MET:SD	1:A:86:MET:CE	1.03	1.13
1:A:86:MET:SD	1:A:86:MET:HE2	1.63	1.05
1:A:86:MET:SD	1:A:86:MET:HE3	1.63	1.02
1:A:86:MET:SD	1:A:86:MET:HE1	1.63	1.01

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:86:MET:HE2	1:A:86:MET:HG3	1.56	0.85
1:A:244:MET:HE3	1:A:244:MET:HA	1.61	0.82
1:A:86:MET:CG	1:A:86:MET:HE2	1.85	0.81
1:A:208:PHE:HE1	1:A:244:MET:CE	1.94	0.81
1:A:208:PHE:CE1	1:A:244:MET:HE2	2.16	0.80
1:A:225:MET:HE2	1:A:230:CYS:HA	1.62	0.80
1:A:279:GLU:O	1:A:279:GLU:HG2	1.82	0.79
1:A:159:MET:HE2	1:A:163:LYS:HG3	1.66	0.78
1:A:208:PHE:CE1	1:A:244:MET:CE	2.70	0.74
1:A:159:MET:HE3	1:A:163:LYS:HG2	1.72	0.72
1:A:208:PHE:HE1	1:A:244:MET:HE1	1.55	0.72
1:A:243:ARG:O	1:A:247:ILE:HG13	1.91	0.71
1:A:228:ASN:O	1:A:231:LYS:HB3	1.92	0.70
1:A:274:LEU:O	1:A:274:LEU:HG	1.92	0.70
1:A:164:LEU:HD22	1:A:217:ASN:ND2	2.09	0.67
1:A:105:ASN:HD22	1:A:105:ASN:C	1.98	0.66
1:A:151:GLY:O	1:A:157:ARG:HD3	1.96	0.65
1:A:270:ALA:N	1:A:271:PRO:HD3	2.13	0.63
1:A:158:THR:O	1:A:158:THR:HG22	2.00	0.61
1:A:147:LYS:HE3	2:A:2083:HOH:O	2.02	0.59
1:A:225:MET:CE	1:A:230:CYS:HA	2.33	0.59
1:A:159:MET:CE	1:A:163:LYS:CG	2.81	0.58
1:A:144:ASP:OD1	1:A:146:THR:OG1	2.21	0.58
1:A:237:TYR:CE1	1:A:241:LEU:HD11	2.38	0.58
1:A:105:ASN:C	1:A:105:ASN:ND2	2.57	0.57
1:A:159:MET:CE	1:A:163:LYS:HG3	2.35	0.56
1:A:168:VAL:N	1:A:169:PRO:CD	2.70	0.55
1:A:229:GLN:O	1:A:232:GLU:N	2.40	0.54
1:A:159:MET:CE	1:A:163:LYS:HG2	2.39	0.52
1:A:163:LYS:O	1:A:166:LYS:HB2	2.11	0.51
1:A:159:MET:HE3	1:A:163:LYS:CG	2.39	0.50
1:A:160:ASN:OD1	1:A:162:GLU:N	2.43	0.50
1:A:229:GLN:O	1:A:230:CYS:C	2.51	0.49
1:A:222:TYR:HE2	1:A:276:ASP:HB2	1.78	0.48
1:A:91:GLU:OE2	1:A:138:TYR:OH	2.27	0.48
1:A:233:GLY:O	1:A:234:LEU:C	2.50	0.48
1:A:222:TYR:CE2	1:A:276:ASP:HB2	2.49	0.48
1:A:55:ASN:OD1	1:A:57:PRO:HD2	2.14	0.48
1:A:147:LYS:CE	2:A:2083:HOH:O	2.59	0.48
1:A:276:ASP:C	1:A:278:LEU:H	2.16	0.47
1:A:86:MET:CB	1:A:86:MET:CE	2.88	0.46
1:A:229:GLN:O	1:A:231:LYS:N	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:167:THR:O	1:A:170:ILE:N	2.44	0.46
1:A:254:ALA:HB1	1:A:259:ILE:HD12	1.97	0.46
1:A:51:GLU:HB3	2:A:2018:HOH:O	2.15	0.46
1:A:274:LEU:O	1:A:274:LEU:CG	2.63	0.45
1:A:232:GLU:O	1:A:233:GLY:C	2.55	0.45
1:A:160:ASN:ND2	2:A:2088:HOH:O	2.44	0.45
1:A:162:GLU:HG2	1:A:163:LYS:N	2.32	0.44
1:A:53:ASN:HB2	2:A:2018:HOH:O	2.17	0.44
1:A:257:VAL:O	1:A:257:VAL:CG1	2.65	0.43
1:A:244:MET:CE	1:A:244:MET:HA	2.35	0.43
1:A:160:ASN:OD1	1:A:160:ASN:C	2.57	0.43
1:A:229:GLN:C	1:A:231:LYS:N	2.71	0.43
1:A:150:ARG:NH2	2:A:2086:HOH:O	2.50	0.43
1:A:276:ASP:C	1:A:278:LEU:N	2.72	0.42
1:A:239:LYS:O	1:A:243:ARG:HD2	2.19	0.42
1:A:217:ASN:OD1	1:A:221:LYS:HD2	2.20	0.42
1:A:206:ARG:HD3	1:A:206:ARG:HH11	1.68	0.42
1:A:234:LEU:HD12	1:A:234:LEU:HA	1.76	0.42
1:A:157:ARG:HH21	1:A:216:ILE:HG21	1.85	0.41
1:A:273:SER:C	1:A:275:LEU:H	2.24	0.41
1:A:273:SER:O	1:A:275:LEU:N	2.53	0.41
1:A:105:ASN:ND2	1:A:105:ASN:O	2.45	0.41
1:A:208:PHE:CZ	1:A:244:MET:HE2	2.54	0.41
1:A:269:GLN:C	1:A:271:PRO:HD3	2.41	0.41
1:A:150:ARG:NH2	1:A:206:ARG:HD2	2.36	0.41
1:A:208:PHE:CE1	1:A:244:MET:HE1	2.42	0.40
1:A:157:ARG:HH11	1:A:157:ARG:HD3	1.71	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:2044:HOH:O	2:A:2044:HOH:O[7_556]	2.02	0.18

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/289 (90%)	242 (93%)	15 (6%)	4 (2%)	15 6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	270	ALA
1	A	230	CYS
1	A	273	SER
1	A	271	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	235/256 (92%)	224 (95%)	11 (5%)	36 29

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	SER
1	A	76	PHE
1	A	105	ASN
1	A	137	SER
1	A	157	ARG
1	A	159	MET
1	A	162	GLU
1	A	166	LYS
1	A	211	TYR
1	A	269	GLN
1	A	281	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	105	ASN
1	A	140	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	212	ASN
1	A	269	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/289 (91%)	0.26	22 (8%) 11 10	22, 41, 90, 125	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	HIS	6.0
1	A	224	ASP	4.2
1	A	278	LEU	3.9
1	A	234	LEU	3.8
1	A	19	GLY	3.8
1	A	153	ASP	3.2
1	A	230	CYS	3.0
1	A	223	PHE	3.0
1	A	155	VAL	2.9
1	A	227	LYS	2.9
1	A	280	GLN	2.8
1	A	222	TYR	2.7
1	A	269	GLN	2.7
1	A	52	MET	2.7
1	A	277	ALA	2.5
1	A	275	LEU	2.4
1	A	270	ALA	2.4
1	A	150	ARG	2.2
1	A	218	LEU	2.2
1	A	219	LEU	2.2
1	A	274	LEU	2.1
1	A	101	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.