



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 05:19 AM GMT

PDB ID : 4HGH
Title : Crystal structure of P450 BM3 5F5 heme domain variant complexed with styrene (dataset I)
Authors : Shehzad, A.; Panneerselvam, S.; Bocola, M.; Mueller-Dieckmann, J.; Wilmanns, M.; Schwaneberg, U.
Deposited on : 2012-10-08
Resolution : 1.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1097 (1.42-1.38)
Clashscore	79885	1246 (1.42-1.38)
Ramachandran outliers	78287	1206 (1.42-1.38)
Sidechain outliers	78261	1205 (1.42-1.38)
RSRZ outliers	66119	1097 (1.42-1.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	455	
1	B	455	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	GOL	A	503	-	X
4	GOL	A	506	-	X
5	MES	A	504	-	X
6	PEG	A	505	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 8824 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional P-450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	455	Total	C	N	O	S	0	28	0
			3817	2458	636	704	19			
1	B	455	Total	C	N	O	S	0	33	0
			3851	2480	642	710	19			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	87	ALA	PHE	ENGINEERED MUTATION	UNP P14779
A	235	ALA	THR	ENGINEERED MUTATION	UNP P14779
B	87	ALA	PHE	ENGINEERED MUTATION	UNP P14779
B	235	ALA	THR	ENGINEERED MUTATION	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is ETHENYLBENZENE (three-letter code: SYN) (formula: C_8H_8).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			8	8		
3	B	1	Total	C	0	0
			8	8		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONICACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	536	Total	O	0	0
			536	536		
7	B	475	Total	O	0	0
			475	475		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional P-450/NADPH-P450 reductase

Chain A:



- Molecule 1: Bifunctional P-450/NADPH-P450 reductase

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.09Å 149.27Å 65.16Å 90.00° 98.38° 90.00°	Depositor
Resolution (Å)	19.87 – 1.40 19.87 – 1.40	Depositor EDS
% Data completeness (in resolution range)	96.3 (19.87-1.40) 96.4 (19.87-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.190 , 0.204 0.190 , 0.204	Depositor DCC
R_{free} test set	792 reflections (0.38%)	DCC
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 210565 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8824	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, SYN, MES, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.29	8/3978 (0.2%)	1.17	22/5373 (0.4%)
1	B	1.29	13/4016 (0.3%)	1.13	14/5425 (0.3%)
All	All	1.29	21/7994 (0.3%)	1.15	36/10798 (0.3%)

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	GLU	CD-OE2	-8.58	1.16	1.25
1	A	409[A]	GLU	CG-CD	-8.32	1.39	1.51
1	A	409[B]	GLU	CG-CD	-8.32	1.39	1.51
1	B	137	GLU	CD-OE1	-6.99	1.18	1.25
1	B	143	GLU	CD-OE2	-6.11	1.19	1.25
1	B	35	GLU	CG-CD	5.92	1.60	1.51
1	B	137	GLU	CD-OE2	-5.86	1.19	1.25
1	B	293	GLU	CD-OE1	-5.84	1.19	1.25
1	B	348	GLU	CD-OE2	-5.75	1.19	1.25
1	A	352	GLU	CD-OE2	5.72	1.31	1.25
1	B	156	CYS	C-O	5.54	1.33	1.23
1	B	143	GLU	CD-OE1	-5.53	1.19	1.25
1	A	375	ARG	CZ-NH1	5.47	1.40	1.33
1	B	352	GLU	CG-CD	-5.41	1.43	1.51
1	A	166	TYR	CG-CD2	5.37	1.46	1.39
1	A	131	GLU	CD-OE2	5.30	1.31	1.25
1	A	429	TYR	CG-CD1	5.18	1.45	1.39
1	A	275	PHE	CG-CD2	5.03	1.46	1.38
1	B	304	SER	CB-OG	5.02	1.48	1.42
1	B	310[A]	GLN	CG-CD	5.00	1.62	1.51
1	B	310[B]	GLN	CG-CD	5.00	1.62	1.51

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296[A]	ARG	NE-CZ-NH1	7.47	124.03	120.30
1	A	296[B]	ARG	NE-CZ-NH1	7.47	124.03	120.30
1	A	84	ASP	CB-CG-OD2	-6.72	112.25	118.30
1	B	251	ASP	CB-CG-OD2	6.35	124.02	118.30
1	B	50	ARG	NE-CZ-NH2	-5.95	117.32	120.30
1	B	296	ARG	NE-CZ-NH1	5.93	123.26	120.30
1	B	263[A]	ILE	CB-CA-C	-5.91	99.77	111.60
1	B	263[B]	ILE	CB-CA-C	-5.91	99.77	111.60
1	B	23	ASP	CB-CG-OD1	5.86	123.57	118.30
1	A	223	ARG	NE-CZ-NH1	5.79	123.19	120.30
1	A	354[A]	MET	CG-SD-CE	5.78	109.45	100.20
1	A	354[B]	MET	CG-SD-CE	5.78	109.45	100.20
1	A	250	ASP	CB-CG-OD1	5.77	123.49	118.30
1	A	179	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	B	217	ASP	CB-CG-OD1	5.69	123.42	118.30
1	B	223	ARG	NE-CZ-NH1	5.67	123.14	120.30
1	A	77	PHE	CB-CG-CD2	-5.64	116.85	120.80
1	A	199	ASP	CB-CG-OD1	5.63	123.37	118.30
1	A	63	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	338	ASP	CB-CG-OD1	5.56	123.30	118.30
1	A	239	ASN	CB-CA-C	-5.53	99.33	110.40
1	B	425	ASP	CB-CG-OD2	-5.49	113.36	118.30
1	A	375	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	B	237	MET	CG-SD-CE	5.44	108.90	100.20
1	B	23	ASP	CB-CG-OD2	-5.44	113.41	118.30
1	B	242	ASP	CB-CG-OD2	-5.27	113.56	118.30
1	A	250	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	63	ASP	CB-CG-OD1	5.19	122.97	118.30
1	B	66	ARG	NE-CZ-NH2	-5.19	117.71	120.30
1	A	199	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	A	47	ARG	NE-CZ-NH2	5.17	122.89	120.30
1	A	251	ASP	CB-CG-OD1	5.16	122.95	118.30
1	A	42	PHE	CB-CG-CD2	-5.15	117.19	120.80
1	A	67	PHE	CB-CG-CD1	-5.10	117.23	120.80
1	A	238	LEU	CB-CG-CD2	5.09	119.66	111.00
1	B	305	TYR	CB-CG-CD1	-5.04	117.97	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3817	0	0	10	0
1	B	3851	0	0	27	0
2	A	43	0	0	0	0
2	B	43	0	0	0	0
3	A	8	0	0	0	0
3	B	8	0	0	0	0
4	A	12	0	0	0	0
5	A	12	0	0	0	0
5	B	12	0	0	2	0
6	A	7	0	0	1	0
7	A	536	0	0	8	0
7	B	475	0	0	8	0
All	All	8824	0	0	38	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (38) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:150[A]:LEU:CD1	1:B:174:ILE:CD1	1.99	1.39
1:B:309[A]:LYS:NZ	7:B:968:HOH:O	1.61	1.33
1:B:375[B]:ARG:NH2	5:B:503:MES:O2S	1.84	1.09
1:B:261:PHE:O	1:B:265[B]:GLY:N	1.87	1.07
1:B:150[B]:LEU:CD1	1:B:266[B]:HIS:CE1	2.53	0.92
1:A:267[B]:GLU:OE2	7:A:1021:HOH:O	1.95	0.85
1:B:370:ASP:OD2	1:B:375[A]:ARG:NH1	2.10	0.84
1:B:375[A]:ARG:NE	7:B:877:HOH:O	2.13	0.81
1:A:118[B]:MET:SD	7:A:866:HOH:O	2.40	0.79
1:B:118[A]:MET:SD	7:B:802:HOH:O	2.46	0.73
1:A:136:ASP:O	1:A:447:LYS:NZ	2.24	0.71
1:B:158[A]:PHE:CE1	1:B:258:ILE:CG1	2.75	0.70
1:B:261:PHE:O	1:B:265[B]:GLY:CA	2.39	0.69
1:A:3:LYS:CE	7:A:1004:HOH:O	2.41	0.68
1:B:158[B]:PHE:CE2	1:B:262[B]:LEU:CD2	2.83	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:266[A]:HIS:CE1	1:B:267[A]:GLU:OE2	2.54	0.59
1:A:23[A]:ASP:OD2	7:A:1061:HOH:O	2.15	0.59
1:A:438[B]:THR:CG2	1:A:439:LEU:N	2.67	0.57
1:B:267[B]:GLU:CG	1:B:438[B]:THR:CG2	2.83	0.56
1:B:375[A]:ARG:CG	7:B:877:HOH:O	2.54	0.55
1:B:262[B]:LEU:O	1:B:266[B]:HIS:ND1	2.39	0.55
1:B:150[B]:LEU:CD1	1:B:266[B]:HIS:ND1	2.68	0.55
1:A:296[A]:ARG:NH1	7:A:995:HOH:O	2.40	0.54
1:B:128[A]:GLN:NE2	7:B:955:HOH:O	2.42	0.53
1:B:23:ASP:OD2	7:B:1004:HOH:O	2.19	0.52
1:B:194:ASP:OD1	1:B:194:ASP:N	2.45	0.49
1:B:375[A]:ARG:CD	7:B:877:HOH:O	2.60	0.49
1:A:210:LYS:NZ	1:A:214:ASP:OD2	2.48	0.47
1:B:233[B]:LEU:CD2	1:B:261:PHE:CD2	2.99	0.46
1:B:177:MET:SD	1:B:263[B]:ILE:CG1	3.05	0.45
1:B:47:ARG:NE	7:B:900:HOH:O	2.50	0.44
1:A:404[B]:GLN:NE2	7:A:981:HOH:O	2.50	0.44
6:A:505:PEG:C1	7:A:768:HOH:O	2.66	0.43
1:B:150[B]:LEU:CD1	1:B:266[B]:HIS:NE2	2.82	0.43
1:B:150[A]:LEU:CD1	1:B:174:ILE:CG1	2.90	0.42
1:B:375[B]:ARG:NH1	5:B:503:MES:O1S	2.54	0.41
1:B:150[B]:LEU:CD1	1:B:266[B]:HIS:CG	3.05	0.40
1:A:59[B]:LYS:NZ	7:A:693:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/455 (106%)	466 (97%)	15 (3%)	0	100	100
1	B	486/455 (107%)	468 (96%)	18 (4%)	0	100	100
All	All	967/910 (106%)	934 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/397 (107%)	419 (99%)	5 (1%)	82	56
1	B	428/397 (108%)	415 (97%)	13 (3%)	53	16
All	All	852/794 (107%)	834 (98%)	18 (2%)	68	29

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	4	GLU
1	A	13	GLU
1	A	45	PRO
1	A	229	GLN
1	B	3	LYS
1	B	4	GLU
1	B	41	LYS
1	B	148	LEU
1	B	150[A]	LEU
1	B	150[B]	LEU
1	B	185	MET
1	B	192	ASN
1	B	194	ASP
1	B	230	SER
1	B	231	ASP
1	B	267[A]	GLU
1	B	267[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	501	1,7	49,50,50	3.12	19 (38%)	46,82,82	2.46	15 (32%)
3	SYN	A	502	-	8,8,8	1.58	2 (25%)	9,9,9	0.77	0
4	GOL	A	503	-	5,5,5	0.57	0	5,5,5	1.15	1 (20%)
5	MES	A	504	-	12,12,12	1.63	2 (16%)	16,16,16	2.52	6 (37%)
6	PEG	A	505	-	6,6,6	1.02	0	5,5,5	1.62	2 (40%)
4	GOL	A	506	-	5,5,5	0.58	0	5,5,5	0.93	0
2	HEM	B	501	1,7	49,50,50	3.05	19 (38%)	46,82,82	2.82	15 (32%)
3	SYN	B	502	-	8,8,8	1.64	2 (25%)	9,9,9	2.16	2 (22%)
5	MES	B	503	-	12,12,12	2.05	4 (33%)	16,16,16	3.16	8 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1,7	-	0/14/114/114	0/0/8/8
3	SYN	A	502	-	-	0/2/2/2	0/1/1/1
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
5	MES	A	504	-	-	0/6/14/14	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	A	505	-	-	0/4/4/4	0/0/0/0
4	GOL	A	506	-	-	0/4/4/4	0/0/0/0
2	HEM	B	501	1,7	-	0/14/114/114	0/0/8/8
3	SYN	B	502	-	-	0/2/2/2	0/1/1/1
5	MES	B	503	-	-	0/6/14/14	0/1/1/1

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C2D-C1D	-13.44	1.41	1.44
2	B	501	HEM	C3D-C4D	-12.81	1.41	1.44
2	A	501	HEM	C3B-C2B	-6.71	1.32	1.43
2	B	501	HEM	C2D-C1D	-6.42	1.43	1.44
2	A	501	HEM	C2B-C1B	-5.70	1.43	1.44
2	B	501	HEM	C3B-C2B	-5.52	1.34	1.43
2	A	501	HEM	C3C-C2C	-5.46	1.34	1.43
5	B	503	MES	C8-S	-5.27	1.68	1.78
2	B	501	HEM	C3C-C2C	-5.20	1.34	1.43
2	A	501	HEM	FE-NA	4.84	2.13	1.92
2	B	501	HEM	FE-NC	4.56	2.15	1.97
2	A	501	HEM	C3B-CAB	4.44	1.54	1.40
5	A	504	MES	C8-S	-4.35	1.70	1.78
2	B	501	HEM	FE-ND	4.34	2.13	1.97
2	A	501	HEM	FE-NC	4.12	2.13	1.97
2	B	501	HEM	FE-NB	4.06	2.12	1.97
2	B	501	HEM	C3C-CAC	3.93	1.52	1.40
2	A	501	HEM	FE-ND	3.87	2.11	1.97
2	B	501	HEM	CHB-C1B	3.65	1.41	1.35
2	B	501	HEM	FE-NA	3.57	2.07	1.92
2	A	501	HEM	C3C-CAC	3.52	1.51	1.40
2	B	501	HEM	C3B-CAB	3.49	1.51	1.40
2	B	501	HEM	O2A-CGA	-3.48	1.17	1.30
2	A	501	HEM	CHB-C1B	3.22	1.40	1.35
2	A	501	HEM	CAA-C2A	3.21	1.57	1.52
2	B	501	HEM	C4C-NC	-3.20	1.33	1.38
3	A	502	SYN	CAA-CAB	3.14	1.52	1.28
2	A	501	HEM	C3D-C4D	-2.95	1.43	1.44
2	A	501	HEM	O2D-CGD	-2.92	1.19	1.30
2	B	501	HEM	CHA-C4D	2.86	1.39	1.35
5	B	503	MES	C5-N4	-2.83	1.42	1.49
2	A	501	HEM	O2A-CGA	-2.72	1.20	1.30
2	B	501	HEM	CAA-C2A	2.65	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	HEM	C3D-C2D	2.63	1.48	1.43
3	B	502	SYN	CAA-CAB	2.48	1.47	1.28
2	B	501	HEM	CMC-C2C	2.42	1.54	1.47
2	A	501	HEM	CMB-C2B	2.42	1.54	1.47
3	B	502	SYN	CAG-CAH	2.41	1.44	1.39
2	B	501	HEM	C4A-C3A	2.36	1.43	1.40
2	A	501	HEM	CMD-C2D	2.34	1.54	1.47
3	A	502	SYN	CAF-CAH	2.30	1.43	1.39
2	A	501	HEM	CHC-C1C	2.28	1.40	1.36
5	A	504	MES	C5-N4	-2.28	1.43	1.49
5	B	503	MES	C3-N4	-2.22	1.44	1.49
2	A	501	HEM	FE-NB	2.20	2.05	1.97
2	B	501	HEM	CHC-C1C	2.13	1.40	1.36
2	A	501	HEM	CMC-C2C	2.11	1.53	1.47
5	B	503	MES	C7-N4	-2.04	1.45	1.50

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	HEM	C3B-C4B-NB	-12.32	105.18	114.00
2	A	501	HEM	C3B-C4B-NB	-9.96	106.87	114.00
5	B	503	MES	C8-C7-N4	-6.85	100.72	112.44
2	B	501	HEM	C4D-ND-C1D	6.66	111.97	105.16
5	B	503	MES	C5-N4-C3	6.46	122.57	109.75
2	B	501	HEM	C1B-NB-C4B	6.28	111.59	105.16
2	A	501	HEM	CMA-C3A-C4A	-4.87	121.13	128.62
5	A	504	MES	C8-C7-N4	4.81	120.67	112.44
5	A	504	MES	C5-N4-C3	4.76	119.19	109.75
5	A	504	MES	C7-N4-C5	4.59	123.67	111.66
2	A	501	HEM	C3A-C4A-NA	-4.52	106.00	109.41
3	B	502	SYN	CAC-CAD-CAF	4.41	127.52	120.17
2	B	501	HEM	C2D-C1D-ND	-4.32	107.83	112.93
5	B	503	MES	O1-C6-C5	-4.18	106.36	111.34
2	A	501	HEM	C4D-ND-C1D	3.91	109.16	105.16
5	A	504	MES	C2-C3-N4	3.74	114.75	109.96
5	B	503	MES	C7-C8-S	3.56	122.18	112.49
5	A	504	MES	O1-C2-C3	-3.53	107.13	111.34
2	B	501	HEM	CMA-C3A-C4A	-3.52	123.21	128.62
2	A	501	HEM	C2A-C1A-NA	-3.51	104.86	109.73
2	A	501	HEM	C4C-NC-C1C	3.44	109.11	105.53
5	B	503	MES	C7-N4-C3	3.43	120.64	111.66
2	A	501	HEM	C4A-C3A-C2A	3.35	109.33	107.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	HEM	C4A-C3A-C2A	3.34	109.32	107.00
2	A	501	HEM	CHD-C1D-ND	3.34	127.36	124.58
3	B	502	SYN	CAD-CAF-CAH	-3.09	116.67	120.64
2	A	501	HEM	C2D-C1D-ND	-3.00	109.39	112.93
2	B	501	HEM	C3A-C4A-NA	-2.98	107.16	109.41
2	B	501	HEM	C4C-NC-C1C	2.86	108.51	105.53
2	A	501	HEM	CBD-CAD-C3D	-2.77	108.33	114.37
2	A	501	HEM	CHD-C4C-NC	-2.70	122.38	124.73
2	A	501	HEM	C4A-NA-C1A	2.64	110.24	106.76
2	B	501	HEM	CBD-CAD-C3D	-2.55	108.80	114.37
2	B	501	HEM	CMC-C2C-C3C	2.53	132.12	126.16
2	A	501	HEM	C1B-NB-C4B	2.50	107.72	105.16
2	A	501	HEM	C1A-C2A-C3A	2.49	109.50	106.92
2	A	501	HEM	CMA-C3A-C2A	2.43	129.53	124.94
5	B	503	MES	O1-C2-C3	2.43	114.24	111.34
6	A	505	PEG	O2-C2-C1	2.42	122.26	110.61
2	B	501	HEM	O1A-CGA-CBA	-2.42	114.72	123.03
5	B	503	MES	O1S-S-C8	2.41	113.78	106.36
5	B	503	MES	O3S-S-O2S	-2.38	106.35	112.48
2	B	501	HEM	CHC-C4B-NB	2.25	126.46	124.58
6	A	505	PEG	O2-C3-C4	2.23	121.32	110.61
2	B	501	HEM	CMB-C2B-C3B	2.21	131.37	126.16
2	B	501	HEM	O1D-CGD-CBD	-2.15	115.63	123.03
5	A	504	MES	C7-N4-C3	2.10	117.14	111.66
4	A	503	GOL	C3-C2-C1	-2.07	102.11	111.26
2	B	501	HEM	C4A-CHB-C1B	2.06	130.18	127.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	455/455 (100%)	-0.20	12 (2%) 53 58	7, 13, 29, 107	0
1	B	455/455 (100%)	0.07	27 (5%) 22 21	7, 14, 51, 142	0
All	All	910/910 (100%)	-0.07	39 (4%) 35 34	7, 14, 41, 142	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	THR	7.2
1	B	191	ALA	6.6
1	B	1	THR	6.6
1	B	193	PRO	6.1
1	B	2	ILE	5.5
1	B	198	TYR	4.8
1	B	194	ASP	4.5
1	B	136	ASP	4.2
1	B	196	PRO	4.1
1	B	3	LYS	4.0
1	A	2	ILE	3.9
1	B	197	ALA	3.9
1	B	195	ASP	3.8
1	B	192	ASN	3.5
1	B	4	GLU	3.4
1	B	229	GLN	3.3
1	B	169	GLN	3.2
1	B	201	ASN	3.2
1	A	191	ALA	3.0
1	B	199	ASP	2.9
1	B	265[A]	GLY	2.8
1	B	189	GLN	2.7
1	A	229	GLN	2.6
1	B	225	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	227	GLY	2.6
1	A	136	ASP	2.5
1	A	4	GLU	2.4
1	B	231	ASP	2.4
1	A	230	SER	2.3
1	B	239	ASN	2.3
1	B	172	PRO	2.2
1	B	46	GLY	2.2
1	A	158[A]	PHE	2.1
1	A	46	GLY	2.1
1	A	225	ALA	2.1
1	B	214	ASP	2.1
1	A	3	LYS	2.0
1	A	169	GLN	2.0
1	B	158[A]	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MES	A	504	12/12	0.28	13.79	30,43,80,123	0
4	GOL	A	503	6/6	0.16	13.54	21,34,45,47	0
6	PEG	A	505	7/7	0.17	4.87	29,32,38,41	0
4	GOL	A	506	6/6	0.14	3.54	26,28,30,41	0
5	MES	B	503	12/12	0.16	1.85	34,45,71,80	0
3	SYN	B	502	8/8	0.13	1.55	26,31,41,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SYN	A	502	8/8	0.10	1.24	23,26,29,30	0
2	HEM	B	501	43/43	0.06	-0.47	6,8,10,22	0
2	HEM	A	501	43/43	0.05	-0.59	5,7,10,17	0

6.5 Other polymers ⓘ

There are no such residues in this entry.