



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 10:22 AM GMT

PDB ID : 1HM3
Title : ACTIVE SITE OF CHONDROITINASE AC LYASE REVEALED BY THE
STRUCTURE OF ENZYME-OLIGOSACCHARIDECOMPLEXES AND
MUTAGENESIS
Authors : Huang, W.; Boju, L.; Tkalec, L.; Su, H.; Yang, H.O.; Gunay, N.S.; Linhardt,
R.J.; Kim, Y.S.; Matte, A.; Cygler, M.
Deposited on : 2000-12-04
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	700	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5822 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHONDROITINASE AC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	674	Total	C	N	O	S	0	0	0
			5382	3445	919	1004	14			

- Molecule 2 is a polymer of unknown type called SUGAR (RAM-MAN-GCU-XYS-MXY).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	5	Total	C	O	0	0
			53	30	23		

- Molecule 3 is a polymer of unknown type called SUGAR (RAM-MAN-GCU).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	3	Total	C	O	0	0
			33	18	15		

- Molecule 4 is a polymer of unknown type called SUGAR (HYALURONIC ACID DISACCHARIDE).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			27	14	1	12		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	326	Total 326	O 326	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CHONDROITINASE AC

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.90Å 86.90Å 192.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.80 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.10) 96.6 (19.80-2.10)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.68 (at 2.09Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.226 , 0.254 0.224 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.676	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 43544 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5822	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, NAG, MXY, CA, RAM, GCU, BDP, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/5520	0.61	1/7487 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	1	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	405	GLU	N-CA-C	-5.30	96.68	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	702	RAM	C1

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5382	0	5240	101	0
2	A	53	0	43	2	0
3	A	33	0	25	1	0
4	A	27	0	21	0	0
5	A	1	0	0	0	0
6	A	326	0	0	1	0
All	All	5822	0	5329	101	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (101) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:499:SER:HB3	1:A:520:LEU:HD23	1.64	0.79
1:A:212:TYR:H	1:A:217:GLN:HE22	1.30	0.78
1:A:212:TYR:N	1:A:217:GLN:HE22	1.85	0.75
1:A:106:PHE:CE2	1:A:149:LYS:HE3	2.28	0.69
1:A:544:GLY:O	1:A:558:VAL:HG22	1.93	0.68
1:A:181:LEU:O	1:A:184:PHE:HB3	1.94	0.67
1:A:247:ASN:HB2	1:A:311:ILE:HD11	1.77	0.67
1:A:212:TYR:H	1:A:217:GLN:NE2	1.95	0.65
1:A:71:ASP:O	1:A:124:ARG:HG2	1.97	0.65
1:A:203:LEU:HD13	1:A:245:LEU:HD22	1.78	0.63
1:A:54:LEU:HD21	1:A:105:VAL:HG22	1.81	0.63
1:A:690:GLN:HB3	1:A:691:GLU:OE2	1.99	0.63
1:A:169:GLU:CD	1:A:169:GLU:H	2.02	0.62
1:A:509:ILE:HD13	1:A:510:THR:N	2.15	0.62
1:A:224:GLN:O	1:A:225:HIS:HB2	2.01	0.61
1:A:205:TYR:HB3	1:A:206:PRO:HD3	1.82	0.61
1:A:368:ARG:HD3	1:A:430:GLN:HA	1.83	0.60
1:A:415:ARG:HD3	1:A:480:ALA:CB	2.33	0.59
1:A:149:LYS:HB2	1:A:149:LYS:NZ	2.18	0.59
1:A:36:MET:HG3	1:A:146:TYR:CD1	2.39	0.58
1:A:403:VAL:CG1	1:A:558:VAL:HG23	2.34	0.57
1:A:32:MET:HE2	1:A:188:LEU:HB3	1.86	0.57
1:A:509:ILE:C	1:A:509:ILE:HD13	2.25	0.57
1:A:685:ILE:HG12	1:A:699:LEU:HD11	1.87	0.56
1:A:506:ARG:HG3	1:A:561:ASP:OD1	2.07	0.55
1:A:104:LYS:O	1:A:108:GLN:HG3	2.07	0.54
1:A:50:ALA:O	1:A:54:LEU:HB2	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:532:GLY:O	1:A:533:ALA:HB2	2.09	0.53
1:A:509:ILE:HB	1:A:540:GLN:NE2	2.23	0.53
1:A:231:ILE:HB	1:A:285:VAL:HB	1.90	0.53
1:A:32:MET:CE	1:A:188:LEU:HB3	2.38	0.52
1:A:479:ASN:OD1	2:A:701:MAN:H5	2.08	0.52
1:A:106:PHE:CE1	1:A:149:LYS:HG3	2.44	0.52
1:A:106:PHE:CD1	1:A:149:LYS:HG3	2.45	0.52
1:A:613:TYR:CE1	1:A:650:ILE:HD11	2.44	0.52
1:A:133:THR:HB	1:A:134:PRO:HD3	1.92	0.51
1:A:85:LEU:O	1:A:89:ILE:HG13	2.10	0.51
1:A:356:TYR:CD2	1:A:584:PRO:HB2	2.46	0.51
1:A:47:ASP:HB3	1:A:98:HIS:NE2	2.25	0.50
1:A:400:ILE:HA	1:A:492:TRP:CZ2	2.47	0.50
1:A:228:GLN:HB2	1:A:368:ARG:HB2	1.94	0.50
1:A:415:ARG:HD3	1:A:480:ALA:HB3	1.93	0.50
1:A:415:ARG:HH21	1:A:454:ASP:HB3	1.77	0.49
1:A:156:VAL:O	1:A:160:THR:HG23	2.12	0.49
1:A:506:ARG:HG2	1:A:507:GLY:N	2.27	0.49
1:A:363:VAL:HG22	1:A:364:SER:N	2.28	0.49
1:A:583:LEU:HD12	1:A:583:LEU:N	2.27	0.49
1:A:102:ASP:OD2	1:A:104:LYS:HB3	2.12	0.49
1:A:601:LYS:HE2	1:A:613:TYR:HD2	1.78	0.49
1:A:247:ASN:CB	1:A:311:ILE:HD11	2.41	0.48
1:A:647:ILE:HG12	1:A:656:ILE:HG12	1.94	0.48
1:A:547:PHE:CG	1:A:555:LYS:HG2	2.49	0.48
1:A:415:ARG:HH22	2:A:703:GCU:C6	2.27	0.48
1:A:603:LEU:HD11	1:A:613:TYR:HB2	1.95	0.47
1:A:131:ILE:O	1:A:135:GLN:HG3	2.14	0.47
1:A:588:LYS:NZ	1:A:588:LYS:HB2	2.29	0.47
1:A:629:LYS:HD3	1:A:636:GLU:OE2	2.14	0.47
1:A:499:SER:HB3	1:A:520:LEU:CD2	2.42	0.47
1:A:56:THR:HG23	1:A:63:TRP:HE1	1.80	0.47
1:A:402:PRO:CG	1:A:558:VAL:HG21	2.45	0.46
1:A:601:LYS:HE3	1:A:603:LEU:HD23	1.97	0.46
1:A:56:THR:HG23	1:A:63:TRP:CD1	2.51	0.46
1:A:601:LYS:HE3	1:A:603:LEU:CD2	2.45	0.46
1:A:302:LYS:HA	1:A:305:LEU:HB2	1.97	0.46
1:A:372:SER:HB2	1:A:423:LEU:HD13	1.97	0.46
1:A:43:LEU:HD13	1:A:94:GLU:HG2	1.98	0.45
1:A:473:GLY:HA3	1:A:578:TYR:CE2	2.51	0.45
1:A:261:LEU:HD13	1:A:261:LEU:C	2.37	0.45
1:A:56:THR:HG23	1:A:63:TRP:NE1	2.31	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:381:ARG:HA	1:A:423:LEU:HD21	1.99	0.45
1:A:293:PRO:HG3	1:A:546:TRP:CD2	2.51	0.45
1:A:120:ASP:N	1:A:121:PRO:HD3	2.31	0.45
1:A:637:ILE:HD13	1:A:647:ILE:HD12	1.97	0.45
1:A:228:GLN:O	1:A:230:GLN:HG2	2.16	0.45
1:A:593:LYS:HD3	1:A:593:LYS:O	2.16	0.45
1:A:149:LYS:HB2	1:A:149:LYS:HZ2	1.81	0.44
1:A:297:ASN:HB2	1:A:398:TYR:CE1	2.52	0.44
1:A:224:GLN:HG2	1:A:234:TYR:CG	2.53	0.44
1:A:415:ARG:HD2	1:A:482:GLU:OE1	2.18	0.43
1:A:529:PHE:CE2	1:A:535:LEU:HD21	2.54	0.43
1:A:134:PRO:HB2	1:A:182:HIS:CD2	2.53	0.43
1:A:126:TRP:CZ3	1:A:171:LYS:HE2	2.54	0.43
1:A:314:LYS:HG3	6:A:853:HOH:O	2.19	0.43
1:A:328:SER:O	3:A:706:MAN:H5	2.19	0.43
1:A:409:ILE:O	1:A:412:ILE:HG12	2.19	0.42
1:A:114:LYS:NZ	1:A:118:ASP:OD1	2.52	0.42
1:A:676:LYS:HE2	1:A:677:THR:HG22	2.01	0.42
1:A:149:LYS:HA	1:A:150:PRO:HD3	1.86	0.42
1:A:372:SER:HB2	1:A:423:LEU:CD1	2.49	0.42
1:A:168:PRO:HG2	1:A:169:GLU:OE2	2.19	0.42
1:A:160:THR:O	1:A:163:MET:HB2	2.19	0.42
1:A:210:VAL:HB	1:A:213:GLU:CG	2.49	0.42
1:A:210:VAL:HB	1:A:213:GLU:HG2	2.02	0.41
1:A:218:TYR:CD1	1:A:691:GLU:HB3	2.55	0.41
1:A:105:VAL:HA	1:A:108:GLN:HE21	1.86	0.41
1:A:252:THR:HB	1:A:253:PRO:HD2	2.02	0.41
1:A:478:SER:O	1:A:572:ARG:HA	2.21	0.40
1:A:363:VAL:HG22	1:A:364:SER:H	1.86	0.40
1:A:507:GLY:HA2	1:A:540:GLN:HA	2.02	0.40
1:A:506:ARG:HD2	1:A:561:ASP:OD1	2.21	0.40
1:A:413:THR:HA	1:A:485:THR:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	672/700 (96%)	635 (94%)	35 (5%)	2 (0%)	50	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	TYR
1	A	501	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	563/598 (94%)	551 (98%)	12 (2%)	66	70

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	HIS
1	A	137	LEU
1	A	149	LYS
1	A	169	GLU
1	A	195	LEU
1	A	268	TYR
1	A	306	LEU
1	A	401	MET
1	A	404	TRP
1	A	468	GLU
1	A	509	ILE
1	A	543	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	108	GLN
1	A	217	GLN
1	A	540	GLN
1	A	551	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MAN	A	701	1,2	10,11,12	0.42	0	11,15,17	0.64	0
2	RAM	A	702	2	9,10,11	1.52	1 (11%)	10,14,16	0.78	0
2	GCU	A	703	2	11,12,13	1.61	3 (27%)	13,17,19	1.39	2 (15%)
2	XYP	A	704	2	8,9,10	0.79	0	8,12,14	0.88	1 (12%)
2	MXY	A	705	2	10,11,12	0.87	0	11,15,17	0.66	0
3	MAN	A	706	1,3	10,11,12	0.40	0	11,15,17	0.57	0
3	RAM	A	707	3	9,10,11	1.14	1 (11%)	10,14,16	0.89	0
3	GCU	A	708	3	11,12,13	1.56	2 (18%)	13,17,19	1.31	2 (15%)
4	NAG	A	709	4	15,15,15	0.47	0	21,21,21	0.74	0
4	BDP	A	710	4	11,12,13	1.46	2 (18%)	13,17,19	1.55	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	A	701	1,2	-	0/2/19/22	0/1/1/1
2	RAM	A	702	2	1/1/4/5	0/0/17/20	0/1/1/1
2	GCU	A	703	2	-	0/4/21/24	0/1/1/1
2	XYP	A	704	2	-	0/0/14/17	0/1/1/1
2	MXY	A	705	2	-	0/2/19/22	0/1/1/1
3	MAN	A	706	1,3	-	0/2/19/22	0/1/1/1
3	RAM	A	707	3	-	0/0/17/20	0/1/1/1
3	GCU	A	708	3	-	0/4/21/24	0/1/1/1
4	NAG	A	709	4	-	0/6/26/26	0/1/1/1
4	BDP	A	710	4	-	0/4/21/24	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	708	GCU	O6A-C6	4.06	1.36	1.22
4	A	710	BDP	O6A-C6	3.80	1.35	1.22
2	A	703	GCU	O6A-C6	3.75	1.35	1.22
2	A	702	RAM	C4-C5	3.30	1.59	1.52
2	A	703	GCU	O6B-C6	-2.51	1.21	1.30
4	A	710	BDP	O6B-C6	-2.40	1.21	1.30
3	A	707	RAM	C4-C5	2.21	1.57	1.52
2	A	703	GCU	C4-C5	2.19	1.57	1.53
3	A	708	GCU	O6B-C6	-2.17	1.22	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	710	BDP	C4-C5-C6	2.87	115.89	110.84
4	A	710	BDP	O6B-C6-C5	2.64	123.13	113.49
3	A	708	GCU	O6B-C6-C5	2.62	123.04	113.49
2	A	703	GCU	O6B-C6-C5	2.62	123.03	113.49
4	A	710	BDP	C4-C3-C2	-2.58	107.04	110.50
2	A	703	GCU	C4-C5-C6	2.46	115.15	110.84
2	A	704	XYP	C5B-C4B-C3B	2.04	112.28	109.71
3	A	708	GCU	C4-C3-C2	-2.02	107.79	110.50

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	702	RAM	C1

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	674/700 (96%)	0.03	16 (2%) 56 61	15, 25, 38, 47	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ALA	5.4
1	A	43	LEU	3.5
1	A	26	GLY	3.4
1	A	593	LYS	3.3
1	A	212	TYR	3.2
1	A	51	GLU	3.0
1	A	509	ILE	3.0
1	A	597	GLY	2.9
1	A	193	GLU	2.9
1	A	598	THR	2.6
1	A	98	HIS	2.6
1	A	676	LYS	2.4
1	A	652	GLY	2.4
1	A	74	THR	2.2
1	A	96	ASP	2.2
1	A	675	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	RAM	A	707	10/11	0.15	10.33	27,28,29,29	0
4	BDP	A	710	12/13	0.22	10.11	33,35,36,36	0
2	RAM	A	702	10/11	0.61	5.57	49,51,51,52	0
3	MAN	A	706	11/12	0.16	3.39	27,28,29,31	0
3	GCU	A	708	12/13	0.17	2.05	30,32,34,34	0
2	MAN	A	701	11/12	0.19	1.95	37,40,42,46	0
2	GCU	A	703	12/13	0.11	-0.28	37,39,40,41	0
4	NAG	A	709	15/15	0.09	-0.53	24,26,27,29	0
2	XYP	A	704	9/10	0.20	-	43,44,46,47	0
2	MXY	A	705	11/12	0.35	-	48,49,50,50	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CA	A	1801	1/1	0.03	-3.57	22,22,22,22	0

6.5 Other polymers ⓘ

There are no such residues in this entry.