



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 12:39 PM GMT

PDB ID : 3HM7  
Title : Crystal structure of allantoinase from *Bacillus halodurans* C-125  
Authors : Patskovsky, Y.; Romero, R.; Rutter, M.; Miller, S.; Wasserman, S.R.; Sauder, J.M.; Raushel, F.M.; Burley, S.K.; Almo, S.C.; New York Structural Genomix Research Consortium (Nysgxrc); New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-05-28  
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	448	
1	B	448	
1	C	448	
1	D	448	
1	E	448	
1	F	448	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	ZN	E	447	-	X
2	ZN	F	447	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20901 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allantoinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	4	0
			3425	2166	593	649	17			
1	B	437	Total	C	N	O	S	0	5	0
			3437	2173	594	653	17			
1	C	435	Total	C	N	O	S	0	3	0
			3410	2154	588	651	17			
1	D	436	Total	C	N	O	S	0	2	0
			3415	2157	591	650	17			
1	E	436	Total	C	N	O	S	0	2	0
			3412	2156	590	649	17			
1	F	436	Total	C	N	O	S	0	4	0
			3425	2165	592	651	17			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
A	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
A	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
A	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
A	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
A	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
B	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
B	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
B	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
B	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
B	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
C	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
C	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
C	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
C	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
C	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
D	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
D	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
D	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
D	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
D	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
E	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
E	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
E	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
E	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
E	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
F	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
F	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
F	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
F	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	F	1	Total Zn 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	53	Total O 53 53	0	0
3	B	69	Total O 69 69	0	0
3	C	39	Total O 39 39	0	0
3	D	47	Total O 47 47	0	0
3	E	74	Total O 74 74	0	0
3	F	89	Total O 89 89	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Allantoinase

Chain A:




#### • Molecule 1: Allantoinase


Chain B:



#### • Molecule 1: Allantoinase


Chain C:






• Molecule 1: Allantoinase

Chain D:




• Molecule 1: Allantoinase

Chain E:



• Molecule 1: Allantoinase

Chain F:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.66Å 157.66Å 418.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.60 27.39 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.60) 100.0 (27.39-2.60)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.94 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, $R_{free}$	0.240 , 0.267 0.244 , 0.269	Depositor DCC
$R_{free}$ test set	2853 reflections (3.11%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	37 of 94701 reflections (0.039%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20901	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.84 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.4571e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/3507	0.60	0/4749
1	B	0.44	0/3523	0.61	1/4770 (0.0%)
1	C	0.43	0/3485	0.62	0/4720
1	D	0.42	0/3488	0.59	0/4724
1	E	0.43	0/3488	0.59	0/4724
1	F	0.41	0/3508	0.61	1/4752 (0.0%)
All	All	0.42	0/20999	0.60	2/28439 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	395	LEU	CA-CB-CG	6.19	129.54	115.30
1	F	262	VAL	CB-CA-C	-5.13	101.65	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	159	ASP	Peptide
1	A	160	ASP	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3404	27	0
1	B	3437	0	3411	25	0
1	C	3410	0	3376	32	0
1	D	3415	0	3379	29	0
1	E	3412	0	3380	16	0
1	F	3425	0	3393	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	53	0	0	0	0
3	B	69	0	0	0	0
3	C	39	0	0	0	0
3	D	47	0	0	0	0
3	E	74	0	0	0	0
3	F	89	0	0	0	0
All	All	20901	0	20343	138	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:67:ASN:ND2	1:D:98:ASN:HB2	1.96	0.80
1:F:75:GLU:HG2	1:F:318:PRO:HG3	1.69	0.73
1:B:75:GLU:HG2	1:B:318:PRO:HG3	1.76	0.68
1:F:67:ASN:ND2	1:F:98:ASN:HB2	2.09	0.67
1:E:67:ASN:ND2	1:E:98:ASN:HB2	2.09	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/448 (98%)	410 (94%)	25 (6%)	3 (1%)	30	58
1	B	440/448 (98%)	425 (97%)	13 (3%)	2 (0%)	38	67
1	C	436/448 (97%)	397 (91%)	35 (8%)	4 (1%)	25	49
1	D	436/448 (97%)	416 (95%)	18 (4%)	2 (0%)	38	67
1	E	436/448 (97%)	419 (96%)	15 (3%)	2 (0%)	38	67
1	F	438/448 (98%)	424 (97%)	12 (3%)	2 (0%)	38	67
All	All	2624/2688 (98%)	2491 (95%)	118 (4%)	15 (1%)	33	63

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	PRO
1	A	242	HIS
1	B	242	HIS
1	C	242	HIS
1	D	242	HIS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/382 (98%)	343 (92%)	32 (8%)	15	29
1	B	377/382 (99%)	351 (93%)	26 (7%)	22	42
1	C	373/382 (98%)	336 (90%)	37 (10%)	11	21
1	D	373/382 (98%)	340 (91%)	33 (9%)	14	27
1	E	373/382 (98%)	350 (94%)	23 (6%)	26	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	375/382 (98%)	347 (92%)	28 (8%)	19	36
All	All	2246/2292 (98%)	2067 (92%)	179 (8%)	17	33

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	304	LEU
1	D	113	ARG
1	F	242	HIS
1	C	328	THR
1	C	393	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	418	GLN
1	D	352	HIS
1	F	383	GLN
1	D	162	GLN
1	B	67	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	436/448 (97%)	0.06	19 (4%) 33 29	22, 38, 54, 77	0
1	B	437/448 (97%)	-0.19	9 (2%) 60 58	28, 37, 52, 66	0
1	C	435/448 (97%)	1.07	83 (19%) 2 1	22, 39, 51, 74	0
1	D	436/448 (97%)	0.02	25 (5%) 23 19	26, 38, 53, 70	0
1	E	436/448 (97%)	-0.27	9 (2%) 60 58	25, 37, 47, 68	0
1	F	436/448 (97%)	-0.31	2 (0%) 88 90	26, 37, 48, 64	0
All	All	2616/2688 (97%)	0.06	147 (5%) 24 20	22, 37, 52, 77	0

The worst 5 of 147 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	322	GLN	6.8
1	C	158	THR	6.7
1	C	402	ASN	6.5
1	C	156	CYS	6.2
1	A	157	GLY	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	F	447	1/1	0.38	4.28	39,39,39,39	1
2	ZN	E	447	1/1	0.29	2.19	39,39,39,39	1
2	ZN	D	447	1/1	0.28	1.57	40,40,40,40	1
2	ZN	B	447	1/1	0.23	0.81	39,39,39,39	1
2	ZN	A	447	1/1	0.18	-1.26	43,43,43,43	1
2	ZN	C	447	1/1	0.16	-3.13	50,50,50,50	1

## 6.5 Other polymers

There are no such residues in this entry.