



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 12:42 PM GMT

PDB ID : 3HMX  
Title : Crystal structure of ustekinumab FAB/IL-12 complex  
Authors : Luo, J.  
Deposited on : 2009-05-29  
Resolution : 3.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	306	
2	B	197	
3	L	214	
4	H	226	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7078 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	0	0
			2331	1475	380	464	12			

- Molecule 2 is a protein called Interleukin-12 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	176	Total	C	N	O	S	0	0	0
			1351	852	229	254	16			

- Molecule 3 is a protein called USTEKINUMAB FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	214	Total	C	N	O	S	0	0	0
			1649	1034	274	336	5			

- Molecule 4 is a protein called USTEKINUMAB FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	217	Total	C	N	O	S	0	0	0
			1642	1045	273	316	8			

- Molecule 5 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	15	Total	O	0	0
			15	15		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total 2	O 2	0	0
6	L	11	Total 11	O 11	0	0
6	H	16	Total 16	O 16	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-12 subunit beta




- Molecule 2: Interleukin-12 subunit alpha



- Molecule 3: USTEKINUMAB FAB LIGHT CHAIN



- Molecule 4: USTEKINUMAB FAB HEAVY CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.09Å 116.44Å 182.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.66 – 3.00 26.66 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (26.66-3.00) 99.7 (26.66-3.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.73 (at 3.00Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.211 , 0.303 0.207 , 0.297	Depositor DCC
$R_{free}$ test set	1168 reflections (4.83%)	DCC
Wilson B-factor (Å <sup>2</sup> )	87.0	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 57.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 24242 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7078	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2388	0.47	0/3249
2	B	0.26	0/1369	0.44	0/1845
3	L	0.29	0/1686	0.49	0/2290
4	H	0.30	0/1685	0.49	0/2294
All	All	0.28	0/7128	0.47	0/9678

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2331	0	0	10	0
2	B	1351	0	0	4	0
3	L	1649	0	0	10	0
4	H	1642	0	0	5	0
5	A	61	0	0	0	0
6	A	15	0	0	1	0
6	B	2	0	0	0	0
6	H	16	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	L	11	0	0	0	0
All	All	7078	0	0	28	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (28) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:H:88:ALA:O	4:H:91:THR:CG2	2.55	0.55
4:H:153:PRO:O	4:H:206:HIS:CE1	2.61	0.54
1:A:5:LYS:NZ	6:A:307:HOH:O	2.40	0.53
2:B:91:SER:C	2:B:93:LYS:N	2.62	0.53
3:L:165:GLU:O	3:L:166:GLN:C	2.50	0.49
3:L:6:GLN:N	3:L:100:GLN:OE1	2.46	0.49
1:A:236:TYR:CD1	1:A:244:HIS:CG	3.02	0.48
1:A:181:GLU:OE1	2:B:183:ARG:NH1	2.48	0.46
1:A:141:SER:O	1:A:142:ASP:C	2.53	0.46
3:L:188:LYS:O	3:L:189:HIS:CG	2.69	0.46
4:H:133:SER:OG	4:H:134:SER:N	2.49	0.46
3:L:136:LEU:CD1	3:L:136:LEU:N	2.80	0.45
1:A:15:TRP:CA	1:A:15:TRP:CE3	2.99	0.44
3:L:210:ASN:O	3:L:211:ARG:CG	2.65	0.44
2:B:38:GLU:O	2:B:40:TYR:N	2.51	0.44
1:A:201:TYR:CD1	1:A:201:TYR:N	2.84	0.44
3:L:198:HIS:CD2	3:L:200:GLY:N	2.88	0.42
3:L:30:SER:O	3:L:68:GLY:N	2.54	0.41
3:L:61:ARG:NH1	3:L:82:ASP:OD2	2.54	0.41
1:A:142:ASP:OD1	1:A:142:ASP:O	2.38	0.41
4:H:207:LYS:O	4:H:209:SER:N	2.53	0.41
1:A:159:ARG:CG	1:A:160:GLY:N	2.83	0.41
3:L:108:ARG:NE	3:L:109:THR:O	2.53	0.41
3:L:125:LEU:O	3:L:127:SER:N	2.54	0.41
4:H:98:ARG:NH1	4:H:107:ASP:OD2	2.54	0.41
1:A:103:ASN:O	1:A:104:LYS:C	2.60	0.40
2:B:87:SER:OG	2:B:88:CYS:N	2.53	0.40
1:A:98:GLN:O	1:A:100:GLU:N	2.54	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/306 (94%)	240 (83%)	36 (12%)	13 (4%)	4	22
2	B	168/197 (85%)	134 (80%)	29 (17%)	5 (3%)	7	34
3	L	212/214 (99%)	181 (85%)	27 (13%)	4 (2%)	12	51
4	H	213/226 (94%)	190 (89%)	18 (8%)	5 (2%)	10	43
All	All	882/943 (94%)	745 (84%)	110 (12%)	27 (3%)	7	34

All (27) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	PRO
1	A	217	LYS
1	A	255	VAL
2	B	54	LYS
3	L	138	ASN
3	L	211	ARG
2	B	96	PHE
1	A	87	ASP
1	A	102	LYS
1	A	161	ASP
1	A	295	SER
2	B	92	ARG
2	B	129	ARG
2	B	150	PHE
3	L	40	PRO
4	H	41	PRO
4	H	133	SER
4	H	167	SER
1	A	34	ASP
1	A	141	SER
3	L	166	GLN
4	H	208	PRO
1	A	142	ASP
1	A	224	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
4	H	72	VAL
1	A	160	GLY
1	A	31	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/277 (95%)	236 (90%)	26 (10%)	11	40
2	B	147/183 (80%)	134 (91%)	13 (9%)	14	48
3	L	187/188 (100%)	168 (90%)	19 (10%)	11	38
4	H	184/195 (94%)	174 (95%)	10 (5%)	31	74
All	All	780/843 (92%)	712 (91%)	68 (9%)	15	49

All (68) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLU
1	A	15	TRP
1	A	18	ASP
1	A	24	VAL
1	A	36	ILE
1	A	39	THR
1	A	42	GLN
1	A	44	SER
1	A	56	GLN
1	A	58	LYS
1	A	89	ILE
1	A	94	ILE
1	A	107	LEU
1	A	119	THR
1	A	124	THR
1	A	131	THR
1	A	135	LYS
1	A	204	SER
1	A	211	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	219	LEU
1	A	235	GLU
1	A	241	SER
1	A	248	SER
1	A	249	LEU
1	A	279	ARG
1	A	303	VAL
2	B	16	LEU
2	B	54	LYS
2	B	64	LEU
2	B	70	LYS
2	B	94	THR
2	B	96	PHE
2	B	133	LEU
2	B	135	GLN
2	B	136	ASN
2	B	173	LEU
2	B	183	ARG
2	B	191	MET
2	B	195	ASN
3	L	11	LEU
3	L	42	LYS
3	L	56	SER
3	L	77	SER
3	L	95	PRO
3	L	100	GLN
3	L	105	GLU
3	L	108	ARG
3	L	133	VAL
3	L	150	VAL
3	L	154	LEU
3	L	161	GLU
3	L	163	VAL
3	L	172	THR
3	L	175	LEU
3	L	180	THR
3	L	205	VAL
3	L	206	THR
3	L	213	GLU
4	H	38	ARG
4	H	50	ILE
4	H	91	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
4	H	96	CYS
4	H	107	ASP
4	H	157	THR
4	H	188	VAL
4	H	189	THR
4	H	198	GLN
4	H	220	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	A	501	1,5	12,14,15	0.55	0	15,19,21	0.88	0
5	NAG	A	502	5	12,14,15	0.68	0	15,19,21	1.37	3 (20%)
5	BMA	A	503	5	10,11,12	0.81	0	11,15,17	2.50	6 (54%)
5	MAN	A	504	5	10,11,12	1.24	1 (10%)	11,15,17	3.17	6 (54%)
5	MAN	A	505	5	10,11,12	0.72	0	11,15,17	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	501	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	502	5	-	0/6/23/26	0/1/1/1
5	BMA	A	503	5	-	0/2/19/22	0/1/1/1
5	MAN	A	504	5	-	0/2/19/22	0/1/1/1
5	MAN	A	505	5	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	504	MAN	C6-C5	2.06	1.59	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	504	MAN	O5-C5-C6	7.67	115.03	106.98
5	A	503	BMA	O5-C5-C6	4.86	112.08	106.98
5	A	504	MAN	O4-C4-C3	3.69	118.64	110.35
5	A	503	BMA	C4-C3-C2	-3.42	105.91	110.50
5	A	504	MAN	O5-C5-C4	-3.12	106.69	110.65
5	A	504	MAN	O2-C2-C3	3.05	116.75	110.18
5	A	503	BMA	O2-C2-C3	3.05	116.75	110.18
5	A	503	BMA	O5-C5-C4	3.02	114.49	110.65
5	A	504	MAN	O6-C6-C5	3.01	121.73	111.36
5	A	502	NAG	O5-C5-C4	2.52	113.86	110.65
5	A	503	BMA	O4-C4-C5	2.50	115.88	109.28
5	A	503	BMA	O6-C6-C5	2.46	119.82	111.36
5	A	502	NAG	C2-N2-C7	-2.30	119.23	123.09
5	A	504	MAN	C4-C3-C2	2.13	113.37	110.50
5	A	502	NAG	C6-C5-C4	-2.05	108.05	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	295/306 (96%)	-0.06	3 (1%) 79 22	60, 97, 200, 255	0
2	B	176/197 (89%)	-0.18	0 100 100	70, 106, 181, 238	0
3	L	214/214 (100%)	-0.13	1 (0%) 88 36	52, 90, 121, 164	0
4	H	217/226 (96%)	-0.17	0 100 100	45, 81, 134, 178	0
All	All	902/943 (95%)	-0.13	4 (0%) 90 41	45, 92, 175, 255	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	LYS	3.1
1	A	228	ARG	2.5
1	A	224	LEU	2.5
3	L	134	CYS	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	MAN	A	504	11/12	0.35	-0.49	147,154,159,159	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MAN	A	505	11/12	0.20	-1.36	119,135,140,144	0
5	BMA	A	503	11/12	0.24	-1.41	109,116,129,137	0
5	NAG	A	501	14/15	0.14	-1.49	84,92,105,106	0
5	NAG	A	502	14/15	0.12	-7.18	80,98,109,111	0

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.