



# wwPDB X-ray Structure Validation Summary Report (i)

Feb 26, 2014 – 03:28 PM GMT

PDB ID : 3HOW  
Title : Complete RNA polymerase II elongation complex III with a T-U mismatch and a frayed RNA 3'-uridine  
Authors : Sydow, J.F.; Brueckner, F.; Cheung, A.C.M.; Damsma, G.E.; Dengl, S.; Lehmann, E.; Vassylyev, D.; Cramer, P.  
Deposited on : 2009-06-03  
Resolution : 3.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1020 (3.86-3.34)
Clashscore	79885	1155 (3.80-3.40)
Ramachandran outliers	78287	1109 (3.80-3.40)
Sidechain outliers	78261	1108 (3.80-3.40)
RSRZ outliers	66119	1000 (3.84-3.36)
RNA backbone	1838	1012 (4.40-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	1733	
2	B	1224	
3	C	347	
4	D	221	
5	E	215	
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
13	2	12	
14	1	26	
15	3	18	

## 2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 31876 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1416	Total	C	N	O	S	0	0	0
			11143	7021	1949	2111	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1112	Total	C	N	O	S	0	0	0
			8838	5594	1550	1639	55			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	266	Total	C	N	O	S	0	0	0
			2095	1317	348	417	13			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-28	MET	-	EXPRESSION TAG	UNP P16370
C	-27	GLY	-	EXPRESSION TAG	UNP P16370
C	-26	SER	-	EXPRESSION TAG	UNP P16370
C	-25	HIS	-	EXPRESSION TAG	UNP P16370
C	-24	HIS	-	EXPRESSION TAG	UNP P16370
C	-23	HIS	-	EXPRESSION TAG	UNP P16370
C	-22	HIS	-	EXPRESSION TAG	UNP P16370
C	-21	HIS	-	EXPRESSION TAG	UNP P16370
C	-20	HIS	-	EXPRESSION TAG	UNP P16370
C	-19	SER	-	EXPRESSION TAG	UNP P16370
C	-18	ASN	-	EXPRESSION TAG	UNP P16370
C	-17	SER	-	EXPRESSION TAG	UNP P16370
C	-16	GLY	-	EXPRESSION TAG	UNP P16370
C	-15	LEU	-	EXPRESSION TAG	UNP P16370

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-14	ASN	-	EXPRESSION TAG	UNP P16370
C	-13	ASP	-	EXPRESSION TAG	UNP P16370
C	-12	ILE	-	EXPRESSION TAG	UNP P16370
C	-11	PHE	-	EXPRESSION TAG	UNP P16370
C	-10	GLU	-	EXPRESSION TAG	UNP P16370
C	-9	ALA	-	EXPRESSION TAG	UNP P16370
C	-8	GLN	-	EXPRESSION TAG	UNP P16370
C	-7	LYS	-	EXPRESSION TAG	UNP P16370
C	-6	ILE	-	EXPRESSION TAG	UNP P16370
C	-5	GLU	-	EXPRESSION TAG	UNP P16370
C	-4	TRP	-	EXPRESSION TAG	UNP P16370
C	-3	HIS	-	EXPRESSION TAG	UNP P16370
C	-2	GLU	-	EXPRESSION TAG	UNP P16370
C	-1	ASP	-	EXPRESSION TAG	UNP P16370
C	0	THR	-	EXPRESSION TAG	UNP P16370
C	1	GLY	-	EXPRESSION TAG	UNP P16370

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	178	Total	C	N	O	S	0	0	0
			1434	887	257	288	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1340	861	222	249	8			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	134	Total	C	N	O	S	0	0	0
			1076	677	182	213	4			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	114	Total	C	N	O	S	0	0	0
			927	571	168	178	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	44	Total	C	N	O	S	0	0	0
			351	217	70	60	4			

- Molecule 13 is a DNA chain called 5'-D(\*AP\*CP\*TP\*AP\*CP\*TP\*TP\*GP\*AP\*GP\*CP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	2	7	Total	C	N	O	P	0	0	0
			137	68	22	41	6			

- Molecule 14 is a DNA chain called 5'-D(\*AP\*GP\*CP\*TP\*C\*AP\*AP\*GP\*TP\*AP\*GP\*TP\*TP\*AP\*TP\*GP\*CP\*CP\*(BRU)P\*GP\*GP\*TP\*CP\*AP\*TP\*T)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
14	1	19	Total	Br	C	N	O	P	0	0	0
			389	1	186	71	113	18			

- Molecule 15 is a RNA chain called 5'-R(\*UP\*GP\*CP\*AP\*UP\*UP\*U\*CP\*AP\*AP\*CP\*CP

\*AP\*GP\*GP\*CP\*UP\*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	3	11	Total	C	N	O	P	0	0	0
			229	104	41	74	10			

- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	J	1	Total	Zn	0	0
			1	1		
16	B	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	C	1	Total	Zn	0	0
			1	1		
16	A	2	Total	Zn	0	0
			2	2		
16	L	1	Total	Zn	0	0
			1	1		

- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	1	Total	Mg	0	0
			1	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase II subunit RPB1

Chain A:














- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F:




- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G:




- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H:



- Molecule 9: DNA-directed RNA polymerase II subunit RPB9


Chain I:



- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5


Chain J:






- Molecule 11: DNA-directed RNA polymerase II subunit RPB11

Chain K:



- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4

Chain L:




- Molecule 13: 5'-D(\*AP\*CP\*TP\*AP\*CP\*TP\*TP\*GP\*AP\*GP\*CP\*T)-3'

Chain 2: 




- Molecule 14: 5'-D(\*AP\*GP\*CP\*TP\*C\*AP\*AP\*GP\*TP\*AP\*GP\*TP\*TP\*AP\*TP\*GP\*CP\*C  
P\*(BRU)P\*GP\*GP\*TP\*CP\*AP\*TP\*T)-3'

Chain 1: 



- Molecule 15: 5'-R(\*UP\*GP\*CP\*AP\*UP\*UP\*U\*CP\*AP\*AP\*CP\*CP\*AP\*GP\*GP\*CP\*UP\*U)-3'

Chain 3: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	222.65Å 395.96Å 283.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.60 48.95 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.60) 99.9 (48.95-3.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 3.57Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.214 , 0.254 0.212 , 0.249	Depositor DCC
$R_{free}$ test set	2843 reflections (2.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	91.4	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 56.7	EDS
Estimated twinning fraction	0.011 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.018 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 144004 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	31876	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/11342	0.80	4/15337 (0.0%)
2	B	0.51	0/9009	0.75	2/12146 (0.0%)
3	C	0.53	0/2133	0.78	1/2891 (0.0%)
4	D	0.48	0/1444	0.76	2/1935 (0.1%)
5	E	0.50	0/1788	0.70	0/2406
6	F	0.60	0/717	0.85	0/967
7	G	0.54	0/1368	0.81	0/1844
8	H	0.48	0/1094	0.77	1/1481 (0.1%)
9	I	0.46	0/945	0.74	0/1273
10	J	0.53	0/541	0.88	0/727
11	K	0.54	0/937	0.72	0/1265
12	L	0.45	0/353	0.77	0/468
13	2	0.68	0/152	0.89	0/232
14	1	0.59	0/413	0.80	0/634
15	3	0.60	0/255	0.89	0/395
All	All	0.52	0/32491	0.78	10/44001 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
14	1	0	3
All	All	0	4

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	26	THR	N-CA-C	-6.09	94.56	111.00
1	A	56	PRO	N-CA-C	-6.08	96.28	112.10
1	A	55	ASP	N-CA-CB	6.07	121.52	110.60
1	A	311	GLN	N-CA-C	5.98	127.14	111.00
2	B	43	LEU	CA-CB-CG	-5.82	101.91	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
14	1	18	DA	Sidechain
14	1	20	DG	Sidechain
14	1	21	DC	Sidechain
3	C	229	TYR	Sidechain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11143	0	11217	1115	0
2	B	8838	0	8870	1052	0
3	C	2095	0	2051	258	0
4	D	1434	0	1460	178	0
5	E	1752	0	1776	175	0
6	F	705	0	731	64	0
7	G	1340	0	1357	142	0
8	H	1076	0	1046	152	0
9	I	927	0	882	122	0
10	J	532	0	542	98	0
11	K	919	0	929	98	0
12	L	351	0	376	59	0
13	2	137	0	82	12	0
14	1	389	0	214	16	0
15	3	229	0	121	12	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	C	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	1	0	0	0	0
All	All	31876	0	31654	3273	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 52.

The worst 5 of 3273 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:882:THR:HG21	2:B:935:ARG:HA	1.24	1.19
1:A:53:LEU:HD23	1:A:54:ASN:N	1.59	1.18
4:D:71:LYS:HA	4:D:74:GLN:HB2	1.23	1.13
7:G:26:LEU:HD12	7:G:56:ILE:HD11	1.22	1.11
1:A:567:LYS:CD	1:A:568:PRO:HD2	1.80	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1406/1733 (81%)	1004 (71%)	279 (20%)	123 (9%)	1	22
2	B	1094/1224 (89%)	774 (71%)	202 (18%)	118 (11%)	1	15
3	C	264/347 (76%)	181 (69%)	48 (18%)	35 (13%)	0	10
4	D	174/221 (79%)	124 (71%)	33 (19%)	17 (10%)	1	18
5	E	212/215 (99%)	144 (68%)	46 (22%)	22 (10%)	1	16
6	F	85/155 (55%)	72 (85%)	8 (9%)	5 (6%)	2	35
7	G	169/171 (99%)	137 (81%)	24 (14%)	8 (5%)	4	42
8	H	130/146 (89%)	84 (65%)	27 (21%)	19 (15%)	0	8

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	I	112/122 (92%)	77 (69%)	26 (23%)	9 (8%)	1	25
10	J	63/70 (90%)	38 (60%)	14 (22%)	11 (18%)	0	4
11	K	112/120 (93%)	86 (77%)	22 (20%)	4 (4%)	5	52
12	L	42/70 (60%)	20 (48%)	13 (31%)	9 (21%)	0	2
All	All	3863/4594 (84%)	2741 (71%)	742 (19%)	380 (10%)	1	18

5 of 380 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	48	ALA
1	A	57	ARG
1	A	58	LEU
1	A	67	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1239/1520 (82%)	1091 (88%)	148 (12%)	8	39
2	B	964/1061 (91%)	846 (88%)	118 (12%)	7	37
3	C	234/299 (78%)	207 (88%)	27 (12%)	8	40
4	D	160/200 (80%)	136 (85%)	24 (15%)	4	27
5	E	196/197 (100%)	176 (90%)	20 (10%)	11	49
6	F	77/137 (56%)	67 (87%)	10 (13%)	6	34
7	G	152/152 (100%)	141 (93%)	11 (7%)	21	68
8	H	118/128 (92%)	99 (84%)	19 (16%)	3	23
9	I	108/116 (93%)	96 (89%)	12 (11%)	9	43
10	J	60/65 (92%)	54 (90%)	6 (10%)	11	50
11	K	99/102 (97%)	86 (87%)	13 (13%)	6	33
12	L	39/57 (68%)	35 (90%)	4 (10%)	10	48
All	All	3446/4034 (85%)	3034 (88%)	412 (12%)	7	38

5 of 412 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	466	TRP
2	B	894	ASP
9	I	34	TYR
2	B	505	ASP
2	B	714	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 104 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	821	GLN
2	B	1084	GLN
8	H	133	ASN
2	B	842	ASN
2	B	957	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
15	3	10/18 (55%)	2 (20%)	1 (10%)

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
15	3	3	A
15	3	11	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
15	3	2	A

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
14	BRU	1	23	15,14	19,21,22	3.80	4 (21%)	22,30,33	1.87	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	1	23	15,14	-	0/5/21/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	1	23	BRU	BR-C5	-15.70	1.50	1.90
14	1	23	BRU	C4-C5	3.53	1.45	1.39
14	1	23	BRU	P-OP1	2.29	1.49	1.46
14	1	23	BRU	C6-N1	2.25	1.39	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	1	23	BRU	C5-C6-N1	6.24	123.78	119.67
14	1	23	BRU	C6-N1-C2	-4.40	121.16	122.41
14	1	23	BRU	BR-C5-C6	2.60	123.13	117.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1416/1733 (81%)	-0.04	6 (0%) 90 76	15, 63, 100, 128	0
2	B	1112/1224 (90%)	0.00	6 (0%) 88 71	15, 74, 111, 122	0
3	C	266/347 (76%)	-0.06	0 100 100	31, 63, 93, 111	0
4	D	178/221 (80%)	0.11	0 100 100	40, 75, 106, 115	0
5	E	214/215 (99%)	0.19	0 100 100	41, 87, 115, 124	0
6	F	87/155 (56%)	-0.28	0 100 100	17, 44, 73, 84	0
7	G	171/171 (100%)	-0.00	0 100 100	44, 64, 92, 100	0
8	H	134/146 (91%)	0.42	0 100 100	67, 95, 112, 117	0
9	I	114/122 (93%)	0.15	0 100 100	62, 93, 109, 114	0
10	J	65/70 (92%)	-0.18	0 100 100	41, 60, 85, 92	0
11	K	114/120 (95%)	0.01	0 100 100	28, 66, 83, 92	0
12	L	44/70 (62%)	0.21	0 100 100	42, 100, 111, 114	0
13	2	7/12 (58%)	1.61	1 (14%) 3 3	112, 116, 123, 124	0
14	1	19/26 (73%)	0.86	5 (26%) 1 2	68, 103, 126, 131	0
15	3	11/18 (61%)	0.26	1 (9%) 9 7	92, 97, 123, 126	0
All	All	3952/4650 (84%)	0.02	19 (0%) 88 71	15, 70, 109, 131	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1455	PRO	3.1
15	3	1	C	3.1
14	1	28	DA	3.0
1	A	149	GLU	2.6
14	1	13	DT	2.5

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
14	BRU	1	23	20/21	0.24	2.69	81,90,95,98	0

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
16	ZN	B	2225	1/1	0.15	-0.56	32,32,32,32	0
16	ZN	A	2457	1/1	0.10	-1.30	35,35,35,35	0
16	ZN	J	1066	1/1	0.17	-1.43	39,39,39,39	0
17	MG	A	2458	1/1	0.12	-1.90	52,52,52,52	0
16	ZN	C	1269	1/1	0.06	-1.95	27,27,27,27	0
16	ZN	I	1121	1/1	0.08	-2.29	74,74,74,74	0
16	ZN	I	1122	1/1	0.10	-2.53	124,124,124,124	0
16	ZN	L	1071	1/1	0.06	-2.65	86,86,86,86	0
16	ZN	A	2456	1/1	0.06	-3.60	70,70,70,70	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.