



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 10:01 PM GMT

PDB ID : 4HON
Title : Crystal structure of human JMJD2D/KDM4D in complex with an H3K9me3 peptide and 2-oxoglutarate
Authors : Krishnan, S.; Trievel, R.C.
Deposited on : 2012-10-22
Resolution : 1.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	330	
1	B	330	
2	F	10	
2	G	10	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
6	SCN	A	404	-	X
6	SCN	A	405	-	X
6	SCN	A	406	-	X
6	SCN	B	405	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
6	SCN	B	406	-	X
7	NO3	A	407	X	X
7	NO3	A	408	X	X
7	NO3	A	409	X	X
7	NO3	A	410	X	X
7	NO3	A	411	X	-
7	NO3	A	412	X	X
7	NO3	A	413	X	X
7	NO3	A	414	X	X
7	NO3	B	407	X	X
7	NO3	B	408	X	X
7	NO3	B	409	X	-
7	NO3	B	410	X	X
8	PDO	A	415	-	X
8	PDO	A	418	-	X
8	PDO	B	411	-	X

2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 5977 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysine-specific demethylase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	4	0
			2672	1719	456	483	14			
1	B	330	Total	C	N	O	S	0	5	0
			2670	1720	455	481	14			

- Molecule 2 is a protein called Histone H3 Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	10	Total	C	N	O	0	1	0
			69	41	14	14			
2	G	7	Total	C	N	O	0	0	0
			43	26	8	9			

- Molecule 3 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: C₅H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ni	0	0
			1	1		
4	A	1	Total	Ni	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	S	0	0
			3	1	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	S	0	0
			3	1	1	1		
6	A	1	Total	C	N	S	0	0
			3	1	1	1		
6	B	1	Total	C	N	S	0	0
			3	1	1	1		
6	B	1	Total	C	N	S	0	0
			3	1	1	1		
6	B	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 7 is NITRATE ION (three-letter code: NO3) (formula: NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	N	O	0	0
			4	1	3		
7	A	1	Total	N	O	0	0
			4	1	3		
7	A	1	Total	N	O	0	0
			4	1	3		
7	A	1	Total	N	O	0	0
			4	1	3		
7	A	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	N	O	0	0
			4	1	3		
7	A	1	Total	N	O	0	0
			4	1	3		
7	B	1	Total	N	O	0	0
			4	1	3		
7	B	1	Total	N	O	0	0
			4	1	3		
7	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 8 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			5	3	2		
8	A	1	Total	C	O	0	0
			5	3	2		
8	A	1	Total	C	O	0	0
			5	3	2		
8	A	1	Total	C	O	0	0
			5	3	2		
8	B	1	Total	C	O	0	0
			5	3	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	232	Total 232	O 232	0	0
9	B	172	Total 172	O 172	0	0
9	F	4	Total 4	O 4	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific demethylase 4D

Chain A: 



- Molecule 1: Lysine-specific demethylase 4D

Chain B: 




- Molecule 2: Histone H3 Peptide

Chain F: 



- Molecule 2: Histone H3 Peptide

Chain G: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.20Å 79.67Å 175.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.93 – 1.80 24.93 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.9 (24.93-1.80) 93.9 (24.93-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.189 , 0.214 0.190 , 0.214	Depositor DCC
R_{free} test set	4486 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 89102 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5977	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI, ZN, SCN, PDO, M3L, AKG, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2769	0.70	0/3762
1	B	0.77	1/2767 (0.0%)	0.72	2/3756 (0.1%)
2	F	0.72	0/59	0.73	0/78
2	G	0.88	0/30	0.58	0/39
All	All	0.76	1/5625 (0.0%)	0.71	2/7635 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218	GLU	CD-OE1	-5.91	1.19	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	B	102	ARG	NE-CZ-NH2	-6.42	117.09	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2672	0	2524	9	0
1	B	2670	0	2532	8	0
2	F	69	0	73	4	0
2	G	43	0	43	2	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	9	0	0	0	0
6	B	9	0	0	0	0
7	A	32	0	0	0	0
7	B	16	0	0	0	0
8	A	20	0	32	10	0
8	B	5	0	8	0	0
9	A	232	0	0	5	0
9	B	172	0	0	2	0
9	F	4	0	0	0	0
All	All	5977	0	5220	22	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (22) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:45:ARG:HG2	9:A:679:HOH:O	1.50	1.12
8:A:418:PDO:H11	1:B:22:HIS:H	1.31	0.95
8:A:418:PDO:H11	1:B:22:HIS:N	1.93	0.84
8:A:418:PDO:C1	1:B:22:HIS:H	2.00	0.74
1:A:20:ILE:H	8:A:418:PDO:H31	1.60	0.65
1:A:192:HIS:CE1	8:A:415:PDO:H22	2.38	0.58
1:A:89[B]:TYR:CD1	2:F:15:ALA:O	2.56	0.58
1:B:234:SER:HB2	1:B:241[A]:PHE:CG	2.41	0.56
9:B:508:HOH:O	2:G:9:M3L:HM12	2.06	0.55
1:A:173:GLU:HG3	9:A:657:HOH:O	2.06	0.55
8:A:415:PDO:H11	9:A:537:HOH:O	2.10	0.52
1:A:256:GLU:HG3	9:A:728:HOH:O	2.11	0.51
1:B:24:THR:OG1	1:B:27[B]:GLU:HG2	2.12	0.49
1:B:241[A]:PHE:HD2	9:B:599:HOH:O	1.95	0.49
1:B:234:SER:HB2	1:B:241[B]:PHE:CG	2.48	0.49
8:A:415:PDO:H31	2:F:11:THR:HA	1.97	0.47
9:A:507:HOH:O	2:F:9:M3L:HM12	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:102:ARG:HD2	1:A:287:PHE:CE2	2.52	0.44
1:A:20:ILE:N	8:A:418:PDO:H31	2.30	0.44
8:A:415:PDO:C3	2:F:11:THR:HA	2.48	0.43
1:B:245:LYS:HB2	2:G:12:GLY:HA2	2.02	0.41
1:A:20:ILE:H	8:A:418:PDO:C3	2.29	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/330 (101%)	330 (99%)	3 (1%)	0	100	100
1	B	333/330 (101%)	330 (99%)	3 (1%)	0	100	100
2	F	8/10 (80%)	8 (100%)	0	0	100	100
2	G	4/10 (40%)	4 (100%)	0	0	100	100
All	All	678/680 (100%)	672 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/282 (97%)	270 (99%)	3 (1%)	84	77
1	B	274/282 (97%)	270 (98%)	4 (2%)	76	66
2	F	5/5 (100%)	5 (100%)	0	100	100
2	G	2/5 (40%)	2 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	554/574 (96%)	547 (99%)	7 (1%)	80	71

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	PHE
1	A	136	TYR
1	A	179	TYR
1	B	28	PHE
1	B	136	TYR
1	B	179	TYR
1	B	311	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	156	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	M3L	F	9	2	11,11,12	5.52	1 (9%)	12,14,16	2.07	1 (8%)
2	M3L	G	9	2	11,11,12	6.34	1 (9%)	12,14,16	1.65	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M3L	F	9	2	-	0/8/10/12	0/0/0/0
2	M3L	G	9	2	-	0/8/10/12	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	9	M3L	O-C	20.88	1.25	1.11
2	F	9	M3L	O-C	18.11	1.23	1.11

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	9	M3L	C-CA-N	-6.40	107.44	113.83
2	G	9	M3L	C-CA-N	-5.13	108.71	113.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 29 ligands modelled in this entry, 4 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AKG	A	401	4	9,9,9	1.05	0	11,11,11	1.24	1 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SCN	A	404	-	2,2,2	2.02	1 (50%)	1,1,1	0.38	0
6	SCN	A	405	-	2,2,2	1.99	1 (50%)	1,1,1	0.07	0
6	SCN	A	406	-	2,2,2	2.03	1 (50%)	1,1,1	0.52	0
7	NO3	A	407	-	3,3,3	3.02	3 (100%)	3,3,3	0.07	0
7	NO3	A	408	-	3,3,3	3.13	3 (100%)	3,3,3	0.11	0
7	NO3	A	409	-	3,3,3	3.09	3 (100%)	3,3,3	0.14	0
7	NO3	A	410	-	3,3,3	3.16	3 (100%)	3,3,3	0.16	0
7	NO3	A	411	-	3,3,3	3.20	3 (100%)	3,3,3	0.39	0
7	NO3	A	412	-	3,3,3	2.94	3 (100%)	3,3,3	0.30	0
7	NO3	A	413	-	3,3,3	3.23	3 (100%)	3,3,3	0.11	0
7	NO3	A	414	-	3,3,3	3.03	3 (100%)	3,3,3	0.52	0
8	PDO	A	415	-	4,4,4	0.41	0	3,3,3	0.32	0
8	PDO	A	416	-	4,4,4	0.40	0	3,3,3	0.51	0
8	PDO	A	417	-	4,4,4	0.43	0	3,3,3	0.29	0
8	PDO	A	418	-	4,4,4	0.30	0	3,3,3	0.49	0
3	AKG	B	401	4	9,9,9	1.14	0	11,11,11	1.09	1 (9%)
6	SCN	B	404	-	2,2,2	1.77	1 (50%)	1,1,1	0.99	0
6	SCN	B	405	-	2,2,2	1.97	1 (50%)	1,1,1	0.43	0
6	SCN	B	406	-	2,2,2	1.63	1 (50%)	1,1,1	0.35	0
7	NO3	B	407	-	3,3,3	3.08	3 (100%)	3,3,3	0.09	0
7	NO3	B	408	-	3,3,3	3.06	3 (100%)	3,3,3	0.35	0
7	NO3	B	409	-	3,3,3	3.03	3 (100%)	3,3,3	0.32	0
7	NO3	B	410	-	3,3,3	2.82	3 (100%)	3,3,3	0.33	0
8	PDO	B	411	-	4,4,4	0.37	0	3,3,3	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AKG	A	401	4	-	0/9/9/9	0/0/0/0
6	SCN	A	404	-	-	0/0/0/0	0/0/0/0
6	SCN	A	405	-	-	0/0/0/0	0/0/0/0
6	SCN	A	406	-	-	0/0/0/0	0/0/0/0
7	NO3	A	407	-	-	0/0/0/0	0/0/0/0
7	NO3	A	408	-	-	0/0/0/0	0/0/0/0
7	NO3	A	409	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NO3	A	410	-	-	0/0/0/0	0/0/0/0
7	NO3	A	411	-	-	0/0/0/0	0/0/0/0
7	NO3	A	412	-	-	0/0/0/0	0/0/0/0
7	NO3	A	413	-	-	0/0/0/0	0/0/0/0
7	NO3	A	414	-	-	0/0/0/0	0/0/0/0
8	PDO	A	415	-	-	0/2/2/2	0/0/0/0
8	PDO	A	416	-	-	0/2/2/2	0/0/0/0
8	PDO	A	417	-	-	0/2/2/2	0/0/0/0
8	PDO	A	418	-	-	0/2/2/2	0/0/0/0
3	AKG	B	401	4	-	0/9/9/9	0/0/0/0
6	SCN	B	404	-	-	0/0/0/0	0/0/0/0
6	SCN	B	405	-	-	0/0/0/0	0/0/0/0
6	SCN	B	406	-	-	0/0/0/0	0/0/0/0
7	NO3	B	407	-	-	0/0/0/0	0/0/0/0
7	NO3	B	408	-	-	0/0/0/0	0/0/0/0
7	NO3	B	409	-	-	0/0/0/0	0/0/0/0
7	NO3	B	410	-	-	0/0/0/0	0/0/0/0
8	PDO	B	411	-	-	0/2/2/2	0/0/0/0

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	408	NO3	O1-N	3.91	1.40	1.24
7	A	409	NO3	O1-N	3.89	1.40	1.24
7	B	407	NO3	O1-N	3.86	1.40	1.24
7	A	413	NO3	O1-N	3.82	1.40	1.24
7	A	408	NO3	O1-N	3.79	1.40	1.24
7	A	411	NO3	O1-N	3.77	1.40	1.24
7	A	410	NO3	O1-N	3.76	1.40	1.24
7	A	412	NO3	O1-N	3.69	1.39	1.24
7	B	409	NO3	O1-N	3.63	1.39	1.24
7	A	407	NO3	O1-N	3.52	1.39	1.24
7	A	414	NO3	O1-N	3.42	1.38	1.24
7	B	410	NO3	O1-N	3.35	1.38	1.24
7	A	414	NO3	O3-N	3.03	1.41	1.25
7	A	411	NO3	O3-N	3.00	1.41	1.25
7	A	413	NO3	O3-N	2.94	1.40	1.25
7	A	410	NO3	O2-N	2.89	1.40	1.25
7	A	413	NO3	O2-N	2.86	1.40	1.25
7	A	407	NO3	O3-N	2.85	1.40	1.25
6	A	406	SCN	C-S	2.84	1.82	1.63
7	B	409	NO3	O2-N	2.81	1.40	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	404	SCN	C-S	2.81	1.81	1.63
7	A	408	NO3	O2-N	2.80	1.40	1.25
7	A	411	NO3	O2-N	2.75	1.39	1.25
7	A	410	NO3	O3-N	2.75	1.39	1.25
6	B	405	SCN	C-S	2.73	1.81	1.63
6	A	405	SCN	C-S	2.70	1.81	1.63
7	A	408	NO3	O3-N	2.67	1.39	1.25
7	B	410	NO3	O3-N	2.67	1.39	1.25
7	A	409	NO3	O3-N	2.66	1.39	1.25
7	B	407	NO3	O3-N	2.65	1.39	1.25
7	A	407	NO3	O2-N	2.61	1.39	1.25
7	A	414	NO3	O2-N	2.59	1.39	1.25
7	A	412	NO3	O3-N	2.58	1.39	1.25
7	B	408	NO3	O3-N	2.56	1.38	1.25
7	B	407	NO3	O2-N	2.56	1.38	1.25
7	A	409	NO3	O2-N	2.55	1.38	1.25
7	B	409	NO3	O3-N	2.52	1.38	1.25
7	B	408	NO3	O2-N	2.50	1.38	1.25
6	B	404	SCN	C-S	2.45	1.79	1.63
7	A	412	NO3	O2-N	2.37	1.37	1.25
7	B	410	NO3	O2-N	2.35	1.37	1.25
6	B	406	SCN	C-S	2.24	1.78	1.63

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	AKG	C3-C4-C5	-2.77	108.40	113.53
3	A	401	AKG	C3-C4-C5	-2.39	109.10	113.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/330 (100%)	-0.06	3 (0%) 81 78	10, 18, 30, 42	0
1	B	330/330 (100%)	0.01	11 (3%) 44 36	10, 19, 33, 44	0
2	F	10/10 (100%)	2.18	3 (30%) 1 1	20, 31, 44, 48	0
2	G	7/10 (70%)	1.89	3 (42%) 1 1	26, 33, 41, 43	0
All	All	677/680 (99%)	0.03	20 (2%) 43 40	10, 19, 33, 48	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	15	ALA	8.1
2	F	6	THR	4.3
1	A	65	ASN	4.2
2	G	7	ALA	4.2
1	B	311	SER	4.0
1	B	235	SER	3.3
2	F	7	ALA	3.3
1	B	236	ARG	2.8
1	B	315	ALA	2.6
1	A	14	PRO	2.4
1	B	14	PRO	2.3
1	A	341	ASP	2.3
2	G	13	GLY	2.2
2	G	11	THR	2.2
1	B	234	SER	2.2
1	B	314	GLU	2.1
1	B	65	ASN	2.1
1	B	317	VAL	2.1
1	B	111	GLN	2.1
1	B	233	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	M3L	G	9	12/13	0.13	0.11	20,24,28,29	0
2	M3L	F	9	12/13	0.11	-0.03	17,19,21,21	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	NO3	A	413	4/4	0.33	25.21	55,55,55,56	0
7	NO3	A	409	4/4	0.23	11.01	38,39,40,40	0
8	PDO	A	418	5/5	0.17	10.49	26,30,33,35	0
6	SCN	A	404	3/3	0.44	8.27	71,71,72,72	0
7	NO3	A	412	4/4	0.22	8.14	31,32,32,35	0
7	NO3	B	410	4/4	0.18	8.12	32,32,33,34	0
7	NO3	A	410	4/4	0.20	6.54	39,40,40,41	0
7	NO3	A	414	4/4	0.30	6.36	45,45,45,47	0
7	NO3	A	408	4/4	0.17	5.68	37,37,37,39	0
8	PDO	B	411	5/5	0.12	4.99	35,40,42,43	0
6	SCN	A	406	3/3	0.28	4.78	47,47,48,48	0
7	NO3	B	408	4/4	0.18	4.14	23,24,25,27	0
8	PDO	A	415	5/5	0.17	3.32	32,33,33,36	0
6	SCN	A	405	3/3	0.19	3.15	46,46,47,48	0
6	SCN	B	406	3/3	0.23	3.09	49,49,50,50	0
7	NO3	A	407	4/4	0.13	2.65	26,27,28,29	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	SCN	B	405	3/3	0.23	2.62	55,55,56,56	0
7	NO3	B	407	4/4	0.17	2.23	29,32,32,33	0
8	PDO	A	417	5/5	0.14	1.81	37,38,39,39	0
7	NO3	A	411	4/4	0.28	1.59	44,44,45,45	0
7	NO3	B	409	4/4	0.11	1.33	24,26,27,29	0
8	PDO	A	416	5/5	0.12	0.98	20,27,32,35	0
6	SCN	B	404	3/3	0.14	0.36	33,33,33,34	0
3	AKG	B	401	10/10	0.09	-1.08	13,18,21,22	0
3	AKG	A	401	10/10	0.10	-1.28	13,15,19,20	0
5	ZN	A	403	1/1	0.04	-1.48	20,20,20,20	0
5	ZN	B	403	1/1	0.05	-1.51	26,26,26,26	0
4	NI	A	402	1/1	0.06	-5.86	13,13,13,13	0
4	NI	B	402	1/1	0.04	-6.76	14,14,14,14	0

6.5 Other polymers ⓘ

There are no such residues in this entry.