



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 10:16 AM GMT

PDB ID : 3HQ7
Title : CcpA from *G. sulfurreducens*, G94K/K97Q/R100I variant
Authors : Hoffmann, M.; Seidel, J.; Einsle, O.
Deposited on : 2009-06-05
Resolution : 2.31 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3293 (2.34-2.30)
Clashscore	79885	4097 (2.34-2.30)
Ramachandran outliers	78287	4055 (2.34-2.30)
Sidechain outliers	78261	4054 (2.34-2.30)
RSRZ outliers	66119	3294 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	345	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2557 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c551 peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	0	0
			2287	1465	388	425	9			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	LYS	GLY	ENGINEERED	UNP Q749D0
A	97	GLN	LYS	ENGINEERED	UNP Q749D0
A	100	ILE	ARG	ENGINEERED	UNP Q749D0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca		
			1	1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	183	Total	O		
			183	183	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome c551 peroxidase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	99.30Å 99.35Å 78.22Å 89.97° 89.98° 90.00°	Depositor
Resolution (Å)	70.00 – 2.31 49.65 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.8 (70.00-2.31) 99.7 (49.65-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.209 , 0.294 0.209 , 0.291	Depositor DCC
R_{free} test set	905 reflections (5.09%)	DCC
Wilson B-factor (Å ²)	36.2	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 17776 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2557	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.49	5/2341 (0.2%)	1.29	13/3177 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	154	TYR	CE1-CZ	6.24	1.46	1.38
1	A	100	ILE	CA-CB	5.61	1.67	1.54
1	A	61	PHE	CE2-CZ	5.42	1.47	1.37
1	A	95	TRP	CB-CG	5.41	1.59	1.50
1	A	171	PHE	CE1-CZ	5.05	1.47	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	342	ARG	NE-CZ-NH2	-12.45	114.08	120.30
1	A	342	ARG	NE-CZ-NH1	11.66	126.13	120.30
1	A	172	ASP	CB-CG-OD1	-8.94	110.25	118.30
1	A	28	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	A	297	MET	CG-SD-CE	-6.37	90.01	100.20
1	A	228	LEU	CB-CG-CD2	6.03	121.24	111.00
1	A	32	LEU	CA-CB-CG	5.95	128.98	115.30
1	A	332	ILE	CA-CB-CG1	-5.53	100.49	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	ASP	CB-CG-OD1	5.52	123.27	118.30
1	A	344	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	345	LEU	CA-CB-CG	5.11	127.05	115.30
1	A	338	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	344	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	218	GLY	Peptide
1	A	230	GLY	Peptide
1	A	311	SER	Peptide
1	A	344	ARG	Peptide
1	A	45	GLY	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2287	0	2298	85	0
2	A	86	0	60	10	0
3	A	1	0	0	1	0
4	A	183	0	0	14	0
All	All	2557	0	2358	87	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

All (87) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:76:CYS:SG	2:A:401:HEM:CAC	2.27	1.22
1:A:193:ASP:O	1:A:197:LYS:HG3	1.45	1.17
1:A:76:CYS:SG	2:A:401:HEM:CBC	2.50	0.99
1:A:214:PHE:O	1:A:219:CYS:HB2	1.62	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:146:LYS:HD3	1:A:341:PRO:HG3	1.56	0.88
1:A:216:ASP:OD1	4:A:496:HOH:O	1.92	0.87
1:A:220:VAL:HG22	1:A:220:VAL:O	1.75	0.83
1:A:285:GLY:O	4:A:470:HOH:O	1.99	0.80
1:A:113:ALA:HB1	1:A:268:ARG:HD3	1.65	0.79
1:A:302:PHE:CE1	2:A:402:HEM:HBC2	2.20	0.76
1:A:76:CYS:SG	2:A:401:HEM:C3C	2.79	0.75
1:A:233:TYR:CE2	1:A:268:ARG:HB2	2.23	0.72
1:A:107:ASN:HD21	1:A:187:THR:H	1.38	0.72
1:A:192:PHE:CZ	1:A:196:LEU:HD21	2.26	0.71
1:A:22:ALA:O	1:A:24:GLU:HB3	1.91	0.70
1:A:28:ARG:NH2	4:A:496:HOH:O	1.99	0.67
1:A:113:ALA:CB	1:A:268:ARG:HD3	2.25	0.66
1:A:26:GLN:NE2	1:A:198:GLY:O	2.30	0.65
1:A:71:ILE:HD11	4:A:387:HOH:O	1.96	0.64
3:A:403:CA:CA	4:A:468:HOH:O	1.72	0.64
1:A:90:SER:HB2	2:A:401:HEM:HBC2	1.79	0.64
1:A:300:ALA:O	1:A:302:PHE:N	2.33	0.62
1:A:213:LEU:O	1:A:217:LYS:HG3	2.01	0.61
1:A:28:ARG:HG3	1:A:32:LEU:HD11	1.83	0.60
1:A:46:ASN:N	1:A:47:PRO:CD	2.65	0.59
1:A:220:VAL:O	1:A:220:VAL:CG2	2.49	0.59
1:A:113:ALA:CB	1:A:268:ARG:CD	2.81	0.58
1:A:22:ALA:HB1	1:A:200:LYS:HD3	1.86	0.58
1:A:256:ALA:HA	1:A:296:VAL:HG13	1.84	0.58
1:A:69:HIS:HD2	4:A:383:HOH:O	1.86	0.58
1:A:186:ILE:HG13	1:A:188:PRO:HD3	1.86	0.58
1:A:195:TYR:HA	1:A:199:LYS:O	2.05	0.56
1:A:28:ARG:NH1	4:A:496:HOH:O	2.39	0.56
1:A:217:LYS:O	1:A:304:ILE:HD11	2.06	0.56
1:A:132:GLN:O	1:A:136:GLU:HG3	2.07	0.55
1:A:94:LYS:HG2	1:A:137:MET:HG2	1.89	0.54
1:A:241:LYS:HB2	4:A:531:HOH:O	2.08	0.54
1:A:113:ALA:HB2	1:A:268:ARG:HD2	1.89	0.53
1:A:113:ALA:HB2	1:A:268:ARG:CD	2.39	0.53
1:A:298:GLY:HA2	1:A:304:ILE:CG2	2.39	0.53
1:A:140:THR:OG1	1:A:142:ASP:HB3	2.09	0.52
1:A:28:ARG:HG3	1:A:32:LEU:CD1	2.40	0.52
1:A:204:ASP:C	1:A:204:ASP:OD1	2.48	0.51
1:A:231:THR:HG22	4:A:356:HOH:O	2.10	0.51
1:A:234:PHE:HB3	1:A:236:PHE:HD1	1.76	0.50
1:A:146:LYS:HD3	1:A:341:PRO:CG	2.35	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:113:ALA:HA	1:A:233:TYR:OH	2.12	0.50
1:A:211:LEU:O	1:A:214:PHE:HB3	2.13	0.48
1:A:189:ASP:N	1:A:189:ASP:OD1	2.40	0.48
1:A:28:ARG:O	1:A:31:GLY:N	2.40	0.48
1:A:26:GLN:HG3	1:A:195:TYR:CZ	2.49	0.48
1:A:231:THR:HG21	4:A:358:HOH:O	2.13	0.48
1:A:33:PHE:CD2	1:A:228:LEU:HD22	2.49	0.47
1:A:41:PRO:O	4:A:422:HOH:O	2.20	0.47
1:A:223:HIS:HE1	2:A:402:HEM:ND	2.01	0.47
1:A:272:LEU:O	1:A:275:VAL:HG12	2.14	0.47
1:A:203:LEU:HA	1:A:207:GLN:HE21	1.78	0.47
1:A:215:LEU:O	1:A:218:GLY:N	2.28	0.47
1:A:298:GLY:HA2	1:A:304:ILE:HG22	1.95	0.47
1:A:46:ASN:N	1:A:47:PRO:HD3	2.31	0.46
1:A:217:LYS:NZ	1:A:305:LYS:O	2.37	0.46
1:A:113:ALA:HB1	1:A:268:ARG:CD	2.39	0.46
1:A:25:LEU:O	1:A:28:ARG:HB3	2.16	0.45
1:A:88:ALA:HB3	4:A:407:HOH:O	2.16	0.45
1:A:306:LEU:HD23	1:A:306:LEU:HA	1.78	0.45
1:A:301:GLN:H	1:A:301:GLN:HG2	1.66	0.45
1:A:143:GLN:HE22	1:A:341:PRO:HB3	1.81	0.45
1:A:90:SER:HB2	2:A:401:HEM:CBC	2.46	0.44
1:A:135:VAL:HG11	1:A:171:PHE:CE1	2.52	0.44
1:A:215:LEU:O	1:A:216:ASP:C	2.56	0.44
1:A:91:THR:HA	1:A:96:GLN:HE22	1.82	0.44
1:A:188:PRO:O	1:A:324:LYS:HG2	2.18	0.44
1:A:71:ILE:CD1	4:A:387:HOH:O	2.63	0.43
1:A:92:GLY:HA3	2:A:401:HEM:C3C	2.53	0.43
1:A:282:PHE:O	1:A:284:SER:N	2.52	0.43
1:A:233:TYR:HB3	1:A:266:VAL:HB	2.02	0.42
2:A:401:HEM:HAC	2:A:401:HEM:HMC1	1.85	0.41
1:A:188:PRO:O	1:A:324:LYS:CG	2.67	0.41
1:A:257:VAL:HG12	1:A:258:THR:N	2.35	0.41
1:A:241:LYS:HE3	1:A:241:LYS:HB3	1.38	0.41
1:A:214:PHE:CE1	1:A:219:CYS:HB3	2.56	0.41
1:A:95:TRP:CD1	1:A:95:TRP:C	2.95	0.41
1:A:236:PHE:CD1	2:A:402:HEM:HBC1	2.56	0.41
1:A:34:LYS:HB3	1:A:34:LYS:HE2	1.82	0.41
1:A:143:GLN:HG3	4:A:491:HOH:O	2.21	0.40
1:A:307:SER:N	1:A:310:GLU:OE1	2.36	0.40
1:A:307:SER:OG	1:A:310:GLU:HG3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/345 (86%)	259 (87%)	25 (8%)	14 (5%)	4 1

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	219	CYS
1	A	301	GLN
1	A	22	ALA
1	A	23	ASP
1	A	24	GLU
1	A	30	GLN
1	A	240	GLU
1	A	283	HIS
1	A	209	ALA
1	A	210	GLY
1	A	215	LEU
1	A	312	GLU
1	A	231	THR
1	A	220	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	244/277 (88%)	217 (89%)	27 (11%)	9 9

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ASP
1	A	24	GLU
1	A	25	LEU
1	A	39	LYS
1	A	94	LYS
1	A	95	TRP
1	A	128	LYS
1	A	137	MET
1	A	154	TYR
1	A	186	ILE
1	A	189	ASP
1	A	196	LEU
1	A	199	LYS
1	A	200	LYS
1	A	212	LYS
1	A	228	LEU
1	A	241	LYS
1	A	262	LYS
1	A	265	TYR
1	A	286	VAL
1	A	291	LYS
1	A	297	MET
1	A	308	ASP
1	A	309	ASP
1	A	312	GLU
1	A	324	LYS
1	A	345	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	HIS
1	A	107	ASN
1	A	143	GLN
1	A	194	GLN
1	A	207	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	401	1	49,50,50	2.96	20 (40%)	46,82,82	2.99	15 (32%)
2	HEM	A	402	1	49,50,50	2.99	20 (40%)	46,82,82	3.16	16 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	401	1	-	0/14/114/114	0/0/8/8
2	HEM	A	402	1	-	0/14/114/114	0/0/8/8

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	HEM	C3D-C4D	-9.77	1.42	1.44
2	A	402	HEM	FE-ND	8.49	2.28	1.97
2	A	402	HEM	FE-NA	7.12	2.23	1.92
2	A	402	HEM	C3D-C2D	6.45	1.55	1.43
2	A	401	HEM	C3C-C2C	-6.03	1.33	1.43
2	A	402	HEM	FE-NC	5.91	2.20	1.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	HEM	C3C-CAC	5.65	1.58	1.40
2	A	402	HEM	C3B-CAB	5.50	1.57	1.40
2	A	401	HEM	C3B-CAB	5.43	1.57	1.40
2	A	402	HEM	C3C-C2C	-5.20	1.34	1.43
2	A	401	HEM	C4A-C3A	5.04	1.46	1.40
2	A	402	HEM	C2B-C1B	-5.04	1.43	1.44
2	A	401	HEM	C3C-CAC	4.81	1.55	1.40
2	A	401	HEM	C3B-C2B	-4.69	1.35	1.43
2	A	401	HEM	C4C-NC	4.66	1.44	1.38
2	A	401	HEM	C3D-C2D	4.35	1.51	1.43
2	A	401	HEM	C2D-C1D	4.33	1.45	1.44
2	A	402	HEM	CHA-C4D	4.25	1.41	1.35
2	A	401	HEM	FE-ND	4.11	2.12	1.97
2	A	401	HEM	FE-NB	3.91	2.12	1.97
2	A	402	HEM	C3B-C2B	-3.71	1.37	1.43
2	A	401	HEM	CAA-C2A	3.56	1.58	1.52
2	A	401	HEM	CMC-C2C	3.38	1.57	1.47
2	A	402	HEM	CHD-C4C	3.27	1.42	1.36
2	A	401	HEM	CHA-C4D	3.21	1.40	1.35
2	A	401	HEM	CMA-C3A	3.15	1.58	1.51
2	A	401	HEM	CHD-C4C	3.12	1.42	1.36
2	A	402	HEM	CHB-C1B	3.07	1.40	1.35
2	A	402	HEM	CMD-C2D	2.99	1.56	1.47
2	A	402	HEM	CMC-C2C	2.91	1.56	1.47
2	A	401	HEM	C1A-NA	2.86	1.42	1.36
2	A	401	HEM	CMB-C2B	2.45	1.55	1.47
2	A	402	HEM	CMB-C2B	2.45	1.55	1.47
2	A	401	HEM	CMD-C2D	2.37	1.54	1.47
2	A	402	HEM	C1A-C2A	2.22	1.47	1.43
2	A	402	HEM	C1D-ND	2.15	1.43	1.37
2	A	402	HEM	CAA-C2A	2.10	1.55	1.52
2	A	401	HEM	O1D-CGD	2.08	1.29	1.22
2	A	402	HEM	FE-NB	2.07	2.05	1.97
2	A	402	HEM	CMA-C3A	2.03	1.55	1.51

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C3B-C4B-NB	-10.90	106.20	114.00
2	A	402	HEM	C4D-ND-C1D	10.12	115.52	105.16
2	A	402	HEM	C3B-C4B-NB	-9.92	106.90	114.00
2	A	401	HEM	CHC-C1C-NC	7.41	131.17	124.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	HEM	C2D-C1D-ND	-6.02	105.82	112.93
2	A	401	HEM	C4A-CHB-C1B	-5.98	119.60	127.47
2	A	401	HEM	C4D-ND-C1D	5.71	111.00	105.16
2	A	402	HEM	C2A-C1A-NA	-5.35	102.31	109.73
2	A	402	HEM	CMA-C3A-C4A	-4.79	121.25	128.62
2	A	402	HEM	C4A-NA-C1A	4.63	112.86	106.76
2	A	401	HEM	CMA-C3A-C4A	-4.39	121.87	128.62
2	A	402	HEM	C4C-NC-C1C	4.27	109.97	105.53
2	A	401	HEM	CHA-C4D-ND	4.18	130.05	124.31
2	A	401	HEM	C1B-NB-C4B	4.16	109.42	105.16
2	A	402	HEM	C1B-NB-C4B	4.16	109.42	105.16
2	A	401	HEM	C2D-C1D-ND	-4.15	108.03	112.93
2	A	401	HEM	C1D-CHD-C4C	-3.94	116.20	126.57
2	A	402	HEM	CMA-C3A-C2A	3.71	131.93	124.94
2	A	402	HEM	O1A-CGA-CBA	-3.57	110.74	123.03
2	A	402	HEM	C1A-C2A-C3A	3.34	110.38	106.92
2	A	401	HEM	CBA-CAA-C2A	-3.33	106.82	112.69
2	A	401	HEM	CAA-CBA-CGA	3.20	123.75	113.47
2	A	402	HEM	C2A-C1A-CHA	3.03	131.75	126.00
2	A	402	HEM	CHC-C1C-NC	2.99	127.33	124.73
2	A	402	HEM	CHB-C1B-NB	2.95	128.36	124.31
2	A	401	HEM	O2A-CGA-CBA	2.57	123.28	114.22
2	A	401	HEM	C4A-C3A-C2A	2.55	108.77	107.00
2	A	401	HEM	O1A-CGA-CBA	-2.53	114.34	123.03
2	A	402	HEM	C3A-C4A-NA	-2.35	107.64	109.41
2	A	401	HEM	CMA-C3A-C2A	2.33	129.34	124.94
2	A	402	HEM	O2A-CGA-CBA	2.19	121.94	114.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/345 (88%)	0.19	14 (4%) 31 41	18, 37, 66, 80	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	265	TYR	5.9
1	A	242	PRO	5.3
1	A	127	ALA	4.5
1	A	113	ALA	4.1
1	A	253	GLY	4.0
1	A	27	GLN	3.0
1	A	302	PHE	2.8
1	A	198	GLY	2.8
1	A	195	TYR	2.4
1	A	97	GLN	2.4
1	A	303	GLY	2.2
1	A	30	GLN	2.2
1	A	262	LYS	2.1
1	A	26	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CA	A	403	1/1	0.18	0.81	19,19,19,19	0
2	HEM	A	401	43/43	0.15	0.19	21,27,49,53	0
2	HEM	A	402	43/43	0.14	-0.02	24,48,53,55	0

6.5 Other polymers

There are no such residues in this entry.