



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 04:22 AM GMT

PDB ID : 3HQ9
Title : CcpA from *G. sulfurreducens*, S134P variant
Authors : Hoffmann, M.; Seidel, J.; Einsle, O.
Deposited on : 2009-06-05
Resolution : 1.52 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1879 (1.54-1.50)
Clashscore	79885	2184 (1.54-1.50)
Ramachandran outliers	78287	2123 (1.54-1.50)
Sidechain outliers	78261	2121 (1.54-1.50)
RSRZ outliers	66119	1880 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	345	
1	B	345	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	IMD	A	346	-	X
6	BU3	A	350	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5448 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c551 peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	0	2	0
			2364	1513	403	439	9			
1	B	306	Total	C	N	O	S	0	3	0
			2313	1480	393	431	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	PRO	SER	ENGINEERED	UNP Q749D0
B	134	PRO	SER	ENGINEERED	UNP Q749D0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	B	1	Total	C	N	0	0
			5	3	2		
4	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is OXALATE ION (three-letter code: OXL) (formula: C_2O_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	2	4		
5	A	1	Total	C	O	0	0
			6	2	4		
5	B	1	Total	C	O	0	0
			6	2	4		
5	B	1	Total	C	O	0	0
			6	2	4		

- Molecule 6 is (R,R)-2,3-BUTANEDIOL (three-letter code: BU3) (formula: $C_4H_{10}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	4	2		
6	B	1	Total	C	O	0	0
			6	4	2		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	244	Total	O	0	0
			244	244		
7	B	297	Total	O	0	0
			297	297		

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome c551 peroxidase

Chain A:



- Molecule 1: Cytochrome c551 peroxidase

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.90Å 55.47Å 68.61Å 104.71° 99.39° 108.97°	Depositor
Resolution (Å)	63.92 – 1.52 26.65 – 1.52	Depositor EDS
% Data completeness (in resolution range)	90.1 (63.92-1.52) 90.2 (26.65-1.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.160 , 0.184 0.169 , 0.190	Depositor DCC
R_{free} test set	4561 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 46.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 91066 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5448	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OXL, BU3, CA, IMD, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2428	0.55	0/3297
1	B	0.46	0/2378	0.59	0/3232
All	All	0.44	0/4806	0.57	0/6529

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2364	0	2359	27	1
1	B	2313	0	2284	16	1
2	A	86	0	60	15	0
2	B	86	0	60	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	9	3	0
4	B	10	0	9	0	0
5	A	12	0	0	0	0
5	B	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	6	0	10	0	0
6	B	6	0	10	2	0
7	A	244	0	0	3	0
7	B	297	0	0	5	0
All	All	5448	0	4801	43	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (43) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:76:CYS:SG	2:A:400:HEM:CAC	2.05	1.45
1:A:219:CYS:SG	2:A:401:HEM:CAB	2.06	1.42
1:B:222:CYS:SG	2:B:401:HEM:CAC	2.15	1.34
1:A:222:CYS:SG	2:A:401:HEM:CAC	2.24	1.25
1:A:219:CYS:SG	2:A:401:HEM:HAB	1.71	1.23
1:A:76:CYS:SG	2:A:400:HEM:HAC	1.78	1.16
1:A:131:VAL:HG13	7:A:494:HOH:O	1.43	1.14
1:A:222:CYS:SG	2:A:401:HEM:HAC	1.97	1.04
1:B:222:CYS:SG	2:B:401:HEM:HAC	1.98	1.02
1:A:222:CYS:HG	2:A:401:HEM:CAC	1.71	0.96
1:B:259[B]:ASN:OD1	7:B:379:HOH:O	1.94	0.85
1:A:76:CYS:SG	2:A:400:HEM:C3C	2.77	0.78
6:B:350:BU3:H42	7:B:572:HOH:O	1.89	0.72
1:A:76:CYS:SG	2:A:400:HEM:CBC	2.77	0.71
1:B:222:CYS:SG	2:B:401:HEM:C3C	2.83	0.71
1:B:222:CYS:SG	2:B:401:HEM:CBC	2.80	0.70
1:A:219:CYS:SG	2:A:401:HEM:CBB	2.80	0.69
1:B:301:GLN:OE1	7:B:476:HOH:O	2.10	0.69
1:A:53:VAL:HG22	1:A:180[A]:VAL:HG12	1.75	0.69
1:B:107:ASN:HD21	1:B:187:THR:H	1.42	0.67
1:A:222:CYS:SG	2:A:401:HEM:C3C	2.88	0.66
1:A:219:CYS:SG	2:A:401:HEM:C3B	2.90	0.62
1:B:23:ASP:OD1	1:B:200:LYS:HE2	2.03	0.58
1:A:131:VAL:HG12	7:A:537:HOH:O	2.02	0.58
1:B:259[B]:ASN:CG	7:B:379:HOH:O	2.40	0.56
1:A:53:VAL:HG22	1:A:180[A]:VAL:CG1	2.36	0.55
1:A:144:VAL:HG11	1:A:174:MET:SD	2.47	0.55
1:B:179:GLU:OE2	6:B:350:BU3:H43	2.07	0.54
1:B:34:LYS:HB2	1:B:226[B]:LEU:HD23	1.92	0.51
1:B:35:PRO:O	1:B:226[B]:LEU:HD21	2.12	0.48
1:A:114:GLN:HE22	1:A:129:GLY:H	1.62	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:131:VAL:HG23	4:A:346:IMD:H4	1.96	0.47
1:A:76:CYS:CB	2:A:400:HEM:C3C	2.98	0.47
1:B:43:LEU:HD21	1:B:109:VAL:CG2	2.45	0.47
1:A:136:GLU:HB3	2:A:400:HEM:HBC2	1.97	0.46
1:A:222:CYS:SG	2:A:401:HEM:CBC	3.00	0.43
1:A:23:ASP:HA	1:A:200:LYS:NZ	2.34	0.43
1:A:131:VAL:CG1	7:A:494:HOH:O	2.27	0.43
1:B:34:LYS:O	1:B:226[B]:LEU:HD22	2.19	0.42
1:B:136:GLU:OE2	7:B:500:HOH:O	2.21	0.42
1:A:95:TRP:CH2	1:B:70:LEU:HD12	2.55	0.41
1:A:277:ILE:O	4:A:347:IMD:H5	2.21	0.41
1:A:131:VAL:HG23	4:A:346:IMD:C4	2.51	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:259:ASN:ND2	1:B:259[B]:ASN:ND2[1_454]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/345 (90%)	306 (99%)	4 (1%)	0	100	100
1	B	303/345 (88%)	300 (99%)	3 (1%)	0	100	100
All	All	613/690 (89%)	606 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/276 (91%)	249 (100%)	1 (0%)	95	85
1	B	243/276 (88%)	242 (100%)	1 (0%)	95	85
All	All	493/552 (89%)	491 (100%)	2 (0%)	95	85

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	227	ASN
1	B	34	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN
1	A	126	GLN
1	B	107	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IMD	A	346	2	5,5,5	1.61	2 (40%)	5,5,5	0.86	0
4	IMD	A	347	-	5,5,5	1.30	0	5,5,5	0.20	0
5	OXL	A	348	-	0,5,5	0.00	-	0,6,6	0.00	-
5	OXL	A	349	-	0,5,5	0.00	-	0,6,6	0.00	-
6	BU3	A	350	-	5,5,5	0.27	0	6,6,6	0.20	0
2	HEM	A	400	1,4	49,50,50	1.97	10 (20%)	46,82,82	2.04	10 (21%)
2	HEM	A	401	1	49,50,50	2.06	13 (26%)	46,82,82	2.44	14 (30%)
4	IMD	B	346	2	5,5,5	1.34	0	5,5,5	0.63	0
4	IMD	B	347	-	5,5,5	1.26	0	5,5,5	0.28	0
5	OXL	B	348	-	0,5,5	0.00	-	0,6,6	0.00	-
5	OXL	B	349	-	0,5,5	0.00	-	0,6,6	0.00	-
6	BU3	B	350	-	5,5,5	0.47	0	6,6,6	0.48	0
2	HEM	B	400	1,4	49,50,50	2.12	14 (28%)	46,82,82	2.08	10 (21%)
2	HEM	B	401	1	49,50,50	2.16	10 (20%)	46,82,82	2.17	12 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	A	346	2	-	0/0/0/0	0/1/1/1
4	IMD	A	347	-	-	0/0/0/0	0/1/1/1
5	OXL	A	348	-	-	0/0/4/4	0/0/0/0
5	OXL	A	349	-	-	0/0/4/4	0/0/0/0
6	BU3	A	350	-	-	0/4/4/4	0/0/0/0
2	HEM	A	400	1,4	-	0/14/114/114	0/0/8/8
2	HEM	A	401	1	-	0/14/114/114	0/0/8/8
4	IMD	B	346	2	-	0/0/0/0	0/1/1/1
4	IMD	B	347	-	-	0/0/0/0	0/1/1/1
5	OXL	B	348	-	-	0/0/4/4	0/0/0/0
5	OXL	B	349	-	-	0/0/4/4	0/0/0/0
6	BU3	B	350	-	-	0/4/4/4	0/0/0/0
2	HEM	B	400	1,4	-	0/14/114/114	0/0/8/8
2	HEM	B	401	1	-	0/14/114/114	0/0/8/8

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	400	HEM	C3B-C2B	-5.98	1.33	1.43
2	A	401	HEM	C3C-C2C	-5.78	1.33	1.43
2	A	400	HEM	C3B-C2B	-5.63	1.33	1.43
2	B	401	HEM	C3D-C4D	5.59	1.46	1.44
2	B	401	HEM	C3B-C2B	-5.52	1.34	1.43
2	B	400	HEM	C3C-C2C	-5.51	1.34	1.43
2	A	400	HEM	C3C-C2C	-5.27	1.34	1.43
2	B	401	HEM	C2D-C1D	-5.18	1.43	1.44
2	A	401	HEM	C3C-CAC	5.16	1.56	1.40
2	B	401	HEM	C3C-CAC	5.09	1.56	1.40
2	A	401	HEM	C3B-C2B	-5.09	1.34	1.43
2	B	400	HEM	C4A-C3A	5.01	1.46	1.40
2	B	401	HEM	C3C-C2C	-4.93	1.35	1.43
2	A	400	HEM	C3C-CAC	4.82	1.55	1.40
2	A	401	HEM	C3B-CAB	4.76	1.55	1.40
2	A	400	HEM	C3B-CAB	4.62	1.55	1.40
2	A	400	HEM	C4A-C3A	4.49	1.45	1.40
2	B	400	HEM	C3C-CAC	4.45	1.54	1.40
2	B	400	HEM	C3B-CAB	4.38	1.54	1.40
2	B	401	HEM	C3B-CAB	4.14	1.53	1.40
2	A	401	HEM	C3D-C2D	3.70	1.50	1.43
2	A	401	HEM	C4A-C3A	3.62	1.44	1.40
2	B	401	HEM	C4A-C3A	3.61	1.44	1.40
2	B	400	HEM	C3D-C2D	3.56	1.50	1.43
2	A	400	HEM	C3D-C2D	3.46	1.49	1.43
2	B	400	HEM	C2B-C1B	3.08	1.45	1.44
2	B	401	HEM	C3D-C2D	3.00	1.49	1.43
2	B	400	HEM	FE-NA	2.85	2.04	1.92
2	A	401	HEM	C2D-C1D	-2.63	1.43	1.44
2	B	401	HEM	CMB-C2B	2.56	1.55	1.47
2	A	401	HEM	FE-NB	2.42	2.06	1.97
2	A	401	HEM	CMB-C2B	2.42	1.54	1.47
2	A	401	HEM	CMC-C2C	2.40	1.54	1.47
4	A	346	IMD	C2-N1	2.34	1.36	1.31
2	B	401	HEM	CMC-C2C	2.33	1.54	1.47
2	B	400	HEM	CMB-C2B	2.29	1.54	1.47
2	A	401	HEM	C4A-NA	2.25	1.41	1.36
2	B	400	HEM	CAA-C2A	2.22	1.55	1.52
4	A	346	IMD	C2-N3	2.19	1.36	1.31
2	A	400	HEM	CMC-C2C	2.16	1.54	1.47
2	B	400	HEM	CHB-C1B	2.16	1.38	1.35
2	A	400	HEM	CMD-C2D	2.14	1.54	1.47
2	A	400	HEM	C2B-C1B	-2.12	1.44	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	400	HEM	CMA-C3A	2.09	1.56	1.51
2	B	400	HEM	CMC-C2C	2.08	1.53	1.47
2	A	401	HEM	CMD-C2D	2.05	1.53	1.47
2	B	400	HEM	C1A-C2A	2.04	1.47	1.43
2	A	400	HEM	CMB-C2B	2.03	1.53	1.47
2	A	401	HEM	FE-NA	2.02	2.01	1.92

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C3B-C4B-NB	-10.21	106.69	114.00
2	A	400	HEM	C3B-C4B-NB	-8.43	107.97	114.00
2	B	400	HEM	C3B-C4B-NB	-8.37	108.01	114.00
2	B	401	HEM	C3B-C4B-NB	-8.35	108.02	114.00
2	B	401	HEM	C4D-ND-C1D	6.47	111.79	105.16
2	B	400	HEM	C4D-ND-C1D	5.85	111.15	105.16
2	A	401	HEM	C4D-ND-C1D	5.61	110.90	105.16
2	A	400	HEM	C4D-ND-C1D	5.29	110.58	105.16
2	A	401	HEM	C1B-NB-C4B	4.21	109.47	105.16
2	B	400	HEM	CBD-CAD-C3D	-3.92	105.82	114.37
2	A	400	HEM	CBD-CAD-C3D	-3.91	105.84	114.37
2	B	401	HEM	CBD-CAD-C3D	-3.68	106.34	114.37
2	B	401	HEM	C2D-C1D-ND	-3.67	108.60	112.93
2	B	400	HEM	C2D-C1D-ND	-3.56	108.73	112.93
2	A	401	HEM	C2D-C1D-ND	-3.34	108.99	112.93
2	A	401	HEM	CHD-C1D-ND	3.31	127.34	124.58
2	B	401	HEM	CHD-C1D-ND	3.07	127.13	124.58
2	A	400	HEM	CMA-C3A-C4A	-3.01	123.99	128.62
2	A	400	HEM	C4A-CHB-C1B	-2.96	123.58	127.47
2	A	400	HEM	C2D-C1D-ND	-2.95	109.44	112.93
2	A	401	HEM	CBA-CAA-C2A	-2.87	107.64	112.69
2	A	401	HEM	CBD-CAD-C3D	-2.86	108.13	114.37
2	B	401	HEM	CBA-CAA-C2A	-2.80	107.76	112.69
2	B	400	HEM	CHD-C1D-ND	2.75	126.87	124.58
2	A	401	HEM	C4A-CHB-C1B	-2.71	123.91	127.47
2	A	401	HEM	CHC-C4B-NB	2.66	126.79	124.58
2	A	400	HEM	C1B-NB-C4B	2.48	107.70	105.16
2	A	401	HEM	C4C-NC-C1C	2.46	108.09	105.53
2	B	400	HEM	C1B-NB-C4B	2.37	107.58	105.16
2	A	401	HEM	CHD-C4C-NC	2.36	126.78	124.73
2	B	401	HEM	CMA-C3A-C4A	-2.35	125.00	128.62
2	B	401	HEM	C1A-CHA-C4D	-2.33	124.41	127.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C3A-C4A-NA	-2.32	107.66	109.41
2	A	401	HEM	CMA-C3A-C4A	-2.31	125.06	128.62
2	B	400	HEM	CAD-C3D-C4D	2.30	128.66	124.53
2	B	401	HEM	C4A-CHB-C1B	-2.25	124.51	127.47
2	B	401	HEM	C4A-C3A-C2A	2.24	108.56	107.00
2	A	401	HEM	CHA-C4D-ND	2.24	127.38	124.31
2	A	400	HEM	CHC-C4B-NB	2.23	126.44	124.58
2	B	401	HEM	CHA-C4D-ND	2.22	127.36	124.31
2	A	400	HEM	C4A-C3A-C2A	2.21	108.53	107.00
2	B	400	HEM	C1A-CHA-C4D	-2.20	124.58	127.47
2	A	400	HEM	CAD-C3D-C4D	2.18	128.45	124.53
2	B	400	HEM	O2A-CGA-CBA	2.13	121.76	114.22
2	B	401	HEM	C1B-NB-C4B	2.11	107.32	105.16
2	B	400	HEM	CMA-C3A-C4A	-2.04	125.49	128.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/345 (90%)	0.73	28 (8%) 10 10	12, 19, 30, 37	0
1	B	306/345 (88%)	0.40	18 (5%) 22 23	11, 16, 29, 38	0
All	All	618/690 (89%)	0.56	46 (7%) 14 15	11, 17, 30, 38	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	GLY	8.3
1	B	123	LEU	7.6
1	A	23	ASP	7.0
1	B	253	GLY	6.9
1	A	22	ALA	6.6
1	A	166	LYS	4.6
1	B	122	ASP	4.4
1	A	164	GLY	4.3
1	B	21	ALA	4.1
1	B	132	GLN	4.1
1	A	24	GLU	4.0
1	A	201	LYS	3.9
1	B	241	LYS	3.8
1	A	44	LYS	3.7
1	A	254	ARG	3.5
1	A	163	PRO	3.3
1	A	27	GLN	3.2
1	A	167	ASP	3.0
1	A	48	ALA	2.9
1	B	119	ARG	2.9
1	B	121	LYS	2.8
1	A	240	GLU	2.7
1	B	120	ALA	2.6
1	B	338	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	25	LEU	2.5
1	A	200	LYS	2.4
1	A	34	LYS	2.4
1	A	160	LYS	2.4
1	B	254	ARG	2.4
1	B	81	LEU	2.4
1	A	156	ALA	2.2
1	B	114	GLN	2.2
1	A	84	GLY	2.2
1	B	240	GLU	2.2
1	A	309	ASP	2.1
1	A	212	LYS	2.1
1	B	277	ILE	2.1
1	B	48	ALA	2.1
1	A	32	LEU	2.1
1	B	86	LEU	2.1
1	A	202	ALA	2.0
1	A	194	GLN	2.0
1	B	275	VAL	2.0
1	A	198	GLY	2.0
1	A	308	ASP	2.0
1	A	26	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	IMD	A	346	5/5	0.19	4.46	16,17,20,21	0
6	BU3	A	350	6/6	0.15	2.61	25,26,27,28	0
5	OXL	A	348	6/6	0.14	1.88	27,28,30,33	0
2	HEM	A	400	43/43	0.13	1.08	11,13,21,26	0
5	OXL	B	349	6/6	0.14	0.99	40,41,42,43	0
2	HEM	A	401	43/43	0.12	0.93	13,14,18,19	0
5	OXL	A	349	6/6	0.13	0.56	31,33,34,35	0
6	BU3	B	350	6/6	0.17	0.28	38,41,42,43	0
4	IMD	B	347	5/5	0.12	0.16	18,20,20,20	0
4	IMD	A	347	5/5	0.11	0.04	20,20,22,22	0
5	OXL	B	348	6/6	0.10	-0.10	34,36,37,38	0
2	HEM	B	401	43/43	0.10	-0.22	9,11,13,14	0
2	HEM	B	400	43/43	0.10	-0.43	10,13,22,28	0
4	IMD	B	346	5/5	0.11	-1.42	22,23,25,25	0
3	CA	A	402	1/1	0.09	-1.64	13,13,13,13	0
3	CA	B	402	1/1	0.09	-2.37	10,10,10,10	0

6.5 Other polymers ⓘ

There are no such residues in this entry.