



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 07:18 AM GMT

PDB ID : 2HT1
Title : The closed ring structure of the Rho transcription termination factor in complex with nucleic acid in the motor domains
Authors : Skordalakes, E.; Berger, J.M.
Deposited on : 2006-07-24
Resolution : 3.51 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance






The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1256 (3.74-3.30)
Clashscore	79885	1020 (3.68-3.36)
Ramachandran outliers	78287	1082 (3.70-3.34)
Sidechain outliers	78261	1082 (3.70-3.34)
RSRZ outliers	66119	1256 (3.74-3.30)
RNA backbone	1838	1008 (4.26-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	J	2	
1	K	2	
2	M	5	
3	A	433	
3	B	433	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5223 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*UP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	J	2	Total	C	N	O	P	0	0	0
			37	18	5	13	1			
1	K	2	Total	C	N	O	P	0	0	0
			37	18	5	13	1			

- Molecule 2 is a RNA chain called 5'-R(*UP*CP*UP*CP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	5	Total	C	N	O	P	0	0	0
			97	45	12	36	4			

- Molecule 3 is a protein called Transcription termination factor rho.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	324	Total	C	N	O	S	0	0	0
			2521	1585	454	474	8			
3	B	324	Total	C	N	O	S	0	0	0
			2521	1585	454	474	8			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	INITIATING METHIONINE	UNP P0AG30
A	-20	GLY	-	CLONING ARTIFACT	UNP P0AG30
A	-19	SER	-	CLONING ARTIFACT	UNP P0AG30
A	-18	SER	-	CLONING ARTIFACT	UNP P0AG30
A	-17	HIS	-	EXPRESSION TAG	UNP P0AG30
A	-16	HIS	-	EXPRESSION TAG	UNP P0AG30
A	-15	HIS	-	EXPRESSION TAG	UNP P0AG30
A	-14	HIS	-	EXPRESSION TAG	UNP P0AG30
A	-13	HIS	-	EXPRESSION TAG	UNP P0AG30
A	-12	HIS	-	EXPRESSION TAG	UNP P0AG30

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	SER	-	CLONING ARTIFACT	UNP P0AG30
A	-10	SER	-	CLONING ARTIFACT	UNP P0AG30
A	-9	GLY	-	CLONING ARTIFACT	UNP P0AG30
A	-8	GLU	-	CLONING ARTIFACT	UNP P0AG30
A	-7	ASN	-	CLONING ARTIFACT	UNP P0AG30
A	-6	LEU	-	CLONING ARTIFACT	UNP P0AG30
A	-5	TYR	-	CLONING ARTIFACT	UNP P0AG30
A	-4	PHE	-	CLONING ARTIFACT	UNP P0AG30
A	-3	GLN	-	CLONING ARTIFACT	UNP P0AG30
A	-2	ALA	-	CLONING ARTIFACT	UNP P0AG30
A	-1	GLY	-	CLONING ARTIFACT	UNP P0AG30
A	0	HIS	-	CLONING ARTIFACT	UNP P0AG30
B	-21	MET	-	INITIATING METHIONINE	UNP P0AG30
B	-20	GLY	-	CLONING ARTIFACT	UNP P0AG30
B	-19	SER	-	CLONING ARTIFACT	UNP P0AG30
B	-18	SER	-	CLONING ARTIFACT	UNP P0AG30
B	-17	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-16	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-15	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-14	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-13	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-12	HIS	-	EXPRESSION TAG	UNP P0AG30
B	-11	SER	-	CLONING ARTIFACT	UNP P0AG30
B	-10	SER	-	CLONING ARTIFACT	UNP P0AG30
B	-9	GLY	-	CLONING ARTIFACT	UNP P0AG30
B	-8	GLU	-	CLONING ARTIFACT	UNP P0AG30
B	-7	ASN	-	CLONING ARTIFACT	UNP P0AG30
B	-6	LEU	-	CLONING ARTIFACT	UNP P0AG30
B	-5	TYR	-	CLONING ARTIFACT	UNP P0AG30
B	-4	PHE	-	CLONING ARTIFACT	UNP P0AG30
B	-3	GLN	-	CLONING ARTIFACT	UNP P0AG30
B	-2	ALA	-	CLONING ARTIFACT	UNP P0AG30
B	-1	GLY	-	CLONING ARTIFACT	UNP P0AG30
B	0	HIS	-	CLONING ARTIFACT	UNP P0AG30

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-R(*UP*C)-3'

Chain J: 




- Molecule 1: 5'-R(*UP*C)-3'

Chain K: 




- Molecule 2: 5'-R(*UP*CP*UP*CP*U)-3'

Chain M: 




- Molecule 3: Transcription termination factor rho

Chain A: 



- Molecule 3: Transcription termination factor rho

Age Group	Percentage
18-29	90%
30-49	83%
50-64	70%
65+	56%



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	257.69Å 257.69Å 257.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.51 20.00 – 3.51	Depositor EDS
% Data completeness (in resolution range)	91.9 (20.00-3.51) 90.3 (20.00-3.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 3.52Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.294 , 0.328 0.274 , 0.323	Depositor DCC
R_{free} test set	854 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	108.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 77.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 16675 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	5223	wwPDB-VP
Average B, all atoms (Å ²)	118.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	J	0.71	0/40	1.37	0/60
1	K	0.74	0/40	1.24	0/60
2	M	0.72	0/106	1.27	0/162
3	A	0.33	0/2559	0.51	0/3450
3	B	0.33	0/2559	0.51	0/3450
All	All	0.35	0/5304	0.56	0/7182

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	37	0	23	1	0
1	K	37	0	23	0	0
2	M	97	0	54	12	0
3	A	2521	0	2580	35	0
3	B	2521	0	2580	41	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5223	0	5260	86	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (86) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:M:3:U:C2'	2:M:4:C:H5''	1.87	1.04
2:M:3:U:C3'	2:M:4:C:H5''	1.93	0.99
2:M:3:U:O2'	2:M:4:C:H5''	1.65	0.94
2:M:2:C:H4'	2:M:3:U:OP1	1.68	0.93
3:A:66:ARG:HH11	3:A:66:ARG:HG2	1.36	0.89
2:M:3:U:C3'	2:M:4:C:C5'	2.58	0.81
2:M:2:C:C4'	2:M:3:U:OP1	2.29	0.79
2:M:3:U:H3'	2:M:4:C:C5'	2.13	0.77
3:B:266:SER:HB2	3:B:269:ARG:HB2	1.69	0.75
3:A:66:ARG:NH1	3:A:66:ARG:HG2	1.99	0.70
3:A:55:LEU:HB2	3:A:97:ILE:HD12	1.74	0.69
3:B:171:GLY:H	3:B:314:THR:HG22	1.58	0.68
3:A:56:GLU:O	3:A:63:GLY:HA3	1.95	0.67
3:B:188:LEU:HD22	3:B:263:LEU:HD21	1.77	0.66
3:A:39:LEU:HD22	3:A:49:ILE:HD12	1.76	0.66
3:B:316:ILE:HG13	3:B:316:ILE:O	1.94	0.66
3:B:161:VAL:HG21	3:B:191:ILE:HD12	1.77	0.66
2:M:1:U:H4'	2:M:2:C:OP1	1.92	0.63
3:A:223:VAL:HG12	3:A:225:GLY:H	1.64	0.61
3:B:89:PHE:HB2	3:B:91:LEU:HG	1.82	0.60
2:M:3:U:H3'	2:M:4:C:H5'	1.86	0.58
3:B:171:GLY:H	3:B:314:THR:CG2	2.16	0.57
3:A:39:LEU:HD12	3:A:111:PHE:CE1	2.40	0.57
3:B:39:LEU:HD22	3:B:49:ILE:HD12	1.87	0.56
3:A:83:PRO:HA	3:A:86:ILE:HD12	1.88	0.55
3:B:210:ASP:HB3	3:B:269:ARG:HG2	1.89	0.55
2:M:3:U:HO2'	2:M:4:C:H5''	1.71	0.55
3:B:263:LEU:HG	3:B:316:ILE:HD11	1.90	0.54
3:B:212:ARG:HB3	3:B:215:GLU:HB2	1.90	0.53
3:A:62:PHE:HB3	3:A:83:PRO:HG3	1.90	0.53
3:A:66:ARG:HH11	3:A:66:ARG:CG	2.16	0.52
3:A:191:ILE:O	3:A:195:ILE:HG12	2.10	0.52
3:A:321:ILE:HD11	3:A:332:TYR:CE2	2.45	0.52
3:B:272:ARG:HD3	3:B:327:MET:SD	2.50	0.51
3:A:254:VAL:HG21	3:A:313:LEU:HB2	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:261:ILE:HG12	3:A:314:THR:HG23	1.94	0.50
2:M:5:U:O5'	2:M:5:U:H6	1.95	0.50
3:B:252:ARG:NH1	3:B:255:GLU:OE2	2.44	0.49
3:B:162:LEU:HD11	3:B:195:ILE:HG13	1.95	0.49
3:B:247:ILE:HB	3:B:300:PHE:CE1	2.48	0.48
3:A:177:VAL:HG13	3:A:321:ILE:HD12	1.94	0.48
3:A:176:ILE:HB	3:A:318:THR:HG22	1.95	0.47
3:B:31:LYS:HA	3:B:34:ILE:HD12	1.95	0.47
1:J:2:C:H42	3:B:66:ARG:HH12	1.62	0.47
3:B:230:SER:OG	3:B:239:HIS:HD2	1.98	0.47
3:B:207:LEU:HA	3:B:228:VAL:O	2.15	0.47
3:B:65:LEU:HB2	3:B:79:ILE:HB	1.97	0.47
3:A:247:ILE:HB	3:A:300:PHE:CE1	2.50	0.46
3:B:136:LEU:HD22	3:B:309:GLU:HG3	1.96	0.46
3:B:268:THR:HA	3:B:331:ILE:HG21	1.98	0.46
3:A:54:VAL:HG22	3:A:96:THR:HA	1.97	0.46
3:A:136:LEU:HD23	3:A:307:VAL:HG21	1.97	0.46
3:B:57:ILE:H	3:B:93:THR:HB	1.80	0.46
3:A:278:VAL:O	3:B:283:LYS:NZ	2.49	0.46
3:B:82:SER:O	3:B:86:ILE:HG12	2.16	0.45
2:M:3:U:HO2'	2:M:4:C:H6	1.63	0.45
3:B:345:LEU:HB3	3:B:346:SER:H	1.68	0.44
3:A:187:LEU:O	3:A:191:ILE:HG12	2.16	0.44
3:A:184:LYS:HG3	3:A:185:THR:N	2.31	0.44
3:B:177:VAL:HB	3:B:344:HIS:HD2	1.83	0.44
3:B:275:ASN:HA	3:B:293:ALA:HB1	2.00	0.44
3:B:248:GLU:OE2	3:B:252:ARG:NH2	2.51	0.44
3:A:39:LEU:HD13	3:A:49:ILE:HG21	1.99	0.44
3:A:210:ASP:HB3	3:A:269:ARG:HG3	2.00	0.43
3:B:209:ILE:HD13	3:B:270:LEU:HB2	2.00	0.43
3:B:187:LEU:O	3:B:191:ILE:HG12	2.18	0.43
3:B:105:LYS:O	3:B:108:GLU:HB2	2.19	0.43
3:B:250:ALA:HB2	3:B:262:ILE:HD11	2.01	0.43
3:B:206:VAL:HG22	3:B:263:LEU:HB3	2.00	0.43
3:A:240:VAL:HG13	3:A:274:TYR:CE1	2.53	0.43
3:A:173:ARG:HB3	3:A:301:PHE:CZ	2.53	0.43
3:B:188:LEU:HD11	3:B:318:THR:HG21	2.01	0.43
3:A:161:VAL:HB	3:A:191:ILE:HD12	2.00	0.43
3:A:307:VAL:HG12	3:A:309:GLU:H	1.83	0.42
3:B:167:PRO:HD3	3:B:199:HIS:CE1	2.54	0.42
3:A:328:ASP:HA	3:A:331:ILE:HG22	2.01	0.41
3:A:294:LEU:O	3:A:297:PRO:HD2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:B:187:LEU:HD11	3:B:191:ILE:HD11	2.02	0.41
3:B:272:ARG:O	3:B:276:THR:HG23	2.21	0.41
3:B:171:GLY:N	3:B:314:THR:HG22	2.32	0.41
3:A:259:ASP:HA	3:A:312:SER:H	1.86	0.41
3:A:321:ILE:HD11	3:A:332:TYR:CD2	2.56	0.40
3:A:172:GLN:H	3:A:314:THR:HB	1.86	0.40
3:A:269:ARG:HD2	3:A:269:ARG:HA	1.90	0.40
3:A:275:ASN:O	3:B:283:LYS:NZ	2.52	0.40
3:B:250:ALA:HB1	3:B:313:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	322/433 (74%)	283 (88%)	36 (11%)	3 (1%)	25	80
3	B	322/433 (74%)	295 (92%)	23 (7%)	4 (1%)	19	75
All	All	644/866 (74%)	578 (90%)	59 (9%)	7 (1%)	21	77

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	121	PHE
3	B	125	GLU
3	B	141	ALA
3	A	346	SER
3	B	48	ASP
3	A	288	GLY
3	A	51	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	272/369 (74%)	252 (93%)	20 (7%)	20	66
3	B	272/369 (74%)	258 (95%)	14 (5%)	33	80
All	All	544/738 (74%)	510 (94%)	34 (6%)	25	73

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	60	ASP
3	A	66	ARG
3	A	77	ASP
3	A	101	ILE
3	A	120	ASN
3	A	122	ASP
3	A	131	ILE
3	A	202	CYS
3	A	222	LEU
3	A	252	ARG
3	A	263	LEU
3	A	267	ILE
3	A	269	ARG
3	A	276	THR
3	A	286	THR
3	A	314	THR
3	A	321	ILE
3	A	328	ASP
3	A	331	ILE
3	A	340	ASN
3	B	134	GLU
3	B	148	GLU
3	B	161	VAL
3	B	186	MET
3	B	215	GLU
3	B	222	LEU
3	B	242	VAL
3	B	267	ILE
3	B	284	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	286	THR
3	B	305	ARG
3	B	314	THR
3	B	326	LYS
3	B	341	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	32	GLN
3	B	90	ASN
3	B	198	ASN
3	B	199	HIS
3	B	239	HIS
3	B	344	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	J	1/2 (50%)	0	0
1	K	1/2 (50%)	1 (100%)	0
2	M	5/5 (100%)	4 (80%)	2 (40%)
All	All	7/9 (77%)	5 (71%)	2 (28%)

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	K	2	C
2	M	2	C
2	M	3	U
2	M	4	C
2	M	5	U

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	M	1	U
2	M	2	C

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	600	-	4,4,4	0.18	0	6,6,6	0.07	0
4	SO4	B	601	-	4,4,4	0.20	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	600	-	-	0/0/0/0	0/0/0/0
4	SO4	B	601	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	J	2/2 (100%)	3.26	2 (100%) 0 0	198, 198, 198, 216	0
1	K	2/2 (100%)	3.74	2 (100%) 0 0	210, 210, 210, 231	0
2	M	5/5 (100%)	6.13	5 (100%) 0 0	100, 104, 130, 175	5 (100%)
3	A	324/433 (74%)	0.14	17 (5%) 26 11	66, 115, 211, 266	0
3	B	324/433 (74%)	0.07	12 (3%) 39 18	66, 89, 204, 247	0
All	All	657/875 (75%)	0.17	38 (5%) 22 10	66, 101, 210, 266	5 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	4	C	9.3
2	M	1	U	6.5
2	M	5	U	5.9
2	M	2	C	5.5
3	B	353	ARG	4.6
1	K	2	C	4.0
3	A	149	ARG	4.0
3	B	51	GLY	3.7
3	B	354	VAL	3.7
3	B	48	ASP	3.6
3	A	51	GLY	3.5
2	M	3	U	3.5
1	K	1	U	3.4
3	A	47	GLU	3.4
3	A	349	ILE	3.4
1	J	2	C	3.4
3	A	98	SER	3.2
1	J	1	U	3.1
3	A	352	LYS	3.1
3	B	128	ARG	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	42	HIS	2.9
3	A	348	LYS	2.8
3	A	123	LYS	2.7
3	B	352	LYS	2.7
3	B	44	LYS	2.6
3	B	47	GLU	2.6
3	A	156	ASP	2.6
3	A	128	ARG	2.5
3	A	44	LYS	2.5
3	B	122	ASP	2.4
3	A	48	ASP	2.4
3	B	351	GLU	2.3
3	B	87	ARG	2.2
3	A	351	GLU	2.2
3	A	110	TYR	2.2
3	A	87	ARG	2.2
3	A	155	GLU	2.2
3	B	110	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	A	600	5/5	0.13	-0.36	99,99,99,99	0
4	SO4	B	601	5/5	0.11	-1.49	94,94,94,94	0

6.5 Other polymers ⓘ

There are no such residues in this entry.