



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 01:45 AM GMT

PDB ID : 4HT1  
Title : Human TWEAK in complex with the Fab fragment of a neutralizing antibody  
Authors : Lammens, A.  
Deposited on : 2012-10-31  
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	T	154	
2	H	227	
3	L	218	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4373 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	T	131	Total	C	N	O	S	40	0	0
			1033	667	184	179	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
T	1	MET	-	INITIATING METHIONINE	UNP O43508

- Molecule 2 is a protein called chimeric antibody Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	210	Total	C	N	O	S	19	0	0
			1580	1009	255	310	6			

- Molecule 3 is a protein called chimeric antibody Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	217	Total	C	N	O	S	8	1	0
			1660	1040	279	334	7			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	T	20	Total	O	0	0
			20	20		
4	H	30	Total	O	0	0
			30	30		
4	L	50	Total	O	0	0
			50	50		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 12

Chain T: 




- Molecule 2: chimeric antibody Fab

Chain H: 



- Molecule 3: chimeric antibody Fab

Chain L: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.63Å 101.63Å 57.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.34 – 2.50 29.34 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.34-2.50) 99.4 (29.34-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.27 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.183 , 0.211 0.183 , 0.211	Depositor DCC
$R_{free}$ test set	2293 reflections (10.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 9.9	EDS
Estimated twinning fraction	0.400 for -k,-h,-l 0.087 for -h,-k,l 0.097 for h,-h-k,-l 0.379 for -k,-h,-l	Xtriage
Reported twinning fraction	0.400 for -k,-h,-l	Depositor
L-test for twinning	$\langle  L  \rangle = 0.40$ , $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 22932 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4373	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	T	0.63	0/1056	0.83	0/1430
2	H	0.60	0/1621	0.76	0/2214
3	L	0.62	0/1699	0.75	1/2312 (0.0%)
All	All	0.62	0/4376	0.77	1/5956 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	77	GLY	N-CA-C	-5.17	100.19	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	T	1033	0	1035	22	0
2	H	1580	0	1527	45	0
3	L	1660	0	1614	37	0
4	H	30	0	0	1	0
4	L	50	0	0	1	0
4	T	20	0	0	1	0
All	All	4373	0	4176	100	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (100) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:L:149:LYS:HB3	3:L:201:THR:HB	1.58	0.85
2:H:67:ARG:HG2	2:H:68:PHE:CD1	2.13	0.84
2:H:67:ARG:HG2	2:H:68:PHE:HD1	1.38	0.84
1:T:133:THR:HG22	1:T:134:LEU:H	1.44	0.81
1:T:91:VAL:HG21	3:L:96:ARG:NH2	2.00	0.77
2:H:202:ILE:HG22	2:H:217:LYS:HB3	1.69	0.75
3:L:2:ILE:HD13	3:L:29:ILE:HD11	1.69	0.75
2:H:158:THR:HG22	2:H:206:ASN:HB3	1.69	0.73
2:H:128:VAL:HG21	2:H:214:VAL:HG11	1.72	0.72
2:H:67:ARG:HD2	2:H:68:PHE:HE1	1.56	0.70
2:H:200:THR:HG23	4:H:302:HOH:O	1.92	0.69
2:H:150:LYS:HG3	2:H:184:SER:OG	1.93	0.68
3:L:21:ILE:HG12	3:L:106:THR:HG21	1.77	0.67
3:L:151:GLN:HG3	3:L:199:GLU:HB3	1.80	0.63
3:L:126:ASP:O	3:L:130:LYS:HG3	1.99	0.62
2:H:133:PRO:HG2	2:H:220:PRO:HA	1.80	0.62
2:H:37:VAL:HG22	2:H:47:TRP:HA	1.82	0.62
1:T:83:LYS:O	1:T:133:THR:HG23	1.99	0.62
2:H:170:VAL:HG22	2:H:189:VAL:HG22	1.84	0.60
1:T:73:HIS:CE1	1:T:114:LEU:HD13	2.36	0.60
2:H:81:LEU:CD2	2:H:83:MET:HG2	2.32	0.60
2:H:157:VAL:HG22	2:H:207:HIS:HB2	1.83	0.59
2:H:68:PHE:CD2	2:H:81:LEU:HD21	2.36	0.59
3:L:61:ARG:HB2	3:L:76:SER:O	2.02	0.59
3:L:61:ARG:NH1	3:L:82:ASP:OD2	2.33	0.58
2:H:48:ILE:HG23	2:H:64:LEU:HD11	1.85	0.57
1:T:91:VAL:HG21	3:L:96:ARG:HH22	1.68	0.57
3:L:159:GLN:HG3	3:L:162:ASN:HD21	1.71	0.56
2:H:31:THR:HG22	2:H:54:ARG:NH1	2.21	0.55
1:T:63:ALA:HB2	1:T:125:PRO:HG3	1.88	0.55
1:T:32:ASP:OD1	1:T:34:THR:OG1	2.24	0.54
2:H:9:GLY:H	2:H:114:THR:HG21	1.74	0.53
2:H:202:ILE:HG22	2:H:217:LYS:CB	2.38	0.52
1:T:16:TYR:CE1	1:T:40:GLU:HG3	2.45	0.52
2:H:64:LEU:O	2:H:67:ARG:HB3	2.10	0.52
2:H:31:THR:HG22	2:H:54:ARG:HH11	1.75	0.52
2:H:154:PRO:HD2	2:H:209:PRO:HD2	1.89	0.52
3:L:199:GLU:OE1	3:L:210:THR:OG1	2.28	0.51
2:H:81:LEU:HD23	2:H:83:MET:HG2	1.91	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:67:ARG:HD2	2:H:68:PHE:CE1	2.43	0.51
3:L:154:VAL:HG12	3:L:159:GLN:OE1	2.10	0.51
3:L:171:ASP:HB3	3:L:174:ASP:OD2	2.11	0.51
1:T:31:VAL:HG22	1:T:136:TRP:CH2	2.47	0.50
3:L:61:ARG:HH22	3:L:82:ASP:CG	2.14	0.50
2:H:103:TYR:CZ	3:L:97:PRO:HB3	2.47	0.50
3:L:205:LEU:HD22	3:L:209:VAL:HG21	1.94	0.50
2:H:88:ALA:O	2:H:91:THR:HB	2.11	0.50
3:L:202:HIS:HB3	3:L:205:LEU:HD12	1.94	0.50
2:H:126:PRO:HG2	2:H:212:THR:HG21	1.94	0.49
3:L:142:ASN:N	3:L:142:ASN:OD1	2.45	0.49
2:H:173:PHE:HB3	3:L:166:SER:OG	2.13	0.49
3:L:36:TYR:HE2	3:L:89:GLN:HB3	1.77	0.49
2:H:38:ARG:HB3	2:H:48:ILE:HD11	1.94	0.48
3:L:93:TYR:CD1	3:L:97:PRO:HA	2.49	0.48
3:L:129:LEU:HB3	3:L:187:LYS:HE3	1.96	0.48
3:L:47:LEU:HA	3:L:58:VAL:HG21	1.94	0.48
1:T:142:ALA:HB3	1:T:145:LEU:HB2	1.96	0.47
2:H:166:LEU:HD21	2:H:189:VAL:HG11	1.95	0.47
2:H:2:GLU:OE1	2:H:98:LYS:NZ	2.48	0.47
2:H:83:MET:HB2	2:H:86:LEU:HD21	1.97	0.47
3:L:188:ALA:O	3:L:192:LYS:HG3	2.15	0.47
2:H:29:PHE:O	2:H:53:VAL:HG21	2.15	0.47
2:H:30:SER:O	2:H:53:VAL:HG23	2.16	0.46
3:L:124:PRO:HD3	3:L:136:VAL:HG22	1.97	0.46
3:L:85:THR:HG22	3:L:107:GLU:HG2	1.98	0.46
3:L:48:MET:HB2	3:L:53:TYR:O	2.15	0.46
3:L:154:VAL:O	3:L:154:VAL:HG13	2.16	0.45
2:H:28:ASP:OD1	2:H:30:SER:OG	2.33	0.45
1:T:130:ARG:HD3	4:T:209:HOH:O	2.15	0.45
3:L:23:CYS:HB2	3:L:35:TRP:CH2	2.52	0.45
1:T:73:HIS:CD2	1:T:112:LEU:HD11	2.51	0.45
3:L:60:SER:O	3:L:63:LYS:HE3	2.17	0.45
1:T:71:GLN:HG3	1:T:114:LEU:HD11	1.98	0.45
1:T:73:HIS:HD2	1:T:112:LEU:HD11	1.82	0.44
2:H:124:LYS:HD2	2:H:151:ASP:O	2.18	0.44
3:L:158:LEU:HD22	3:L:158:LEU:HA	1.85	0.44
1:T:58:PHE:O	1:T:128:SER:HA	2.17	0.44
1:T:80:VAL:HG13	1:T:81:TYR:HD2	1.82	0.44
2:H:6:GLU:OE2	2:H:111:GLY:HA3	2.18	0.43
1:T:81:TYR:OH	1:T:134:LEU:HD21	2.19	0.43
3:L:120:PHE:HE1	3:L:141:ASN:OD1	2.01	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:L:140:LEU:HD12	3:L:140:LEU:N	2.34	0.43
1:T:27:ALA:HB3	1:T:139:LEU:HD11	2.01	0.43
3:L:146[A]:ARG:NH1	4:L:315:HOH:O	2.52	0.43
2:H:83:MET:CB	2:H:86:LEU:HD21	2.49	0.42
1:T:48:PRO:HG3	1:T:66:TYR:CE1	2.53	0.42
3:L:48:MET:HB3	3:L:48:MET:HE2	1.83	0.42
3:L:19:VAL:HG12	3:L:75:ILE:HB	2.01	0.42
2:H:156:PRO:O	2:H:207:HIS:CD2	2.73	0.42
2:H:70:ILE:HG12	2:H:81:LEU:HD12	2.01	0.42
2:H:146:GLY:HA2	2:H:161:TRP:CH2	2.55	0.41
1:T:144:PHE:CD2	1:T:145:LEU:HD13	2.55	0.41
3:L:127:GLU:O	3:L:130:LYS:HB2	2.19	0.41
2:H:198:THR:HG23	2:H:199:GLN:N	2.35	0.41
2:H:12:VAL:O	2:H:118:VAL:HA	2.21	0.41
1:T:31:VAL:H	1:T:31:VAL:HG23	1.64	0.41
2:H:182:LEU:HA	2:H:182:LEU:HD12	1.81	0.41
2:H:156:PRO:O	2:H:207:HIS:HD2	2.04	0.40
1:T:144:PHE:CE2	1:T:145:LEU:HD13	2.56	0.40
2:H:8:GLY:O	2:H:18:LEU:HD21	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	T	125/154 (81%)	118 (94%)	7 (6%)	0	100	100
2	H	204/227 (90%)	197 (97%)	6 (3%)	1 (0%)	38	60
3	L	216/218 (99%)	207 (96%)	9 (4%)	0	100	100
All	All	545/599 (91%)	522 (96%)	22 (4%)	1 (0%)	56	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	156	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	T	106/123 (86%)	98 (92%)	8 (8%)	19	34
2	H	175/189 (93%)	159 (91%)	16 (9%)	14	24
3	L	185/185 (100%)	171 (92%)	14 (8%)	19	33
All	All	466/497 (94%)	428 (92%)	38 (8%)	17	29

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	T	44	ASN
1	T	86	LEU
1	T	91	VAL
1	T	98	GLU
1	T	115	CYS
1	T	117	VAL
1	T	140	LYS
1	T	150	LEU
2	H	1	GLN
2	H	7	SER
2	H	25	SER
2	H	29	PHE
2	H	30	SER
2	H	43	LYS
2	H	67	ARG
2	H	91	THR
2	H	150	LYS
2	H	167	THR
2	H	182	LEU
2	H	184	SER
2	H	193	SER
2	H	203	CYS
2	H	211	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	H	215	ASP
3	L	5	THR
3	L	19	VAL
3	L	29	ILE
3	L	50	THR
3	L	52	SER
3	L	65	SER
3	L	68	ARG
3	L	133	THR
3	L	142	ASN
3	L	158	LEU
3	L	168	THR
3	L	172	SER
3	L	173	LYS
3	L	174	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	T	131/154 (85%)	-0.30	1 (0%) 83 84	7, 14, 37, 49	10 (7%)
2	H	210/227 (92%)	-0.23	0 100 100	11, 24, 41, 67	6 (2%)
3	L	217/218 (99%)	-0.26	0 100 100	11, 21, 35, 43	3 (1%)
All	All	558/599 (93%)	-0.26	1 (0%) 93 94	7, 21, 38, 67	19 (3%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	T	77	GLY	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.