



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 05:56 AM GMT

PDB ID : 2I2X
Title : Crystal structure of methanol:cobalamin methyltransferase complex MtaBC from Methanosarcina barkeri
Authors : Hagemeier, C.H.; Kruer, M.; Thauer, R.K.; Warkentin, E.; Ermler, U.
Deposited on : 2006-08-17
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	461	
1	C	461	
1	E	461	
1	G	461	
1	I	461	
1	K	461	
1	M	461	
1	O	461	
2	B	258	
2	D	258	
2	F	258	
2	H	258	
2	J	258	
2	L	258	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	N	258	
2	P	258	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 45566 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	C	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	E	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	G	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	I	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	K	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	M	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			
1	O	459	Total	C	N	O	S	0	0	0
			3534	2216	589	695	34			

- Molecule 2 is a protein called Methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			
2	D	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			
2	F	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			
2	H	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			
2	J	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			
2	L	258	Total	C	N	O	S	0	0	0
			1951	1238	311	389	13			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	258	Total 1951	C 1238	N 311	O 389	S 13	0	0	0
2	P	258	Total 1951	C 1238	N 311	O 389	S 13	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total 1	Zn 1	0	0
3	K	1	Total 1	Zn 1	0	0
3	E	2	Total 2	Zn 2	0	0
3	I	2	Total 2	Zn 2	0	0
3	C	1	Total 1	Zn 1	0	0
3	A	2	Total 2	Zn 2	0	0
3	O	1	Total 1	Zn 1	0	0
3	M	2	Total 2	Zn 2	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total 1	K 1	0	0
4	K	1	Total 1	K 1	0	0
4	E	1	Total 1	K 1	0	0
4	I	1	Total 1	K 1	0	0
4	C	1	Total 1	K 1	0	0
4	A	1	Total 1	K 1	0	0
4	O	1	Total 1	K 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	M	1	Total K 1 1	0	0

- Molecule 5 is 5-HYDROXYBENZIMIDAZOLYLCOB(III)AMIDE (three-letter code: B13) (formula: $C_{60}H_{88}CoN_{13}O_{15}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	B	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	D	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	F	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	H	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	J	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	L	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	N	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		
5	P	1	Total	C	Co	N	O	P	0	0
			90	60	1	13	15	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	162	Total 162	O 162	0	0
6	B	36	Total 36	O 36	0	0
6	C	183	Total 183	O 183	0	0
6	D	27	Total 27	O 27	0	0
6	E	20	Total 20	O 20	0	0
6	F	6	Total 6	O 6	0	0
6	G	47	Total 47	O 47	0	0
6	H	1	Total 1	O 1	0	0
6	I	93	Total 93	O 93	0	0
6	J	15	Total 15	O 15	0	0
6	K	99	Total 99	O 99	0	0
6	L	18	Total 18	O 18	0	0
6	M	95	Total 95	O 95	0	0
6	N	18	Total 18	O 18	0	0
6	O	111	Total 111	O 111	0	0
6	P	15	Total 15	O 15	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methyltransferase 1

Chain A:




- Molecule 1: Methyltransferase 1

Chain C:




- Molecule 1: Methyltransferase 1

Chain E:




- Molecule 1: Methyltransferase 1

Chain G:




• Molecule 1: Methyltransferase 1

Chain I:




• Molecule 1: Methyltransferase 1

Chain K:




• Molecule 1: Methyltransferase 1

Chain M:




• Molecule 1: Methyltransferase 1

Chain O: 




• Molecule 2: Methyltransferase 1

Chain B: 




• Molecule 2: Methyltransferase 1

Chain D: 




• Molecule 2: Methyltransferase 1


Chain F: 




- Molecule 2: Methyltransferase 1

Chain H: 


- Molecule 2: Methyltransferase 1


Chain J: 

- Molecule 2: Methyltransferase 1

Chain L: 


- Molecule 2: Methyltransferase 1

Chain N: 



● Molecule 2: Methyltransferase 1

Chain P:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.75Å 172.85Å 190.54Å 90.00° 98.86° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 20.02 – 2.51	Depositor EDS
% Data completeness (in resolution range)	98.5 (20.00-2.50) 98.5 (20.02-2.51)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.182 , 0.231 0.182 , 0.230	Depositor DCC
R_{free} test set	10921 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 218403 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	45566	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, K, B13

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.72	1/3595 (0.0%)	0.78	5/4852 (0.1%)
1	C	0.70	0/3595	0.78	5/4852 (0.1%)
1	E	0.51	0/3595	0.62	0/4852
1	G	0.54	0/3595	0.64	1/4852 (0.0%)
1	I	0.60	0/3595	0.70	3/4852 (0.1%)
1	K	0.60	0/3595	0.68	2/4852 (0.0%)
1	M	0.60	0/3595	0.69	2/4852 (0.0%)
1	O	0.62	0/3595	0.72	6/4852 (0.1%)
2	B	0.48	0/1980	0.66	1/2682 (0.0%)
2	D	0.47	0/1980	0.65	0/2682
2	F	0.46	0/1980	0.59	0/2682
2	H	0.45	0/1980	0.59	0/2682
2	J	0.47	0/1980	0.63	1/2682 (0.0%)
2	L	0.46	0/1980	0.60	0/2682
2	N	0.48	0/1980	0.61	0/2682
2	P	0.48	0/1980	0.65	0/2682
All	All	0.57	1/44600 (0.0%)	0.68	26/60272 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	GLU	CG-CD	5.18	1.59	1.51

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	361	ARG	NE-CZ-NH2	-7.96	116.32	120.30
1	A	361	ARG	NE-CZ-NH2	-7.96	116.32	120.30
1	C	353	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	M	123	ARG	NE-CZ-NH2	-7.65	116.48	120.30
1	I	361	ARG	NE-CZ-NH2	-6.96	116.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3534	0	3462	49	0
1	C	3534	0	3462	56	0
1	E	3534	0	3462	54	0
1	G	3534	0	3462	63	0
1	I	3534	0	3462	51	0
1	K	3534	0	3462	61	0
1	M	3534	0	3462	54	0
1	O	3534	0	3462	57	0
2	B	1951	0	1952	64	0
2	D	1951	0	1952	55	0
2	F	1951	0	1952	54	0
2	H	1951	0	1952	52	0
2	J	1951	0	1952	53	0
2	L	1951	0	1952	49	0
2	N	1951	0	1952	50	0
2	P	1951	0	1952	47	0
3	A	2	0	0	0	0
3	C	1	0	0	0	0
3	E	2	0	0	0	0
3	G	1	0	0	0	0
3	I	2	0	0	0	0
3	K	1	0	0	0	0
3	M	2	0	0	0	0
3	O	1	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	G	1	0	0	0	0
4	I	1	0	0	0	0
4	K	1	0	0	0	0
4	M	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	O	1	0	0	0	0
5	B	90	0	81	9	0
5	D	90	0	81	13	0
5	F	90	0	81	7	0
5	H	90	0	81	10	0
5	J	90	0	81	13	0
5	L	90	0	81	10	0
5	N	90	0	81	11	0
5	P	90	0	81	9	0
6	A	162	0	0	4	0
6	B	36	0	0	1	0
6	C	183	0	0	3	0
6	D	27	0	0	0	0
6	E	20	0	0	0	0
6	F	6	0	0	0	0
6	G	47	0	0	3	0
6	H	1	0	0	0	0
6	I	93	0	0	3	0
6	J	15	0	0	0	0
6	K	99	0	0	4	0
6	L	18	0	0	1	0
6	M	95	0	0	4	0
6	N	18	0	0	0	0
6	O	111	0	0	4	0
6	P	15	0	0	0	0
All	All	45566	0	43960	865	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 865 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:P:233:ASP:O	2:P:237:ILE:HG22	1.56	1.05
1:M:353:ARG:HA	1:M:356:MET:CE	1.92	0.99
1:K:353:ARG:HA	1:K:356:MET:CE	1.95	0.97
2:L:134:ASP:HB2	2:L:187:THR:HG21	1.46	0.94
2:B:17:ASN:HD22	2:B:17:ASN:H	1.14	0.91

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/461 (99%)	442 (97%)	15 (3%)	0	100	100
1	C	457/461 (99%)	440 (96%)	17 (4%)	0	100	100
1	E	457/461 (99%)	433 (95%)	21 (5%)	3 (1%)	30	50
1	G	457/461 (99%)	434 (95%)	22 (5%)	1 (0%)	56	79
1	I	457/461 (99%)	439 (96%)	17 (4%)	1 (0%)	56	79
1	K	457/461 (99%)	439 (96%)	18 (4%)	0	100	100
1	M	457/461 (99%)	439 (96%)	18 (4%)	0	100	100
1	O	457/461 (99%)	439 (96%)	18 (4%)	0	100	100
2	B	256/258 (99%)	239 (93%)	13 (5%)	4 (2%)	14	23
2	D	256/258 (99%)	224 (88%)	28 (11%)	4 (2%)	14	23
2	F	256/258 (99%)	227 (89%)	26 (10%)	3 (1%)	19	32
2	H	256/258 (99%)	235 (92%)	18 (7%)	3 (1%)	19	32
2	J	256/258 (99%)	237 (93%)	14 (6%)	5 (2%)	11	17
2	L	256/258 (99%)	235 (92%)	15 (6%)	6 (2%)	10	14
2	N	256/258 (99%)	229 (90%)	23 (9%)	4 (2%)	14	23
2	P	256/258 (99%)	240 (94%)	13 (5%)	3 (1%)	19	32
All	All	5704/5752 (99%)	5371 (94%)	296 (5%)	37 (1%)	33	55

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	256	HIS
2	F	256	HIS
2	H	256	HIS
2	L	37	ASP
2	L	256	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/374 (100%)	344 (92%)	29 (8%)	18	32
1	C	373/374 (100%)	344 (92%)	29 (8%)	18	32
1	E	373/374 (100%)	339 (91%)	34 (9%)	14	24
1	G	373/374 (100%)	346 (93%)	27 (7%)	21	36
1	I	373/374 (100%)	345 (92%)	28 (8%)	19	34
1	K	373/374 (100%)	348 (93%)	25 (7%)	23	40
1	M	373/374 (100%)	344 (92%)	29 (8%)	18	32
1	O	373/374 (100%)	343 (92%)	30 (8%)	17	31
2	B	206/206 (100%)	183 (89%)	23 (11%)	9	15
2	D	206/206 (100%)	192 (93%)	14 (7%)	22	39
2	F	206/206 (100%)	189 (92%)	17 (8%)	16	29
2	H	206/206 (100%)	190 (92%)	16 (8%)	18	32
2	J	206/206 (100%)	190 (92%)	16 (8%)	18	32
2	L	206/206 (100%)	190 (92%)	16 (8%)	18	32
2	N	206/206 (100%)	189 (92%)	17 (8%)	16	29
2	P	206/206 (100%)	184 (89%)	22 (11%)	10	17
All	All	4632/4640 (100%)	4260 (92%)	372 (8%)	17	31

5 of 372 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	370	LEU
1	I	325	THR
1	O	347	LYS
2	H	9	LEU
2	H	256	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 112 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	195	ASN
2	J	17	ASN
1	O	156	ASN
2	H	216	GLN
1	I	124	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 28 ligands modelled in this entry, 20 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	B13	B	500	2	100,100,100	2.05	19 (19%)	150,164,164	1.93	29 (19%)
5	B13	D	500	2	100,100,100	2.12	23 (23%)	150,164,164	1.98	36 (24%)
5	B13	F	500	2	100,100,100	2.10	22 (22%)	150,164,164	1.96	39 (26%)
5	B13	H	500	2	100,100,100	2.15	19 (19%)	150,164,164	1.99	32 (21%)
5	B13	J	500	2,6	100,100,100	2.02	19 (19%)	150,164,164	1.93	31 (20%)
5	B13	L	500	2,6	100,100,100	2.10	22 (22%)	150,164,164	1.98	40 (26%)
5	B13	N	500	2	100,100,100	2.04	19 (19%)	150,164,164	2.00	36 (24%)
5	B13	P	500	2	100,100,100	2.01	20 (20%)	150,164,164	2.01	39 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	B13	B	500	2	-	0/56/223/223	0/1/11/11
5	B13	D	500	2	-	0/56/223/223	0/1/11/11
5	B13	F	500	2	-	0/56/223/223	0/1/11/11
5	B13	H	500	2	-	0/56/223/223	0/1/11/11
5	B13	J	500	2,6	-	0/56/223/223	0/1/11/11
5	B13	L	500	2,6	-	0/56/223/223	0/1/11/11
5	B13	N	500	2	-	0/56/223/223	0/1/11/11
5	B13	P	500	2	-	0/56/223/223	0/1/11/11

The worst 5 of 163 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	500	B13	C15-C14	10.32	1.41	1.33
5	D	500	B13	C15-C14	10.00	1.41	1.33
5	F	500	B13	C15-C14	9.65	1.41	1.33
5	L	500	B13	C15-C14	9.31	1.40	1.33
5	N	500	B13	C15-C14	8.70	1.40	1.33

The worst 5 of 282 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	N	500	B13	C2B-N1B-C1R	-8.58	109.48	126.38
5	F	500	B13	O6R-C4R-C5R	8.08	126.41	109.15
5	N	500	B13	O6R-C4R-C5R	8.06	126.38	109.15
5	B	500	B13	O6R-C4R-C5R	8.02	126.29	109.15
5	H	500	B13	C2B-N1B-C1R	-7.81	111.01	126.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	459/461 (99%)	-0.58	0	100	100	3, 11, 26, 38	0
1	C	459/461 (99%)	-0.61	0	100	100	3, 10, 23, 39	0
1	E	459/461 (99%)	0.11	14 (3%)	47	48	25, 33, 42, 49	0
1	G	459/461 (99%)	-0.21	6 (1%)	74	76	18, 29, 40, 49	0
1	I	459/461 (99%)	-0.47	3 (0%)	84	86	9, 20, 34, 50	0
1	K	459/461 (99%)	-0.47	3 (0%)	84	86	12, 20, 33, 46	0
1	M	459/461 (99%)	-0.47	2 (0%)	90	92	10, 20, 34, 48	0
1	O	459/461 (99%)	-0.50	1 (0%)	93	94	7, 18, 32, 45	0
2	B	258/258 (100%)	0.92	44 (17%)	2	2	4, 67, 100, 107	0
2	D	258/258 (100%)	1.00	51 (19%)	2	1	5, 66, 100, 110	0
2	F	258/258 (100%)	1.30	67 (25%)	1	1	18, 72, 102, 112	0
2	H	258/258 (100%)	1.63	82 (31%)	1	1	28, 73, 104, 110	0
2	J	258/258 (100%)	0.90	37 (14%)	3	3	17, 64, 97, 107	0
2	L	258/258 (100%)	1.23	63 (24%)	1	1	12, 71, 102, 113	0
2	N	258/258 (100%)	1.73	109 (42%)	1	0	12, 73, 103, 110	0
2	P	258/258 (100%)	0.96	39 (15%)	3	3	17, 67, 98, 108	0
All	All	5736/5752 (99%)	0.18	521 (9%)	10	8	3, 29, 92, 113	0

The worst 5 of 521 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	258	HIS	7.6
2	J	207	PHE	7.2
2	P	207	PHE	7.1
2	J	178	LEU	7.1
2	H	207	PHE	7.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	K	K	516	1/1	0.14	1.88	31,31,31,31	1
4	K	I	515	1/1	0.12	1.12	28,28,28,28	1
4	K	G	514	1/1	0.13	0.79	39,39,39,39	1
4	K	O	518	1/1	0.10	0.17	35,35,35,35	1
5	B13	N	500	90/90	0.18	0.11	60,66,72,73	0
5	B13	L	500	90/90	0.15	-0.23	41,47,58,59	0
5	B13	F	500	90/90	0.17	-0.26	53,63,71,72	0
5	B13	P	500	90/90	0.14	-0.41	31,40,45,52	0
5	B13	D	500	90/90	0.13	-0.44	20,30,40,42	0
5	B13	H	500	90/90	0.14	-0.50	45,58,68,69	0
5	B13	J	500	90/90	0.13	-0.55	32,37,50,57	0
5	B13	B	500	90/90	0.11	-0.60	9,24,48,49	0
4	K	C	512	1/1	0.07	-0.92	22,22,22,22	1
3	ZN	I	521	1/1	0.04	-1.98	16,16,16,16	0
3	ZN	O	508	1/1	0.05	-1.99	33,33,33,33	0
3	ZN	E	522	1/1	0.05	-2.11	37,37,37,37	0
4	K	A	511	1/1	0.04	-2.39	23,23,23,23	1
3	ZN	A	524	1/1	0.04	-2.39	9,9,9,9	0
3	ZN	G	504	1/1	0.07	-2.49	33,33,33,33	1
3	ZN	I	505	1/1	0.04	-2.54	30,30,30,30	0
3	ZN	K	506	1/1	0.04	-2.70	33,33,33,33	0
3	ZN	M	523	1/1	0.03	-2.84	21,21,21,21	0
4	K	M	517	1/1	0.07	-3.05	33,33,33,33	1
3	ZN	A	501	1/1	0.03	-3.09	20,20,20,20	0
4	K	E	513	1/1	0.10	-3.14	33,33,33,33	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	C	502	1/1	0.03	-3.28	19,19,19,19	0
3	ZN	M	507	1/1	0.03	-4.18	33,33,33,33	0
3	ZN	E	503	1/1	0.05	-6.12	36,36,36,36	1

6.5 Other polymers ⓘ

There are no such residues in this entry.