



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 08:04 AM GMT

PDB ID : 4I2V  
Title : X-ray structure of the unliganded uridine phosphorylase from Yersinia pseudotuberculosis at 2.12Å resolution  
Authors : Lashkov, A.A.; Balaev, V.V.; Prokofev, I.I.; Betzel, C.; Gabdoulkhakov, A.G.; Mikhailov, A.M.  
Deposited on : 2012-11-23  
Resolution : 2.12 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3409 (2.14-2.10)
Clashscore	79885	4090 (2.14-2.10)
Ramachandran outliers	78287	4048 (2.14-2.10)
Sidechain outliers	78261	4049 (2.14-2.10)
RSRZ outliers	66119	3410 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	253	
1	B	253	
1	C	253	
1	D	253	
1	E	253	
1	F	253	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11322 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			
1	B	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			
1	C	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			
1	D	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			
1	E	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			
1	F	250	Total	C	N	O	S	0	0	0
			1875	1173	330	360	12			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	O	0	0
			12	12		
2	B	17	Total	O	0	0
			17	17		
2	C	13	Total	O	0	0
			13	13		
2	D	17	Total	O	0	0
			17	17		
2	E	4	Total	O	0	0
			4	4		
2	F	9	Total	O	0	0
			9	9		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Uridine phosphorylase

Chain A:



#### • Molecule 1: Uridine phosphorylase


Chain B:



#### • Molecule 1: Uridine phosphorylase


Chain C:






• Molecule 1: Uridine phosphorylase

Chain D:




• Molecule 1: Uridine phosphorylase


Chain E:



• Molecule 1: Uridine phosphorylase

Chain F:





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.68Å 156.68Å 48.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.61 – 2.12 29.61 – 2.12	Depositor EDS
% Data completeness (in resolution range)	97.6 (29.61-2.12) 97.7 (29.61-2.12)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 2.12Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.200 , 0.242 0.200 , 0.242	Depositor DCC
$R_{free}$ test set	3680 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.6	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 15.2	EDS
Estimated twinning fraction	0.460 for -h,-k,l 0.427 for -h,-k,l 0.075 for h,-h-k,-l 0.074 for -k,-h,-l	Xtriage
Reported twinning fraction	0.460 for -h,-k,l	Depositor
L-test for twinning	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 73600 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11322	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	1/1908 (0.1%)	0.67	0/2590
1	B	0.45	0/1908	0.69	0/2590
1	C	0.43	0/1908	0.69	0/2590
1	D	0.44	0/1908	0.67	0/2590
1	E	0.41	0/1908	0.69	0/2590
1	F	0.43	0/1908	0.66	0/2590
All	All	0.44	1/11448 (0.0%)	0.68	0/15540

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	136	CYS	CB-SG	-5.05	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	232	GLU	Peptide
1	E	232	GLU	Peptide



## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1875	0	1869	117	0
1	B	1875	0	1869	104	0
1	C	1875	0	1869	113	0
1	D	1875	0	1869	119	0
1	E	1875	0	1869	97	0
1	F	1875	0	1869	94	0
2	A	12	0	0	1	0
2	B	17	0	0	0	0
2	C	13	0	0	1	0
2	D	17	0	0	4	0
2	E	4	0	0	0	0
2	F	9	0	0	1	0
All	All	11322	0	11214	595	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 595 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:122:HIS:HB3	1:B:164:PRO:HB2	1.43	1.00
1:F:16:LEU:HG	1:F:63:ILE:HG13	1.48	0.93
1:D:49:GLU:HG2	1:D:68:GLY:HA2	1.55	0.88
1:E:167:GLU:HG3	1:E:183:SER:HB2	1.54	0.87
1:E:164:PRO:HB2	1:F:122:HIS:HB3	1.57	0.85

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/253 (98%)	228 (92%)	20 (8%)	0	100	100
1	B	248/253 (98%)	236 (95%)	11 (4%)	1 (0%)	43	39
1	C	248/253 (98%)	227 (92%)	17 (7%)	4 (2%)	14	6
1	D	248/253 (98%)	229 (92%)	18 (7%)	1 (0%)	43	39
1	E	248/253 (98%)	228 (92%)	18 (7%)	2 (1%)	27	19
1	F	248/253 (98%)	233 (94%)	15 (6%)	0	100	100
All	All	1488/1518 (98%)	1381 (93%)	99 (7%)	8 (0%)	38	33

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	126	MET
1	C	221	VAL
1	C	116	LEU
1	B	176	VAL
1	C	176	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/203 (99%)	179 (89%)	22 (11%)	9	5
1	B	201/203 (99%)	173 (86%)	28 (14%)	5	2
1	C	201/203 (99%)	174 (87%)	27 (13%)	6	3
1	D	201/203 (99%)	174 (87%)	27 (13%)	6	3
1	E	201/203 (99%)	173 (86%)	28 (14%)	5	2
1	F	201/203 (99%)	181 (90%)	20 (10%)	11	6
All	All	1206/1218 (99%)	1054 (87%)	152 (13%)	7	3

5 of 152 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	172	PHE
1	D	54	ARG
1	F	158	SER
1	C	226	GLN
1	D	9	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	166	GLN
1	F	152	HIS
1	D	194	ASN
1	B	225	GLN
1	D	8	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/253 (98%)	0.22	4 (1%) 68 72	17, 31, 50, 61	0
1	B	250/253 (98%)	0.10	5 (2%) 62 66	16, 30, 44, 55	0
1	C	250/253 (98%)	0.11	4 (1%) 68 72	18, 30, 46, 53	0
1	D	250/253 (98%)	0.10	5 (2%) 62 66	16, 31, 46, 58	0
1	E	250/253 (98%)	0.24	10 (4%) 36 40	20, 32, 53, 66	0
1	F	250/253 (98%)	0.25	7 (2%) 50 55	19, 34, 50, 59	0
All	All	1500/1518 (98%)	0.17	35 (2%) 57 62	16, 32, 50, 66	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	84	LEU	4.7
1	E	169	TYR	4.1
1	A	4	SER	3.8
1	B	229	PRO	3.6
1	D	231	GLU	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.