



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 12:21 PM GMT

PDB ID : 1I36  
Title : Structure of Conserved Protein MTH1747 of Unknown Function Reveals Structural Similarity with 3-Hydroxyacid Dehydrogenases  
Authors : Korolev, S.V.; Dementieva, I.S.; Christendat, D.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2001-02-13  
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	264	
1	B	264	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4443 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CONSERVED HYPOTHETICAL PROTEIN MTH1747.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			1963	1218	361	373	11			
1	B	258	Total	C	N	O	S	0	0	0
			1963	1218	361	373	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	LEU	MET	ENGINEERED	UNP O27779
B	1	LEU	MET	ENGINEERED	UNP O27779

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	219	Total	O	0	0
			219	219		
3	B	202	Total	O	0	0
			202	202		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: CONSERVED HYPOTHETICAL PROTEIN MTH1747

Chain A: 



#### • Molecule 1: CONSERVED HYPOTHETICAL PROTEIN MTH1747

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.87Å 123.46Å 133.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.81 – 2.00 29.43 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.9 (23.81-2.00) 95.9 (29.43-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.97 (at 2.00Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.203 , 0.230 0.202 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	24.1	Xtriage
Anisotropy	0.488	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 51211 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4443	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.18 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7643e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/1985	0.70	0/2670
1	B	0.56	0/1985	0.69	1/2670 (0.0%)
All	All	0.57	0/3970	0.69	1/5340 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	130	ARG	NE-CZ-NH1	-6.29	117.16	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1963	0	1988	26	0
1	B	1963	0	1988	46	0
2	A	48	0	22	2	0
2	B	48	0	24	4	0
3	A	219	0	0	1	0
3	B	202	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4443	0	4022	66	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (66) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:220:TYR:CE2	1:B:243:ILE:HG23	2.10	0.87
1:A:196:THR:HG21	1:B:162:ARG:HD2	1.56	0.86
1:B:160:MET:SD	1:B:229:MET:HG2	2.21	0.80
1:B:13:GLN:NE2	1:B:40:ARG:HH11	1.83	0.76
1:A:114:GLY:HA3	1:A:123:ILE:HD11	1.75	0.68
1:A:192:MET:O	1:A:196:THR:HG22	1.93	0.67
1:B:170:SER:OG	1:B:208:ARG:HD2	1.95	0.66
1:B:13:GLN:HE21	1:B:40:ARG:HH11	1.45	0.62
1:B:89:ASN:HB2	2:B:1750:NAP:O2D	2.00	0.61
1:A:117:ARG:HG2	1:A:117:ARG:HH11	1.65	0.60
1:A:199:ASN:O	1:A:201:PHE:N	2.34	0.60
1:A:27:VAL:HG12	1:A:47:THR:HB	1.84	0.60
1:B:174:TRP:O	1:B:178:THR:HG23	2.02	0.58
1:B:150:ARG:HH11	1:B:150:ARG:CB	2.17	0.58
1:A:199:ASN:C	1:A:201:PHE:H	2.08	0.57
1:B:150:ARG:HH11	1:B:150:ARG:HB2	1.69	0.56
1:A:26:VAL:HG23	1:A:46:VAL:HG12	1.88	0.55
1:B:163:SER:OG	1:B:167:LYS:NZ	2.39	0.55
1:B:137:LYS:O	1:B:140:ARG:HG3	2.06	0.55
1:B:16:ALA:HB1	1:B:26:VAL:HG11	1.88	0.55
1:B:114:GLY:CA	1:B:123:ILE:HD11	2.37	0.54
1:B:9:GLY:HA3	2:B:1750:NAP:O5B	2.07	0.54
1:A:227:GLN:OE1	1:A:239:PRO:HB2	2.08	0.54
1:B:192:MET:O	1:B:195:TYR:HB2	2.09	0.53
1:B:119:LYS:HD2	1:B:119:LYS:N	2.23	0.53
1:B:160:MET:SD	1:B:229:MET:CG	2.96	0.53
1:B:112:ILE:HG23	1:B:123:ILE:HG12	1.92	0.52
1:B:13:GLN:NE2	1:B:40:ARG:NH1	2.57	0.52
1:B:196:THR:HG23	1:B:197:GLU:OE2	2.09	0.52
1:B:89:ASN:OD1	2:B:1750:NAP:H6N	2.10	0.52
1:A:182:ARG:HB2	1:B:233:VAL:CG1	2.41	0.51
1:B:114:GLY:C	1:B:123:ILE:HD11	2.31	0.50
1:A:9:GLY:HA3	2:A:1350:NAP:O5B	2.11	0.50
1:B:174:TRP:CZ2	1:B:178:THR:HG21	2.47	0.49
1:B:229:MET:HG3	1:B:230:LEU:N	2.27	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:233:VAL:HG12	1:B:234:ILE:HG23	1.93	0.49
1:B:114:GLY:HA3	1:B:123:ILE:HD11	1.95	0.48
1:B:35:PRO:HG2	3:B:1799:HOH:O	2.13	0.48
1:A:206:ILE:HG13	1:A:258:ALA:HB2	1.95	0.48
1:A:114:GLY:CA	1:A:123:ILE:HD11	2.44	0.48
1:B:213:CYS:SG	1:B:249:LEU:HD22	2.54	0.47
1:A:217:ARG:NH1	1:A:217:ARG:HB2	2.29	0.47
1:B:220:TYR:HE2	1:B:243:ILE:HG23	1.77	0.46
1:B:1:LEU:HD13	1:B:58:PRO:HG3	1.96	0.46
1:A:167:LYS:HE2	1:A:222:GLU:HB3	1.98	0.45
1:A:182:ARG:CB	1:B:233:VAL:HG13	2.47	0.45
1:A:1:LEU:HD12	1:A:58:PRO:HG3	1.99	0.44
1:A:246:PHE:CD2	1:A:249:LEU:HD12	2.52	0.44
1:B:174:TRP:CE2	1:B:178:THR:HG21	2.52	0.44
1:A:195:TYR:CZ	1:B:124:ARG:HD2	2.52	0.44
1:A:196:THR:CG2	1:B:162:ARG:HD2	2.37	0.44
1:B:34:SER:HB2	1:B:35:PRO:HD2	1.99	0.43
1:A:182:ARG:HH22	1:A:264:ALA:HB3	1.84	0.43
1:A:227:GLN:HE22	1:A:236:PRO:HG2	1.84	0.43
1:A:26:VAL:CG2	1:A:46:VAL:HG12	2.48	0.43
1:A:182:ARG:HB2	1:B:233:VAL:HG13	2.00	0.42
1:B:151:GLU:HB2	1:B:152:PRO:HD2	2.01	0.42
1:A:64:VAL:O	2:A:1350:NAP:H52N	2.18	0.42
1:B:109:ASP:O	1:B:127:ALA:HA	2.19	0.42
1:B:222:GLU:O	1:B:226:VAL:HG13	2.20	0.41
1:B:68:VAL:O	1:B:68:VAL:HG12	2.21	0.41
1:B:8:PHE:HA	1:B:12:ALA:HB3	2.02	0.41
1:B:64:VAL:O	2:B:1750:NAP:H52N	2.20	0.41
1:B:117:ARG:HB2	3:B:1951:HOH:O	2.21	0.41
1:B:130:ARG:NH1	1:B:130:ARG:HB2	2.36	0.41
1:A:242:ILE:HD13	3:A:1561:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/264 (96%)	236 (93%)	13 (5%)	5 (2%)	11	4
1	B	254/264 (96%)	242 (95%)	11 (4%)	1 (0%)	43	36
All	All	508/528 (96%)	478 (94%)	24 (5%)	6 (1%)	19	9

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	ASN
1	A	200	ASP
1	A	201	PHE
1	A	64	VAL
1	A	198	GLY
1	B	64	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/217 (94%)	203 (99%)	2 (1%)	85	88
1	B	205/217 (94%)	200 (98%)	5 (2%)	61	61
All	All	410/434 (94%)	403 (98%)	7 (2%)	73	75

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LEU
1	A	196	THR
1	B	104	LYS
1	B	150	ARG
1	B	226	VAL
1	B	229	MET
1	B	232	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	HIS
1	B	13	GLN
1	B	181	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	1350	-	52,52,52	2.89	20 (38%)	80,80,80	3.65	29 (36%)
2	NAP	B	1750	-	52,52,52	2.72	20 (38%)	80,80,80	3.63	27 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	1350	-	-	0/35/67/67	0/3/5/5
2	NAP	B	1750	-	-	0/35/67/67	0/3/5/5

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1350	NAP	C3B-C4B	-10.44	1.24	1.53
2	B	1750	NAP	C3B-C4B	-9.91	1.25	1.53
2	A	1350	NAP	O4B-C4B	7.40	1.62	1.45
2	A	1350	NAP	C2B-C1B	-6.52	1.41	1.52
2	B	1750	NAP	O4B-C4B	6.46	1.60	1.45
2	B	1750	NAP	C2B-C1B	-6.30	1.41	1.52
2	A	1350	NAP	O2B-C2B	-5.53	1.26	1.44
2	B	1750	NAP	O2B-C2B	-4.98	1.28	1.44
2	A	1350	NAP	PN-O3	4.91	1.70	1.60
2	B	1750	NAP	C2N-N1N	4.85	1.41	1.35
2	A	1350	NAP	O4B-C1B	4.32	1.48	1.41
2	A	1350	NAP	C2N-N1N	4.14	1.40	1.35
2	A	1350	NAP	PA-O3	4.11	1.67	1.59
2	B	1750	NAP	O4B-C1B	3.87	1.47	1.41
2	B	1750	NAP	C2N-C3N	3.87	1.44	1.38
2	A	1350	NAP	C2N-C3N	3.61	1.44	1.38
2	B	1750	NAP	PN-O3	3.60	1.67	1.60
2	B	1750	NAP	PA-O3	3.41	1.66	1.59
2	A	1350	NAP	C4A-N3A	3.16	1.40	1.35
2	B	1750	NAP	P2B-O2X	-3.05	1.43	1.54
2	A	1350	NAP	P2B-O2X	-2.87	1.44	1.54
2	B	1750	NAP	C4A-N3A	2.79	1.39	1.35
2	A	1350	NAP	C3N-C7N	-2.73	1.45	1.50
2	A	1350	NAP	P2B-O3X	-2.58	1.45	1.54
2	B	1750	NAP	P2B-O3X	-2.57	1.45	1.54
2	B	1750	NAP	C5B-C4B	-2.50	1.43	1.51
2	A	1350	NAP	O3B-C3B	-2.48	1.37	1.43
2	A	1350	NAP	C8A-N9A	2.47	1.40	1.36
2	B	1750	NAP	C3N-C7N	-2.47	1.46	1.50
2	A	1350	NAP	O5B-C5B	2.46	1.55	1.44
2	B	1750	NAP	C3B-C2B	-2.45	1.47	1.53
2	B	1750	NAP	C6N-N1N	2.45	1.42	1.35
2	B	1750	NAP	C8A-N9A	2.29	1.40	1.36
2	A	1350	NAP	P2B-O2B	2.24	1.66	1.59
2	B	1750	NAP	O3B-C3B	-2.24	1.37	1.43
2	A	1350	NAP	PA-O2A	-2.18	1.45	1.55
2	A	1350	NAP	C6N-N1N	2.12	1.41	1.35
2	B	1750	NAP	O5B-C5B	2.10	1.53	1.44
2	A	1350	NAP	C3B-C2B	-2.06	1.48	1.53
2	B	1750	NAP	PA-O2A	-2.05	1.45	1.55

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1350	NAP	C2B-C3B-C4B	17.59	143.69	101.94
2	B	1750	NAP	C2B-C3B-C4B	16.69	141.56	101.94
2	B	1750	NAP	O4B-C4B-C3B	-10.99	82.89	105.17
2	B	1750	NAP	O3B-C3B-C4B	-10.65	79.71	111.08
2	A	1350	NAP	O4B-C4B-C3B	-10.25	84.39	105.17
2	A	1350	NAP	O3B-C3B-C4B	-10.11	81.30	111.08
2	B	1750	NAP	C3B-C2B-C1B	-8.40	86.40	102.73
2	A	1350	NAP	C3B-C2B-C1B	-8.24	86.70	102.73
2	B	1750	NAP	O4D-C1D-N1N	7.75	115.88	107.95
2	A	1350	NAP	O4B-C1B-C2B	7.55	114.00	106.95
2	A	1350	NAP	O4D-C1D-N1N	7.13	115.25	107.95
2	B	1750	NAP	O4B-C1B-C2B	6.40	112.93	106.95
2	A	1350	NAP	O2B-C2B-C3B	6.11	135.45	111.54
2	B	1750	NAP	C8A-N9A-C1B	5.88	137.96	126.38
2	B	1750	NAP	C1B-N9A-C4A	-5.87	116.48	126.64
2	A	1350	NAP	C8A-N9A-C1B	5.78	137.76	126.38
2	A	1350	NAP	C1B-N9A-C4A	-5.63	116.91	126.64
2	A	1350	NAP	N3A-C2A-N1A	-5.19	124.37	128.71
2	B	1750	NAP	O3-PA-O1A	-4.96	75.66	111.28
2	A	1350	NAP	O3-PA-O1A	-4.94	75.79	111.28
2	B	1750	NAP	N3A-C2A-N1A	-4.92	124.59	128.71
2	B	1750	NAP	O2B-C2B-C3B	4.55	129.35	111.54
2	A	1350	NAP	P2B-O2B-C2B	-4.53	112.41	121.96
2	B	1750	NAP	O4B-C1B-N9A	4.52	112.65	108.44
2	B	1750	NAP	O2A-PA-O1A	4.52	137.45	112.21
2	B	1750	NAP	P2B-O2B-C2B	-4.33	112.85	121.96
2	A	1350	NAP	O4B-C1B-N9A	4.32	112.45	108.44
2	A	1350	NAP	O2A-PA-O1A	4.19	135.62	112.21
2	B	1750	NAP	C5B-C4B-C3B	4.15	131.85	115.21
2	B	1750	NAP	O4B-C4B-C5B	-4.07	94.84	109.36
2	A	1350	NAP	O4B-C4B-C5B	-3.90	95.45	109.36
2	A	1350	NAP	O2A-PA-O3	-3.82	87.02	105.14
2	B	1750	NAP	O2A-PA-O3	-3.81	87.05	105.14
2	A	1350	NAP	O7N-C7N-N7N	-3.58	117.42	122.59
2	B	1750	NAP	O5B-C5B-C4B	-3.54	95.94	108.94
2	A	1350	NAP	C3N-C7N-N7N	3.52	121.78	117.77
2	A	1350	NAP	C5B-C4B-C3B	3.45	129.04	115.21
2	B	1750	NAP	O7N-C7N-N7N	-3.22	117.94	122.59
2	B	1750	NAP	C3N-C7N-N7N	3.20	121.42	117.77
2	A	1350	NAP	O5B-C5B-C4B	-3.12	97.49	108.94
2	B	1750	NAP	O3-PA-O5B	-3.08	89.61	103.41
2	B	1750	NAP	C3N-C2N-N1N	-2.94	115.40	120.36
2	A	1350	NAP	O3-PA-O5B	-2.89	90.46	103.41

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1350	NAP	O2X-P2B-O2B	-2.76	99.13	107.09
2	A	1350	NAP	C3N-C2N-N1N	-2.72	115.77	120.36
2	A	1350	NAP	O3-PN-O5D	2.59	111.75	101.36
2	A	1350	NAP	O3X-P2B-O2X	2.58	117.67	107.61
2	B	1750	NAP	O3-PN-O5D	2.55	111.58	101.36
2	A	1350	NAP	C5N-C4N-C3N	2.50	123.56	120.32
2	B	1750	NAP	C5N-C4N-C3N	2.46	123.52	120.32
2	A	1350	NAP	C4B-O4B-C1B	-2.34	107.21	109.75
2	B	1750	NAP	O2X-P2B-O2B	-2.28	100.53	107.09
2	A	1350	NAP	O2N-PN-O3	-2.20	102.85	108.79
2	B	1750	NAP	O3X-P2B-O2X	2.19	116.14	107.61
2	A	1350	NAP	C8A-N9A-C4A	-2.13	105.27	106.90
2	B	1750	NAP	C4B-O4B-C1B	-2.05	107.53	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/264 (97%)	0.12	8 (3%) 47 46	10, 25, 57, 76	0
1	B	258/264 (97%)	0.25	12 (4%) 30 29	12, 28, 56, 72	0
All	All	516/528 (97%)	0.19	20 (3%) 37 37	10, 27, 57, 76	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	GLY	8.9
1	B	199	ASN	7.0
1	B	264	ALA	6.1
1	A	199	ASN	6.1
1	B	1	LEU	4.9
1	B	195	TYR	4.8
1	A	263	CYS	4.1
1	B	118	ARG	3.4
1	B	217	ARG	3.4
1	A	1	LEU	3.4
1	A	217	ARG	3.2
1	B	263	CYS	2.7
1	A	200	ASP	2.7
1	B	200	ASP	2.5
1	A	117	ARG	2.4
1	B	196	THR	2.3
1	B	198	GLY	2.3
1	A	195	TYR	2.2
1	B	203	GLU	2.1
1	B	117	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAP	B	1750	48/48	0.13	-0.08	23,35,39,44	0
2	NAP	A	1350	48/48	0.11	-0.17	21,29,35,41	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.