



wwPDB X-ray Structure Validation Summary Report i

Mar 27, 2014 – 02:43 PM EDT

PDB ID : 2I3S
Title : Bub3 complex with Bub1 GLEBS motif
Authors : Larsen, N.A.; Harrison, S.C.
Deposited on : 2006-08-20
Resolution : 1.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1439
EDS : stable22978
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22978

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	349	
1	C	349	
1	E	349	
2	B	36	
2	D	36	
2	F	36	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 9442 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell cycle arrest protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	Total 2646	C 1676	N 441	O 516	S 13	0	0	0
1	C	328	Total 2600	C 1651	N 434	O 502	S 13	0	0	0
1	E	332	Total 2631	C 1670	N 441	O 507	S 13	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	GLU	-	CLONING ARTIFACT	UNP P26449
A	343	LEU	-	CLONING ARTIFACT	UNP P26449
A	344	HIS	-	EXPRESSION TAG	UNP P26449
A	345	HIS	-	EXPRESSION TAG	UNP P26449
A	346	HIS	-	EXPRESSION TAG	UNP P26449
A	347	HIS	-	EXPRESSION TAG	UNP P26449
A	348	HIS	-	EXPRESSION TAG	UNP P26449
A	349	HIS	-	EXPRESSION TAG	UNP P26449
C	342	GLU	-	CLONING ARTIFACT	UNP P26449
C	343	LEU	-	CLONING ARTIFACT	UNP P26449
C	344	HIS	-	EXPRESSION TAG	UNP P26449
C	345	HIS	-	EXPRESSION TAG	UNP P26449
C	346	HIS	-	EXPRESSION TAG	UNP P26449
C	347	HIS	-	EXPRESSION TAG	UNP P26449
C	348	HIS	-	EXPRESSION TAG	UNP P26449
C	349	HIS	-	EXPRESSION TAG	UNP P26449
E	342	GLU	-	CLONING ARTIFACT	UNP P26449
E	343	LEU	-	CLONING ARTIFACT	UNP P26449
E	344	HIS	-	EXPRESSION TAG	UNP P26449
E	345	HIS	-	EXPRESSION TAG	UNP P26449
E	346	HIS	-	EXPRESSION TAG	UNP P26449
E	347	HIS	-	EXPRESSION TAG	UNP P26449
E	348	HIS	-	EXPRESSION TAG	UNP P26449

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	349	HIS	-	EXPRESSION TAG	UNP P26449

- Molecule 2 is a protein called Checkpoint serine/threonine-proteinkinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	36	Total	C	N	O	S	0	0	0
			307	200	47	59	1			
2	D	36	Total	C	N	O	S	0	0	0
			307	200	47	59	1			
2	F	36	Total	C	N	O	S	0	0	0
			307	200	47	59	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	264	Total	O	0	0
			264	264		
3	B	21	Total	O	0	0
			21	21		
3	C	188	Total	O	0	0
			188	188		
3	D	27	Total	O	0	0
			27	27		
3	E	136	Total	O	0	0
			136	136		
3	F	8	Total	O	0	0
			8	8		




- Molecule 2: Checkpoint serine/threonine-proteinkinase

Chain B:



- Molecule 2: Checkpoint serine/threonine-proteinkinase

Chain D:



- Molecule 2: Checkpoint serine/threonine-proteinkinase

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.66Å 79.18Å 142.58Å 90.00° 98.34° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 49.55 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.90) 85.2 (49.55-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.90Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.219 , 0.262 0.218 , 0.261	Depositor DCC
R_{free} test set	3810 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.764	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.1	EDS
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 75104 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9442	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2692	0.68	0/3644
1	C	0.34	0/2644	0.64	0/3576
1	E	0.32	0/2676	0.63	0/3621
2	B	0.38	0/313	0.53	0/420
2	D	0.42	0/313	0.60	0/420
2	F	0.34	0/313	0.53	0/420
All	All	0.34	0/8951	0.64	0/12101

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2646	0	2618	63	0
1	C	2600	0	2591	90	0
1	E	2631	0	2629	105	0
2	B	307	0	301	10	0
2	D	307	0	301	19	0
2	F	307	0	301	8	0
3	A	264	0	0	4	0
3	B	21	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	188	0	0	5	0
3	D	27	0	0	0	0
3	E	136	0	0	4	0
3	F	8	0	0	1	0
All	All	9442	0	8741	278	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:329:GLU:HG2	2:D:330:ASN:H	1.24	1.01
1:E:239:ARG:HG2	1:E:242:ARG:HH21	1.27	0.99
1:C:86:LEU:H	1:C:86:LEU:HD23	1.32	0.94
1:E:47:VAL:HG11	1:E:335:ILE:HD12	1.51	0.93
1:A:322:ILE:H	1:A:322:ILE:HD13	1.36	0.91

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/349 (95%)	311 (94%)	15 (4%)	4 (1%)	19	6
1	C	322/349 (92%)	304 (94%)	17 (5%)	1 (0%)	50	37
1	E	328/349 (94%)	292 (89%)	24 (7%)	12 (4%)	5	0
2	B	34/36 (94%)	28 (82%)	5 (15%)	1 (3%)	7	1
2	D	34/36 (94%)	28 (82%)	4 (12%)	2 (6%)	2	0
2	F	34/36 (94%)	27 (79%)	6 (18%)	1 (3%)	7	1
All	All	1082/1155 (94%)	990 (92%)	71 (7%)	21 (2%)	12	2

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	LYS
2	B	328	PRO
1	C	204	LYS
2	D	329	GLU
1	E	145	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/313 (95%)	289 (97%)	9 (3%)	53	42
1	C	293/313 (94%)	280 (96%)	13 (4%)	39	25
1	E	297/313 (95%)	289 (97%)	8 (3%)	57	47
2	B	34/34 (100%)	33 (97%)	1 (3%)	55	44
2	D	34/34 (100%)	33 (97%)	1 (3%)	55	44
2	F	34/34 (100%)	33 (97%)	1 (3%)	55	44
All	All	990/1041 (95%)	957 (97%)	33 (3%)	50	37

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	112	ASP
1	C	281	GLN
1	E	244	ASN
1	C	205	GLU
1	C	234	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	151	ASN
1	C	249	ASN
1	E	305	ASN
1	C	173	GLN
1	C	288	ASN

5.3.3 RNA

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/349 (95%)	-0.08	7 (2%) 60 62	17, 30, 50, 81	0
1	C	328/349 (93%)	0.13	7 (2%) 60 62	20, 37, 60, 73	0
1	E	332/349 (95%)	0.28	13 (3%) 37 38	26, 43, 67, 83	0
2	B	36/36 (100%)	0.42	4 (11%) 6 5	21, 36, 66, 74	0
2	D	36/36 (100%)	0.19	3 (8%) 11 11	21, 30, 59, 69	0
2	F	36/36 (100%)	0.45	2 (5%) 24 24	30, 43, 73, 79	0
All	All	1102/1155 (95%)	0.13	36 (3%) 44 45	17, 36, 63, 83	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	139	VAL	8.6
1	A	325	THR	8.2
2	F	350	GLN	6.5
1	E	145	ASN	5.3
1	A	326	ILE	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.