



Full wwPDB X-ray Structure Validation Report (i)

Mar 1, 2014 – 12:43 AM GMT

PDB ID : 4I32
Title : Crystal structure of HCV NS3/4A D168V protease complexed with compound 4
Authors : Lemke, C.T.
Deposited on : 2012-11-23
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance (i)


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3039 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1328	829	238	250	11			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	182	Total	C	N	O	S	0	0	0
			1328	829	238	250	11			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP P26662
A	114	ILE	VAL	CONFLICT	UNP P26662
A	132	VAL	ILE	CONFLICT	UNP P26662
A	168	VAL	ASP	ENGINEERED MUTATION	UNP P26662
A	181	ALA	-	EXPRESSION TAG	UNP P26662
A	183	LYS	-	EXPRESSION TAG	UNP P26662
A	184	LYS	-	EXPRESSION TAG	UNP P26662
A	185	LYS	-	EXPRESSION TAG	UNP P26662
A	186	LYS	-	EXPRESSION TAG	UNP P26662
B	0	MET	-	EXPRESSION TAG	UNP P26662
B	114	ILE	VAL	CONFLICT	UNP P26662
B	132	VAL	ILE	CONFLICT	UNP P26662
B	168	VAL	ASP	ENGINEERED MUTATION	UNP P26662
B	181	ALA	-	EXPRESSION TAG	UNP P26662
B	183	LYS	-	EXPRESSION TAG	UNP P26662
B	184	LYS	-	EXPRESSION TAG	UNP P26662
B	185	LYS	-	EXPRESSION TAG	UNP P26662
B	186	LYS	-	EXPRESSION TAG	UNP P26662

- Molecule 2 is a protein called HCV non-structural protein 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	13	Total	C	N	O		0	0	0
			88	57	16	15				

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	D	13	Total C N O 88 57 16 15	0	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	219	LYS	-	INSERTION	UNP P26662
C	220	LYS	-	INSERTION	UNP P26662
C	235	LYS	-	INSERTION	UNP P26662
D	219	LYS	-	INSERTION	UNP P26662
D	220	LYS	-	INSERTION	UNP P26662
D	235	LYS	-	INSERTION	UNP P26662

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is (2R,6S,7E,10E,13AR,14AR,16AS)-2-{{[7-METHOXY-8-METHYL-2-(PROPYL)-AN-2-YLOXY)QUINOLIN-4-YL]OXY}-N-[(1-METHYLCYCLOPROPYL)SULFONYL]-6-{{[(1-METHYL-1H-PYRAZOL-3-YL)CARBONYL]AMINO}-5,16-DIOXO-1,2,3,6,9,12,13,13A,14,15,16,16A-DODECAHYDROCYCLOPROPA[E]PYRRROLO[1,2-A][1,4]DIAZACYCLOPENTADECINE-14A(5H)-CARBOXAMIDE (three-letter code: 1BV) (formula: C₄₁H₅₁N₇O₉S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			58	41	7	9	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	50	Total	O	0	0
			50	50		
5	B	37	Total	O	0	0
			37	37		
5	C	2	Total	O	0	0
			2	2		
5	D	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Genome polyprotein

Chain A:



- Molecule 1: Genome polyprotein

Chain B:



- Molecule 2: HCV non-structural protein 4A

Chain C:



- Molecule 2: HCV non-structural protein 4A

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	93.67Å 93.67Å 81.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.84 – 2.30 57.61 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.84-2.30) 98.8 (57.61-2.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.12 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R , R_{free}	0.197 , 0.248 0.193 , 0.240	Depositor DCC
R_{free} test set	917 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.6	EDS
Estimated twinning fraction	0.041 for h,-h-k,-l	Xtriage
L-test for twinning	$< L > = 0.51$, $< L^2 > = 0.34$	Xtriage
Outliers	0 of 17994 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3039	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 1BV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.40	0/1355	0.59	0/1848
1	B	0.41	0/1355	0.62	0/1848
2	C	0.38	0/87	0.66	0/116
2	D	0.35	0/87	0.61	0/116
All	All	0.40	0/2884	0.61	0/3928

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1328	0	1336	29	0
1	B	1328	0	1336	28	0
2	C	88	0	100	0	0
2	D	88	0	100	0	0
3	A	1	0	0	0	0
4	A	58	0	51	7	0
4	B	58	0	51	6	0
5	A	50	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	37	0	0	1	0
5	C	2	0	0	0	0
5	D	1	0	0	0	0
All	All	3039	0	2974	61	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (61) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:B:201:1BV:O9	4:B:201:1BV:H9	1.69	0.90
1:A:160:THR:HG22	1:A:161:ARG:HG2	1.64	0.79
1:B:104:LEU:HD22	1:B:151:VAL:HG21	1.67	0.76
4:A:202:1BV:H9	4:A:202:1BV:O9	1.87	0.72
1:B:116:VAL:HG22	1:B:126:LEU:HD23	1.80	0.62
1:B:155:ARG:NH1	4:B:201:1BV:H38	2.15	0.62
1:A:14:LEU:O	1:A:18:ILE:HG12	1.99	0.62
1:A:23:GLY:HA3	1:A:70:PRO:HG3	1.83	0.59
1:A:127:LEU:HD23	1:B:102:SER:HB2	1.84	0.58
1:B:115:PRO:HD2	5:B:321:HOH:O	2.04	0.58
1:A:2:PRO:HG2	1:A:3:ILE:H	1.69	0.58
1:A:1:ALA:HA	1:A:2:PRO:C	2.25	0.56
1:A:127:LEU:HD23	1:B:102:SER:CB	2.36	0.56
1:A:136:LYS:HG2	4:A:202:1BV:H11	1.87	0.55
1:A:75:TYR:CZ	1:A:179:MET:HG2	2.42	0.54
1:B:126:LEU:HD12	1:B:164:ALA:O	2.09	0.53
1:B:51:VAL:HG11	1:B:53:TRP:CE2	2.44	0.53
4:A:202:1BV:H29	5:A:329:HOH:O	2.09	0.53
4:A:202:1BV:N45	4:A:202:1BV:C55	2.72	0.52
1:A:32:GLU:HG3	1:A:92:ARG:HB2	1.92	0.51
1:B:132:VAL:CG2	4:B:201:1BV:H25	2.41	0.50
1:B:132:VAL:HG12	1:B:162:GLY:O	2.11	0.50
1:A:117:ARG:NH2	1:B:119:ARG:HH21	2.09	0.49
1:B:75:TYR:CE1	1:B:179:MET:HA	2.48	0.49
1:A:92:ARG:NH1	5:A:343:HOH:O	2.45	0.49
1:B:32:GLU:N	1:B:32:GLU:OE1	2.43	0.48
4:B:201:1BV:O17	4:B:201:1BV:H26	2.14	0.48
1:B:115:PRO:HB2	1:B:127:LEU:HD12	1.96	0.47
1:A:160:THR:HG22	1:A:161:ARG:CG	2.39	0.47
1:B:47:CYS:HA	1:B:51:VAL:O	2.14	0.47
1:A:160:THR:C	1:A:161:ARG:HG2	2.35	0.47
1:B:136:LYS:HA	4:B:201:1BV:H11	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:80:GLN:NE2	1:A:82:LEU:HD22	2.31	0.46
1:A:2:PRO:CG	1:A:3:ILE:H	2.28	0.46
1:B:132:VAL:HG23	4:B:201:1BV:H25	1.97	0.46
1:A:74:MET:HG2	1:A:75:TYR:CE2	2.51	0.45
1:A:17:ILE:O	1:A:20:SER:HB3	2.16	0.45
1:B:98:THR:HA	1:B:99:CYS:HA	1.67	0.45
1:A:104:LEU:HD22	1:A:151:VAL:HG21	1.99	0.45
1:A:80:GLN:OE1	1:A:174:SER:HB3	2.19	0.43
1:A:80:GLN:HE21	1:A:82:LEU:CD2	2.31	0.43
4:A:202:1BV:H26	4:A:202:1BV:H8	1.91	0.42
1:A:74:MET:HB2	1:A:86:PRO:HD3	1.99	0.42
1:B:24:ARG:O	1:B:24:ARG:NE	2.52	0.42
1:B:175:MET:O	1:B:179:MET:HG3	2.20	0.42
1:B:145:CYS:HB2	1:B:146:PRO:HD2	2.02	0.42
1:B:104:LEU:CD2	1:B:151:VAL:HG21	2.45	0.42
1:A:18:ILE:O	1:A:22:THR:HG23	2.19	0.42
1:B:160:THR:HG22	1:B:161:ARG:HG3	2.01	0.42
1:B:49:ASN:N	1:B:93:SER:OG	2.53	0.42
4:A:202:1BV:N45	4:A:202:1BV:H45	2.35	0.42
1:A:132:VAL:HG23	1:A:162:GLY:C	2.40	0.42
1:B:51:VAL:CG1	1:B:53:TRP:CE2	3.03	0.41
1:A:95:THR:HA	1:A:96:PRO:HD3	1.90	0.41
4:A:202:1BV:H36	4:A:202:1BV:H17	1.76	0.41
1:B:114:ILE:HA	1:B:115:PRO:HD3	1.89	0.41
1:A:114:ILE:HA	1:A:115:PRO:HD3	1.74	0.41
1:A:82:LEU:HD13	1:A:170:ILE:HG21	2.01	0.41
1:B:136:LYS:HD2	1:B:136:LYS:HA	1.84	0.41
1:A:127:LEU:HD22	1:B:102:SER:OG	2.21	0.41
1:A:112:ASP:HB2	1:A:134:TYR:OH	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/187 (96%)	177 (98%)	2 (1%)	1 (1%)	33	39
1	B	180/187 (96%)	179 (99%)	1 (1%)	0	100	100
2	C	11/17 (65%)	11 (100%)	0	0	100	100
2	D	11/17 (65%)	11 (100%)	0	0	100	100
All	All	382/408 (94%)	378 (99%)	3 (1%)	1 (0%)	50	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/153 (95%)	141 (97%)	5 (3%)	49	64
1	B	146/153 (95%)	145 (99%)	1 (1%)	91	97
2	C	10/14 (71%)	9 (90%)	1 (10%)	11	12
2	D	10/14 (71%)	9 (90%)	1 (10%)	11	12
All	All	312/334 (93%)	304 (97%)	8 (3%)	59	76

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	17	ILE
1	A	143	LEU
1	A	161	ARG
1	A	176	GLU
1	B	24	ARG
2	C	228	ARG
2	D	228	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	B	41	GLN

5.3.3 RNA (i)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	1BV	A	202	-	64,64,64	2.18	16 (25%)	97,97,97	3.02	39 (40%)
4	1BV	B	201	-	64,64,64	2.29	21 (32%)	97,97,97	2.78	32 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1BV	A	202	-	-	1/55/89/89	0/1/7/7
4	1BV	B	201	-	-	1/55/89/89	0/1/7/7

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	201	1BV	C8-N7	6.36	1.45	1.38
4	B	201	1BV	C23-N22	6.30	1.47	1.34
4	A	202	1BV	C16-N15	6.07	1.47	1.34
4	A	202	1BV	C8-N7	5.99	1.45	1.38
4	B	201	1BV	C16-N15	5.90	1.47	1.34
4	A	202	1BV	C23-N22	5.68	1.46	1.34
4	A	202	1BV	C21-C20	-4.69	1.44	1.53
4	B	201	1BV	C21-C20	-4.57	1.44	1.53
4	A	202	1BV	C13-C14	-3.98	1.38	1.53
4	B	201	1BV	C13-C14	-3.96	1.38	1.53
4	B	201	1BV	C26-C27	3.91	1.50	1.31
4	A	202	1BV	C32-N31	3.74	1.42	1.34
4	B	201	1BV	C32-N31	3.73	1.42	1.34
4	A	202	1BV	C26-C27	3.69	1.49	1.31
4	B	201	1BV	C44-N45	3.43	1.35	1.30
4	A	202	1BV	C44-N45	3.41	1.35	1.30
4	A	202	1BV	C30-C29	3.37	1.50	1.31
4	B	201	1BV	C30-C29	3.29	1.49	1.31
4	B	201	1BV	C1-C3	-3.23	1.49	1.51
4	B	201	1BV	C2-C3	-3.23	1.49	1.51
4	A	202	1BV	O56-C50	3.04	1.42	1.37
4	B	201	1BV	C25-N31	-2.87	1.43	1.47
4	B	201	1BV	C10-C12	-2.81	1.49	1.52
4	B	201	1BV	C25-C23	-2.70	1.51	1.53
4	A	202	1BV	C1-C3	-2.63	1.49	1.51
4	A	202	1BV	C18-C16	-2.57	1.47	1.52
4	A	202	1BV	C10-C12	-2.48	1.50	1.52
4	B	201	1BV	O52-C44	2.48	1.42	1.36
4	B	201	1BV	C18-C16	-2.43	1.47	1.52
4	B	201	1BV	O56-C50	2.43	1.41	1.37
4	B	201	1BV	C13-C12	-2.42	1.46	1.53
4	A	202	1BV	C13-C12	-2.39	1.46	1.53
4	A	202	1BV	O52-C44	2.39	1.42	1.36
4	B	201	1BV	C25-C26	2.35	1.53	1.50
4	A	202	1BV	C25-N31	-2.32	1.44	1.47
4	B	201	1BV	C46-N45	2.26	1.42	1.37
4	B	201	1BV	O41-C42	2.05	1.43	1.37

All (71) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	202	1BV	C8-N7-S4	-9.60	115.95	124.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	202	1BV	O6-S4-O5	-9.45	101.76	120.51
4	A	202	1BV	C44-O52-C53	9.28	127.31	117.86
4	B	201	1BV	O6-S4-O5	-9.11	102.44	120.51
4	A	202	1BV	O56-C50-C51	9.10	122.78	115.13
4	A	202	1BV	C8-C10-N15	8.18	124.47	117.23
4	B	201	1BV	O6-S4-C3	7.75	113.42	107.50
4	B	201	1BV	O56-C50-C51	7.68	121.59	115.13
4	B	201	1BV	C8-N7-S4	-7.44	117.78	124.12
4	B	201	1BV	C11-C10-N15	-7.26	108.79	117.67
4	B	201	1BV	C44-O52-C53	7.19	125.19	117.86
4	B	201	1BV	C8-C10-N15	7.08	123.49	117.23
4	B	201	1BV	C13-C12-C10	-6.14	111.49	122.47
4	A	202	1BV	C11-C10-N15	-6.03	110.30	117.67
4	A	202	1BV	C44-N45-C46	5.53	121.50	116.24
4	A	202	1BV	O6-S4-C3	5.04	111.35	107.50
4	B	201	1BV	C11-C10-C8	-4.97	108.95	116.62
4	A	202	1BV	O5-S4-C3	4.90	111.24	107.50
4	A	202	1BV	C13-C12-C10	-4.68	114.11	122.47
4	A	202	1BV	O56-C50-C49	-4.39	117.12	124.37
4	B	201	1BV	C44-N45-C46	4.25	120.28	116.24
4	B	201	1BV	O56-C50-C49	-4.24	117.36	124.37
4	A	202	1BV	C39-N36-N35	4.13	125.48	120.56
4	B	201	1BV	C2-C3-S4	-3.98	110.68	116.99
4	A	202	1BV	C2-C3-S4	-3.87	110.85	116.99
4	A	202	1BV	C11-C12-C13	-3.84	112.47	121.29
4	A	202	1BV	C11-C10-C8	-3.80	110.75	116.62
4	A	202	1BV	C38-C34-N35	-3.77	107.42	110.69
4	B	201	1BV	C19-C18-C16	3.69	118.96	111.20
4	B	201	1BV	O41-C42-C47	3.66	120.19	114.52
4	A	202	1BV	O52-C44-N45	3.61	122.60	119.84
4	B	201	1BV	C38-C34-N35	-3.56	107.60	110.69
4	A	202	1BV	C37-N36-N35	-3.55	108.93	111.39
4	B	201	1BV	C39-N36-N35	3.44	124.66	120.56
4	A	202	1BV	C25-C23-N22	3.28	120.65	117.47
4	B	201	1BV	C11-C12-C13	-3.27	113.77	121.29
4	A	202	1BV	C42-O41-C20	-2.97	110.42	120.67
4	A	202	1BV	O9-C8-N7	-2.89	117.80	120.95
4	B	201	1BV	C16-C18-N22	-2.80	105.71	112.71
4	A	202	1BV	C43-C44-N45	-2.71	120.75	124.03
4	B	201	1BV	O17-C16-N15	-2.70	116.59	123.00
4	A	202	1BV	O17-C16-N15	-2.66	116.70	123.00
4	B	201	1BV	C43-C44-N45	-2.62	120.86	124.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	202	1BV	C16-C18-N22	-2.56	106.31	112.71
4	A	202	1BV	C13-C14-C30	2.47	120.13	112.58
4	A	202	1BV	C14-C13-C12	2.44	118.56	114.46
4	A	202	1BV	O52-C53-C55	2.44	116.80	107.97
4	A	202	1BV	O9-C8-C10	2.44	123.54	120.72
4	A	202	1BV	C19-C20-C21	2.43	105.99	103.55
4	A	202	1BV	O24-C23-N22	-2.41	117.54	121.38
4	B	201	1BV	C20-C21-N22	2.38	105.34	102.38
4	A	202	1BV	C40-C3-C2	2.34	122.36	116.72
4	B	201	1BV	O52-C44-N45	2.33	121.62	119.84
4	B	201	1BV	C37-N36-N35	-2.32	109.78	111.39
4	B	201	1BV	C27-C28-C29	2.31	119.67	111.65
4	B	201	1BV	C34-C32-N31	2.31	120.19	114.94
4	B	201	1BV	C42-O41-C20	-2.31	112.71	120.67
4	A	202	1BV	C51-C46-N45	2.25	120.07	117.73
4	B	201	1BV	C40-C3-C2	2.22	122.06	116.72
4	B	201	1BV	C40-C3-C1	2.18	121.96	116.72
4	B	201	1BV	C25-C23-N22	2.17	119.58	117.47
4	B	201	1BV	C25-C26-C27	-2.15	116.15	126.12
4	A	202	1BV	C34-C32-N31	2.14	119.82	114.94
4	A	202	1BV	C1-C3-S4	2.10	120.32	116.99
4	A	202	1BV	O41-C42-C47	2.07	117.73	114.52
4	A	202	1BV	C49-C48-C47	-2.07	118.07	121.15
4	B	201	1BV	C23-C25-N31	-2.07	103.88	107.71
4	B	201	1BV	O41-C20-C19	2.05	114.16	108.75
4	A	202	1BV	C18-C16-N15	2.05	122.20	116.41
4	A	202	1BV	C11-C10-C12	2.04	60.15	59.11
4	A	202	1BV	C12-C10-N15	-2.04	112.27	117.92

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	201	1BV	O33-C32-C34-N35
4	A	202	1BV	O33-C32-C34-N35

There are no ring outliers.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/187 (97%)	0.06	2 (1%) 77 85	41, 53, 72, 92	0
1	B	182/187 (97%)	0.23	8 (4%) 33 43	41, 55, 75, 97	0
2	C	13/17 (76%)	-0.08	0 100 100	43, 47, 61, 73	0
2	D	13/17 (76%)	-0.02	0 100 100	43, 49, 71, 75	0
All	All	390/408 (95%)	0.13	10 (2%) 54 63	41, 54, 74, 97	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	PRO	4.8
1	B	158	VAL	4.7
1	B	129	PRO	3.4
1	B	14	LEU	2.7
1	B	105	TYR	2.7
1	B	180	ARG	2.5
1	B	1	ALA	2.5
1	A	80	GLN	2.2
1	B	179	MET	2.2
1	B	102	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	1BV	B	201	58/58	0.15	0.06	42,55,64,68	0
4	1BV	A	202	58/58	0.13	-0.22	37,51,63,68	0
3	NA	A	201	1/1	0.10	-0.35	63,63,63,63	0

6.5 Other polymers (i)

There are no such residues in this entry.