



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 03:59 AM GMT

PDB ID : 4I3R
Title : Crystal structure of the outer domain of HIV-1 gp120 in complex with VRC-PG04 space group P3221
Authors : Joyce, M.G.; Biertumpfel, C.; Nabel, G.J.; Kwong, P.D.
Deposited on : 2012-11-26
Resolution : 3.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	G	190	
2	H	228	
3	L	208	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4874 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer domain of HIV-1 gp120 (KER2018 OD4.2.2).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	190	Total	C	N	O	S	0	1	0
			1464	900	264	291	9			

- Molecule 2 is a protein called Heavy chain of VRC-PG04 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	228	Total	C	N	O	S	0	0	0
			1732	1097	300	329	6			

- Molecule 3 is a protein called Light chain of VRC-PG04 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	208	Total	C	N	O	S	0	0	0
			1621	1019	275	320	7			

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	3	Total	O	0	0
			3	3		
5	L	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer domain of HIV-1 gp120 (KER2018 OD4.2.2)

Chain G: 



- Molecule 2: Heavy chain of VRC-PG04 Fab

Chain H: 



- Molecule 3: Light chain of VRC-PG04 Fab

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	158.91Å 158.91Å 80.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.65 – 3.00 43.65 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (43.65-3.00) 96.9 (43.65-3.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 3.01Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.1_1168)	Depositor
R, R_{free}	0.183 , 0.245 0.186 , 0.247	Depositor DCC
R_{free} test set	1379 reflections (6.02%)	DCC
Wilson B-factor (Å ²)	81.8	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 71.9	EDS
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 22974 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4874	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.44	0/1489	0.64	0/2020
2	H	0.44	0/1777	0.64	0/2421
3	L	0.42	0/1658	0.60	0/2246
All	All	0.43	0/4924	0.63	0/6687

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1464	0	0	11	0
2	H	1732	0	3	12	0
3	L	1621	0	0	5	0
4	G	53	0	46	6	0
5	G	3	0	0	0	0
5	L	1	0	0	0	0
All	All	4874	0	49	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 6.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:L:213:GLU:N	3:L:213:GLU:OE1	2.02	0.92
4:G:902:NAG:H83	4:G:902:NAG:O3	1.84	0.77
1:G:339:ASN:CG	4:G:903:NAG:C1	2.53	0.77
1:G:301:ASN:OD1	1:G:302:ASN:ND2	2.28	0.65
1:G:368:ASP:OD2	2:H:71:ARG:NH1	2.32	0.62
1:G:412:ASP:N	1:G:412:ASP:OD1	2.34	0.60
2:H:6:GLN:O	2:H:105:ARG:NH2	2.35	0.59
2:H:200:HIS:ND1	2:H:203:SER:OG	2.35	0.59
1:G:334:ASN:OD1	4:G:902:NAG:N2	2.37	0.58
1:G:474:ASP:OD2	1:G:476[B]:ARG:NE	2.38	0.56
1:G:350:ARG:NH2	1:G:395(A):ASN:O	2.42	0.53
2:H:21:SER:OG	2:H:79:HIS:ND1	2.43	0.51
2:H:82(A):ARG:NH1	2:H:82(A):ARG:CG	2.73	0.50
1:G:403:ASP:OD1	1:G:403:ASP:N	2.44	0.50
4:G:902:NAG:C3	4:G:902:NAG:H83	2.41	0.49
2:H:66:ARG:NH2	2:H:86:ASP:OD2	2.45	0.49
2:H:52(A):THR:O	2:H:71:ARG:NE	2.48	0.47
1:G:339:ASN:CB	4:G:903:NAG:C1	2.93	0.47
2:H:65:GLN:OE1	2:H:82(A):ARG:NH2	2.48	0.47
2:H:209:LYS:NZ	3:L:123:GLU:OE2	2.48	0.46
4:G:901:NAG:O7	4:G:901:NAG:O3	2.30	0.45
3:L:82:ASP:O	3:L:86:TYR:OH	2.36	0.44
3:L:12:SER:OG	3:L:105:GLU:OE1	2.35	0.43
1:G:273:LYS:NZ	1:G:287:GLN:OE1	2.52	0.43
2:H:94:ARG:NH1	2:H:101:ASP:OD2	2.50	0.43
2:H:6:GLN:NE2	2:H:90:TYR:O	2.52	0.43
2:H:1:GLN:OE1	2:H:1:GLN:N	2.52	0.42
3:L:65:SER:OG	3:L:66:GLN:N	2.53	0.42
1:G:257:THR:OG1	1:G:258:GLN:N	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	189/190 (100%)	169 (89%)	19 (10%)	1 (0%)	38	84
2	H	226/228 (99%)	221 (98%)	4 (2%)	1 (0%)	43	87
3	L	206/208 (99%)	196 (95%)	10 (5%)	0	100	100
All	All	621/626 (99%)	586 (94%)	33 (5%)	2 (0%)	50	90

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	477	ASP
2	H	126	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	166/165 (101%)	153 (92%)	13 (8%)	18	55
2	H	192/193 (100%)	172 (90%)	20 (10%)	10	37
3	L	182/182 (100%)	161 (88%)	21 (12%)	8	32
All	All	540/540 (100%)	486 (90%)	54 (10%)	11	39

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	278	SER
1	G	335	ARG
1	G	339	ASN
1	G	347	ILE
1	G	357	THR
1	G	390	LEU
1	G	401	SER
1	G	402	ILE
1	G	403	ASP
1	G	419	ARG
1	G	450	THR
1	G	464	THR
1	G	480	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	2	VAL
2	H	4	LEU
2	H	7	SER
2	H	25	SER
2	H	60	SER
2	H	65	GLN
2	H	68	SER
2	H	70	THR
2	H	71	ARG
2	H	99	THR
2	H	113	SER
2	H	135	THR
2	H	159	LEU
2	H	161	SER
2	H	187	SER
2	H	189	LEU
2	H	193	THR
2	H	203	SER
2	H	212	GLU
2	H	216	CYS
3	L	3	VAL
3	L	13	LEU
3	L	14	SER
3	L	18	THR
3	L	21	LEU
3	L	27	SER
3	L	52	SER
3	L	61	ARG
3	L	63	SER
3	L	65	SER
3	L	72	THR
3	L	74	THR
3	L	85	ARG
3	L	104	LEU
3	L	107	ARG
3	L	108	ARG
3	L	133	VAL
3	L	145	LYS
3	L	154	LEU
3	L	203	SER
3	L	210	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	900	1	12,14,15	0.52	0	15,19,21	1.22	1 (6%)
4	NAG	G	901	1	12,14,15	0.46	0	15,19,21	0.83	1 (6%)
4	NAG	G	902	1	12,14,15	0.56	0	15,19,21	1.20	2 (13%)
4	NAG	G	903	1	8,11,15	1.14	1 (12%)	6,14,21	1.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	900	1	-	0/6/23/26	1/1/1/1
4	NAG	G	901	1	-	0/6/23/26	0/1/1/1
4	NAG	G	902	1	-	1/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	903	1	-	2/4/15/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	903	NAG	C5-C4	3.07	1.53	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	900	NAG	O5-C5-C6	3.68	110.84	106.98
4	G	902	NAG	O5-C5-C6	2.94	110.06	106.98
4	G	901	NAG	O5-C5-C6	2.35	109.45	106.98
4	G	902	NAG	O5-C5-C4	-2.10	107.99	110.65

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	903	NAG	O7-C7-N2-C2
4	G	902	NAG	O7-C7-N2-C2
4	G	903	NAG	C8-C7-N2-C2

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	900	NAG	C1-C2-C3-C4-C5-O5

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	190/190 (100%)	-0.13	2 (1%) 77 21	54, 91, 188, 210	0
2	H	228/228 (100%)	-0.21	1 (0%) 90 41	48, 92, 140, 227	0
3	L	208/208 (100%)	-0.16	0 100 100	47, 82, 144, 165	0
All	All	626/626 (100%)	-0.17	3 (0%) 88 36	47, 89, 158, 227	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	481	SER	2.8
1	G	404	GLY	2.5
2	H	1	GLN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	G	901	14/15	0.46	-	117,163,180,182	0
4	NAG	G	902	14/15	0.42	-	138,155,171,174	0
4	NAG	G	900	14/15	0.38	-	153,165,170,170	0
4	NAG	G	903	11/15	0.47	-	129,141,163,164	0

6.5 Other polymers

There are no such residues in this entry.