



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 03:37 PM GMT

PDB ID : 1IB1  
Title : CRYSTAL STRUCTURE OF THE 14-3-3 ZETA:SEROTONIN N-ACETYLTRANSFERASE COMPLEX  
Authors : Obsil, T.; Ghirlando, R.; Klein, D.C.; Ganguly, S.; Dyda, F.  
Deposited on : 2001-03-26  
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	245	
1	B	245	
1	C	245	
1	D	245	
2	E	200	
2	F	200	
2	G	200	
2	H	200	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13407 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 ZETA ISOFORM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1821	1142	306	364	9			
1	B	227	Total	C	N	O	S	0	0	0
			1821	1142	306	364	9			
1	C	227	Total	C	N	O	S	0	0	0
			1821	1142	306	364	9			
1	D	227	Total	C	N	O	S	0	0	0
			1821	1142	306	364	9			


- Molecule 2 is a protein called SEROTONIN N-ACETYLTRANSFERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	E	179	Total 1415	C 897	N 263	O 246	P 1	S 8	0	0	0
2	F	179	Total 1415	C 897	N 263	O 246	P 1	S 8	0	0	0
2	G	179	Total 1415	C 897	N 263	O 246	P 1	S 8	0	0	0
2	H	179	Total 1415	C 897	N 263	O 246	P 1	S 8	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	31	TPO	THR	MODIFIED RESIDUE	UNP Q29495
F	31	TPO	THR	MODIFIED RESIDUE	UNP Q29495
G	31	TPO	THR	MODIFIED RESIDUE	UNP Q29495
H	31	TPO	THR	MODIFIED RESIDUE	UNP Q29495

- Molecule 3 is COA-S-ACETYL TRYPTAMINE (three-letter code: COT) (formula: C<sub>33</sub>H<sub>48</sub>N<sub>9</sub>O<sub>17</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	F	1	Total 63	C 33	N 9	O 17	P 3	S 1	0	0
3	E	1	Total 63	C 33	N 9	O 17	P 3	S 1	0	0
3	G	1	Total 63	C 33	N 9	O 17	P 3	S 1	0	0
3	H	1	Total 63	C 33	N 9	O 17	P 3	S 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	28	Total	O	0	0
			28	28		
4	B	20	Total	O	0	0
			20	20		
4	C	42	Total	O	0	0
			42	42		
4	D	31	Total	O	0	0
			31	31		
4	E	24	Total	O	0	0
			24	24		
4	F	21	Total	O	0	0
			21	21		
4	G	23	Total	O	0	0
			23	23		
4	H	22	Total	O	0	0
			22	22		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: 14-3-3 ZETA ISOFORM

Chain A:




#### • Molecule 1: 14-3-3 ZETA ISOFORM

Chain B:




#### • Molecule 1: 14-3-3 ZETA ISOFORM

Chain C:




#### • Molecule 1: 14-3-3 ZETA ISOFORM

Chain D:




• Molecule 2: SEROTONIN N-ACETYLTRANSFERASE

Chain E:




• Molecule 2: SEROTONIN N-ACETYLTRANSFERASE

Chain F:




• Molecule 2: SEROTONIN N-ACETYLTRANSFERASE

Chain G:



• Molecule 2: SEROTONIN N-ACETYLTRANSFERASE

Chain H:



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.72Å 75.08Å 101.78Å 90.14° 90.06° 63.04°	Depositor
Resolution (Å)	20.00 – 2.70	Depositor
% Data completeness (in resolution range)	99.3 (20.00-2.70)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.204 , 0.228	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13407	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, COT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/1846	0.67	0/2482
1	B	0.42	0/1846	0.62	0/2482
1	C	0.42	0/1846	0.64	0/2482
1	D	0.41	0/1846	0.65	0/2482
2	E	0.44	0/1443	0.69	1/1956 (0.1%)
2	F	0.46	0/1443	0.71	1/1956 (0.1%)
2	G	0.44	0/1443	0.68	1/1956 (0.1%)
2	H	0.45	0/1443	0.68	0/1956
All	All	0.43	0/13156	0.67	3/17752 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	19	GLY	N-CA-C	5.19	126.06	113.10
2	F	19	GLY	N-CA-C	5.18	126.05	113.10
2	E	19	GLY	N-CA-C	5.16	125.99	113.10

There are no chirality outliers.



All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	125	TYR	Sidechain
1	C	125	TYR	Sidechain
1	D	125	TYR	Sidechain

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1821	0	1809	67	0
1	B	1821	0	1809	40	1
1	C	1821	0	1809	48	0
1	D	1821	0	1809	49	0
2	E	1415	0	1382	32	0
2	F	1415	0	1382	43	1
2	G	1415	0	1382	44	0
2	H	1415	0	1382	34	0
3	E	63	0	44	1	0
3	F	63	0	44	1	0
3	G	63	0	44	1	0
3	H	63	0	44	0	0
4	A	28	0	0	19	0
4	B	20	0	0	5	0
4	C	42	0	0	7	1
4	D	31	0	0	13	1
4	E	24	0	0	2	0
4	F	21	0	0	6	0
4	G	23	0	0	10	0
4	H	22	0	0	2	0
All	All	13407	0	12940	317	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

The worst 5 of 317 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:80:ARG:HD2	4:D:3635:HOH:O	1.34	1.22
1:D:144:GLN:HG3	4:D:4412:HOH:O	1.37	1.20
1:A:73:GLU:HB2	4:A:2487:HOH:O	1.57	1.03
1:A:75:LYS:HE3	1:B:8:GLN:HG2	1.48	0.94
1:A:226:THR:CB	4:A:3511:HOH:O	2.24	0.86

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:F:177:GLY:N	4:C:2713:HOH:O[1_554]	2.05	0.15
1:B:31:GLU:O	4:D:4373:HOH:O[1_464]	2.09	0.11

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/245 (92%)	216 (96%)	8 (4%)	1 (0%)	43	76
1	B	225/245 (92%)	215 (96%)	8 (4%)	2 (1%)	25	55
1	C	225/245 (92%)	218 (97%)	6 (3%)	1 (0%)	43	76
1	D	225/245 (92%)	216 (96%)	9 (4%)	0	100	100
2	E	176/200 (88%)	167 (95%)	7 (4%)	2 (1%)	21	49
2	F	176/200 (88%)	169 (96%)	5 (3%)	2 (1%)	21	49
2	G	176/200 (88%)	169 (96%)	5 (3%)	2 (1%)	21	49
2	H	176/200 (88%)	168 (96%)	6 (3%)	2 (1%)	21	49
All	All	1604/1780 (90%)	1538 (96%)	54 (3%)	12 (1%)	30	62

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	72	ALA
1	C	134	ALA
1	A	75	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	E	151	PRO
2	F	151	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/209 (94%)	172 (87%)	25 (13%)	6	15
1	B	197/209 (94%)	177 (90%)	20 (10%)	11	24
1	C	197/209 (94%)	174 (88%)	23 (12%)	8	18
1	D	197/209 (94%)	173 (88%)	24 (12%)	7	17
2	E	148/166 (89%)	139 (94%)	9 (6%)	26	54
2	F	148/166 (89%)	140 (95%)	8 (5%)	31	61
2	G	148/166 (89%)	140 (95%)	8 (5%)	31	61
2	H	148/166 (89%)	138 (93%)	10 (7%)	22	48
All	All	1380/1500 (92%)	1253 (91%)	127 (9%)	13	29

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	119	LEU
1	D	68	LYS
2	H	27	GLN
1	C	156	SER
1	C	191	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	219	GLN
2	E	62	ASN
2	H	62	ASN
1	D	108	ASN
1	B	77	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TPO	E	31	2	10,10,11	5.88	4 (40%)	12,14,16	2.36	2 (16%)
2	TPO	F	31	2	10,10,11	5.74	2 (20%)	12,14,16	2.31	3 (25%)
2	TPO	G	31	2	10,10,11	5.53	3 (30%)	12,14,16	2.27	2 (16%)
2	TPO	H	31	2	10,10,11	5.86	3 (30%)	12,14,16	2.35	3 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	E	31	2	-	0/9/11/13	0/0/0/0
2	TPO	F	31	2	-	0/9/11/13	0/0/0/0
2	TPO	G	31	2	-	0/9/11/13	0/0/0/0
2	TPO	H	31	2	-	0/9/11/13	0/0/0/0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	31	TPO	O-C	18.01	1.23	1.11
2	H	31	TPO	O-C	17.89	1.23	1.11
2	F	31	TPO	O-C	17.75	1.23	1.11
2	G	31	TPO	O-C	16.96	1.23	1.11
2	H	31	TPO	CA-C	2.66	1.53	1.48

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	31	TPO	C-CA-N	-6.05	102.11	111.94
2	G	31	TPO	C-CA-N	-5.96	102.25	111.94
2	E	31	TPO	C-CA-N	-5.95	102.26	111.94
2	F	31	TPO	C-CA-N	-5.94	102.28	111.94
2	E	31	TPO	CB-CA-N	4.61	116.05	109.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	COT	E	401	-	67,67,67	0.80	1 (1%)	98,98,98	1.09	3 (3%)
3	COT	F	400	-	67,67,67	0.81	0	98,98,98	1.07	1 (1%)
3	COT	G	402	-	67,67,67	0.81	1 (1%)	98,98,98	1.05	3 (3%)
3	COT	H	403	-	67,67,67	0.77	0	98,98,98	1.07	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COT	E	401	-	-	0/58/74/74	0/1/5/5
3	COT	F	400	-	-	0/58/74/74	0/1/5/5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COT	G	402	-	-	0/58/74/74	0/1/5/5
3	COT	H	403	-	-	0/58/74/74	0/1/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	401	COT	O4B-C1B	2.12	1.44	1.41
3	G	402	COT	O4B-C1B	2.00	1.44	1.41

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	403	COT	N3A-C2A-N1A	-6.42	123.34	128.71
3	F	400	COT	N3A-C2A-N1A	-6.33	123.41	128.71
3	E	401	COT	N3A-C2A-N1A	-6.20	123.52	128.71
3	G	402	COT	N3A-C2A-N1A	-5.97	123.72	128.71
3	E	401	COT	C4B-O4B-C1B	-2.47	107.07	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.