



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 03:33 AM GMT

PDB ID : 2IEW
Title : Crystal structure of Inositol Phosphate Multikinase Ipk2 from *S. cerevisiae*
Authors : Holmes, W.; Jogl, G.
Deposited on : 2006-09-19
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	363	
1	B	363	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4676 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol polyphosphate multikinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			2113	1356	344	403	10			
1	B	256	Total	C	N	O	S	0	0	0
			2091	1342	336	403	10			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	LEU	-	CLONING ARTIFACT	UNP P07250
A	357	GLU	-	CLONING ARTIFACT	UNP P07250
A	358	HIS	-	EXPRESSION TAG	UNP P07250
A	359	HIS	-	EXPRESSION TAG	UNP P07250
A	360	HIS	-	EXPRESSION TAG	UNP P07250
A	361	HIS	-	EXPRESSION TAG	UNP P07250
A	362	HIS	-	EXPRESSION TAG	UNP P07250
A	363	HIS	-	EXPRESSION TAG	UNP P07250
B	356	LEU	-	CLONING ARTIFACT	UNP P07250
B	357	GLU	-	CLONING ARTIFACT	UNP P07250
B	358	HIS	-	EXPRESSION TAG	UNP P07250
B	359	HIS	-	EXPRESSION TAG	UNP P07250
B	360	HIS	-	EXPRESSION TAG	UNP P07250
B	361	HIS	-	EXPRESSION TAG	UNP P07250
B	362	HIS	-	EXPRESSION TAG	UNP P07250
B	363	HIS	-	EXPRESSION TAG	UNP P07250

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	273	Total 273	O 273	0	0
3	B	198	Total 198	O 198	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Inositol polyphosphate multikinase

Chain A:



- Molecule 1: Inositol polyphosphate multikinase

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	186.51Å 186.51Å 50.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.01 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.8 (30.00-2.00) 93.9 (29.01-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.208 , 0.253 0.210 , 0.251	Depositor DCC
R_{free} test set	4868 reflections (8.25%)	DCC
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.7	EDS
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 66965 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4676	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	2/2155 (0.1%)	0.67	0/2899
1	B	0.51	0/2130	0.62	0/2865
All	All	0.56	2/4285 (0.0%)	0.65	0/5764

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	216	GLU	CB-CG	7.49	1.66	1.52
1	A	216	GLU	CD-OE1	6.69	1.33	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2113	0	2090	29	0
1	B	2091	0	2073	31	0
2	A	1	0	0	0	0
3	A	273	0	0	9	0
3	B	198	0	0	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4676	0	4163	59	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (59) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:25:ASP:HB2	3:B:559:HOH:O	1.25	1.30
1:A:62:LEU:HD13	1:A:247:MET:HE1	1.62	0.79
1:A:58:GLY:N	3:A:735:HOH:O	2.16	0.78
1:B:330:GLU:HG2	3:B:489:HOH:O	1.91	0.71
1:A:317:SER:N	3:A:674:HOH:O	2.23	0.71
1:A:174:LYS:HG2	1:A:194:ASP:O	1.92	0.70
1:A:62:LEU:HD13	1:A:247:MET:CE	2.21	0.69
1:A:209:GLN:HG2	3:A:600:HOH:O	1.93	0.69
1:B:268:GLU:HG3	3:B:396:HOH:O	1.93	0.68
1:A:278:LYS:HE2	1:A:282:ASP:OD2	1.95	0.65
1:B:317:SER:N	3:B:554:HOH:O	2.33	0.61
1:B:174:LYS:HD2	1:B:179:LEU:HD11	1.84	0.59
1:A:45:GLN:HE22	1:A:69:TYR:H	1.50	0.58
1:B:134:LEU:HD11	1:B:259:LEU:HG	1.86	0.56
1:A:209:GLN:CG	3:A:632:HOH:O	2.55	0.54
1:B:44:ILE:HG22	3:B:382:HOH:O	2.10	0.52
1:A:185:GLU:OE2	3:A:712:HOH:O	2.19	0.51
1:B:112:LYS:HG3	1:B:113:GLN:N	2.26	0.51
1:B:25:ASP:N	1:B:25:ASP:OD1	2.43	0.51
1:B:285:ILE:HG12	1:B:286:ASP:H	1.76	0.49
1:A:209:GLN:HG3	3:A:632:HOH:O	2.11	0.49
1:A:147:LYS:HB3	3:A:656:HOH:O	2.12	0.49
1:A:134:LEU:HD11	1:A:259:LEU:HG	1.94	0.48
1:A:45:GLN:NE2	1:A:69:TYR:H	2.11	0.48
1:B:62:LEU:HD13	1:B:247:MET:CE	2.43	0.48
1:B:180:ASN:HB2	3:B:432:HOH:O	2.13	0.48
1:A:145:LEU:HA	1:A:148:ARG:HB3	1.95	0.47
1:B:44:ILE:CG2	3:B:382:HOH:O	2.62	0.47
1:A:118:GLU:HG3	1:A:284:PHE:CD1	2.50	0.47
1:B:68:THR:HB	1:B:118:GLU:HB3	1.97	0.46
1:B:62:LEU:HD13	1:B:247:MET:HE3	1.97	0.46
1:B:58:GLY:N	3:B:494:HOH:O	2.48	0.46
1:A:209:GLN:HG2	3:A:632:HOH:O	2.15	0.46
1:A:339:GLU:O	1:A:343:GLU:HG3	2.17	0.45
1:B:66:MET:HG2	1:B:326:PHE:CZ	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:75:GLU:HB3	1:A:112:LYS:H	1.82	0.44
1:B:324:ILE:O	1:B:325:ASP:HB2	2.18	0.44
1:B:215:ILE:O	1:B:218:TYR:HB3	2.18	0.44
1:A:238:LYS:HZ3	1:B:59:ASP:CG	2.21	0.44
1:A:132:ILE:O	1:A:258:SER:HA	2.18	0.43
1:A:62:LEU:CD1	1:A:247:MET:CE	2.93	0.43
1:A:332:THR:HB	1:A:335:LYS:HD2	2.00	0.43
1:B:281:ARG:NH1	3:B:409:HOH:O	2.52	0.43
1:B:246:THR:HG22	1:B:247:MET:HE2	2.00	0.43
1:B:167:ILE:O	1:B:203:GLY:HA3	2.19	0.42
1:A:145:LEU:CD2	1:A:148:ARG:HD2	2.50	0.42
1:B:251:GLU:O	1:B:331:ILE:HD12	2.20	0.42
1:A:44:ILE:HG22	1:A:63:CYS:HB3	2.02	0.42
1:B:285:ILE:HG12	1:B:286:ASP:N	2.34	0.41
1:B:261:PHE:N	1:B:261:PHE:CD2	2.88	0.41
1:B:25:ASP:CB	3:B:559:HOH:O	2.13	0.41
1:B:354:LYS:HG2	3:B:442:HOH:O	2.21	0.41
1:B:183:SER:HB3	1:B:185:GLU:OE2	2.21	0.41
1:B:73:LEU:HB3	1:B:114:TYR:HB2	2.03	0.41
1:B:228:ARG:HG2	1:B:270:TRP:CE3	2.56	0.41
1:A:167:ILE:O	1:A:203:GLY:HA3	2.22	0.40
1:A:32:PRO:HB3	1:A:112:LYS:NZ	2.37	0.40
1:A:175:ASN:HB2	1:A:222:PRO:HD3	2.03	0.40
1:A:31:LYS:HE3	3:A:722:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/363 (69%)	248 (99%)	2 (1%)	0	100	100
1	B	248/363 (68%)	242 (98%)	6 (2%)	0	100	100
All	All	498/726 (69%)	490 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/327 (72%)	230 (97%)	7 (3%)	53	50
1	B	235/327 (72%)	221 (94%)	14 (6%)	27	20
All	All	472/654 (72%)	451 (96%)	21 (4%)	39	32

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ILE
1	A	141	SER
1	A	145	LEU
1	A	147	LYS
1	A	192	ASP
1	A	285	ILE
1	A	342	ILE
1	B	25	ASP
1	B	29	ILE
1	B	31	LYS
1	B	44	ILE
1	B	45	GLN
1	B	113	GLN
1	B	118	GLU
1	B	147	LYS
1	B	149	GLU
1	B	156	GLU
1	B	185	GLU
1	B	204	ARG
1	B	216	GLU
1	B	331	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	209	GLN
1	A	223	HIS
1	B	45	GLN
1	B	223	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/363 (71%)	0.02	14 (5%) 25 24	15, 24, 50, 72	0
1	B	256/363 (70%)	0.51	27 (10%) 7 6	20, 35, 64, 87	0
All	All	514/726 (70%)	0.27	41 (7%) 12 12	15, 30, 61, 87	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	285	ILE	12.4
1	B	286	ASP	9.9
1	B	191	ALA	9.4
1	A	191	ALA	8.5
1	B	145	LEU	7.9
1	B	284	PHE	6.9
1	B	111	SER	6.8
1	A	286	ASP	6.1
1	B	283	ASP	5.3
1	A	285	ILE	5.2
1	B	282	ASP	4.4
1	A	111	SER	4.2
1	B	184	LEU	4.1
1	B	75	GLU	4.0
1	B	192	ASP	3.9
1	B	146	GLU	3.7
1	A	192	ASP	3.7
1	B	156	GLU	3.7
1	A	112	LYS	3.5
1	A	145	LEU	3.4
1	B	113	GLN	3.3
1	B	34	PHE	3.0
1	B	74	ASN	2.9
1	B	274	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	197	PHE	2.7
1	A	190	GLU	2.7
1	A	282	ASP	2.7
1	B	268	GLU	2.6
1	A	283	ASP	2.6
1	B	144	SER	2.6
1	B	190	GLU	2.6
1	B	45	GLN	2.5
1	B	150	ARG	2.4
1	A	75	GLU	2.3
1	A	146	GLU	2.3
1	A	360	HIS	2.3
1	B	272	LEU	2.3
1	B	180	ASN	2.2
1	B	193	SER	2.2
1	A	284	PHE	2.1
1	B	177	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	A	501	1/1	0.04	-2.36	31,31,31,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.